

HEWLETT-PACKARD

Step-by-Step Solutions
For Your HP Calculator
Calculus

$$\frac{b^2 - 4ac}{2a} = P$$
$$\frac{-2.1}{3} = \int_{t_1}^{\infty} \text{Re}(G(t)) dt + i \int_{t_1}^{\infty} \text{Im}(G(t)) dt$$
$$3 = Pr \left[\frac{2 - 2.151}{1.085} < \frac{X - \mu}{\sigma} \leq \frac{3 - 2.151}{1.085} \right]$$
$$\frac{e^{iz}}{z + 1/z} dz$$
$$s = \frac{f(x + \Delta) - f(x - \Delta)}{2\Delta}$$
$$z^n = r^n e^{in\theta}$$
$$z_c = -30i$$
$$P = P\left(\frac{3 - 2.151}{1.085}\right) - P\left(\frac{2 - 2.151}{1.085}\right)$$
$$J_1(x) =$$
$$\int_{t_2}^{t_1} G(t) dt$$
$$\mathbf{B} = \begin{bmatrix} 3 & 3 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ \frac{1}{3} & 1 \end{bmatrix} \begin{bmatrix} 3 & 3 \\ 0 & 0 \end{bmatrix}$$
$$z^n = r^n e^{in\theta}$$
$$G(t) dt$$

HP-28S
HP-28C

 HEWLETT
PACKARD

Help Us Help You!

Please take a moment to complete this postage-paid card, tear it out and put it in the mail. Your responses and comments will help us better understand your needs and will provide you with the best procedures to solve your problems. Thank you!

HELP US HELP YOU!

Book: **Calculus** Date acquired: _____

Name _____

Street _____

City, State, Zip _____

Phone (_____) _____ Business _____ or Home _____

1. What calculator will you use this book with?
004 **HP-28S** 005 **HP-28C** 006 **Other** _____
2. How many other HP solution books have you bought for this calculator? _____
3. What is your OCCUPATION?
101 **Student** 103 **Professional** 109 **Other** _____
4. Where did you purchase this book?
403 **Bookstore** 404 **Discount or Catalog Store**
407 **Mail Order** 410 **HP Direct** 411 **Other** _____
5. How did you first hear about this book?
501 **HP Owner** 503 **Advertising** 506 **Salesperson** 507 **Brochure**
508 **Other** _____
6. To what degree did this book influence your calculator purchase decision?
601 **Major Influence** 602 **Minor Influence** 603 **No Influence**
7. How well does this book cover the material you expected?
701 **Good** 702 **Moderate** 703 **Low**
8. What level of knowledge is required to make use of the topics in this book?
801 **High** 802 **Medium** 803 **Low**
9. How clearly was the material in this book presented?
901 **Good** 902 **Moderate** 903 **Low**
10. How would you rate the value of this book for your money?
111 **High** 112 **Medium** 113 **Low**

Comments: (Please comment on improvements and additional applications or subjects you would like HP to cover in this or another solution book.) _____



NO POSTAGE
NECESSARY
IF MAILED
IN THE
UNITED STATES

BUSINESS REPLY CARD

FIRST CLASS PERMIT NO. 40 CORVALLIS, OR

POSTAGE WILL BE PAID BY ADDRESSEE

Hewlett-Packard Company
ATTN: Calculator Market Research
1000 N.E. Circle Blvd.
Corvallis, OR 97330, U.S.A.

Calculus

**Step-by-Step Solutions
for Your HP-28S or HP-28C Calculator**



**Edition 2 June 1988
Reorder Number 00028-90102**

Notice

This book and any keystroke programs contained herein are provided "as is" and are subject to change without notice. **Hewlett-Packard Company makes no warranty of any kind with regard to this book or the keystroke programs contained herein, including, but not limited to, the implied warranties of merchantability and fitness for a particular purpose.**

Hewlett-Packard Company shall not be liable for any errors or for incidental or consequential damages in connection with the furnishing, performance, or use of this book or the keystroke programs contained herein.

© Hewlett-Packard Company 1987. All rights reserved. Reproduction, adaptation, or translation of this book, including any programs, is prohibited without prior written permission of Hewlett-Packard Company; except as allowed under the copyright laws. Hewlett-Packard Company grants you the right to use any program contained in this book in a Hewlett-Packard calculator.

The programs that control your calculator are copyrighted and all rights are reserved. Reproduction, adaptation, or translation of those programs without prior written permission of Hewlett-Packard Company is also prohibited.

**Corvallis Division
1000 N.E. Circle Blvd.
Corvallis, OR 97330, U.S.A.**

Printing History

Edition 1	November 1987	Mfg No. 00028-90107
Edition 2	June 1988	Mfg No. 00028-90134

Welcome...

... to the HP-28S and HP-28C Step-by-Step Books. These books are designed to help you get the most from your HP-28S or HP-28C calculator.

This book, *Calculus*, provides examples and techniques for solving problems on your HP-28S or HP-28C. A variety of function operations and differential and integral calculus problems are designed to familiarize you with the many functions built into your calculator.

Before you try the examples in this book, you should be familiar with certain concepts from the owner's documentation:

- The basics of your calculator: how to move from menu to menu, how to exit graphics and edit modes, and how to use the menu to assign values to, and solve for, user variables.
- Entering numbers, programs, and algebraic expressions into the calculator.

Please review the section "How To Use This Book." It contains important information on the examples in this book.

For more information about the topics in the *Calculus* book, refer to a basic textbook on the subject. Many references are available in university libraries and in technical and college bookstores. The examples in the book demonstrate approaches to solving certain problems, but they do not cover the many ways to approach solutions to mathematical problems.

Our thanks to Ross Greenley of Oregon State University for developing the problems in this book.

Contents

7 How To Use This Book

- 11 Function Operations**
- 12** Function Definition
- 14** Function Composition
- 16** Function Analysis
- 22** Angle Between Two Lines
- 23** Angle Between Two Curves

27 Differential Calculus

- 28** Minimize Perimeter
- 31** Mimimize Surface Area
- 35** Lines Tangent to a Circle
- 43** Implicit Differentiation With User-Defined Derivative
- 45** Taylor Series Error Term
- 51** Tangent Lines and Taylor Series
- 54** Normal Line
- 57** Implicit Functions

61 Integral Calculus

- 62** Integration and Free Falling Body
- 69** Double Integration
- 72** Area Between Two Curves
- 76** Arc Length
- 80** Surface Area
- 82** Arc Length of Parametric Equations
- 84** Surface Area of Parametric Equations
- 86** Volume of Solid of Revolution: Method of Shells
- 89** Volume of Solids of Revolution: Method of Disks.

How To Use This Book

Please take a moment to familiarize yourself with the formats used in this book.

Keys and Menu Selection: A box represents a key on the calculator keyboard.

ENTER
1/x
STO

ARRAY
PLOT
ALGEBRA

In many cases, a box represents a shifted key on the calculator. In the example problems, the shift key is NOT explicitly shown. (For example, **ARRAY** requires the press of the shift key, followed by the **ARRAY** key, found above the "A" on the left keyboard.)

The "inverse" highlight represents a menu label:

Key:

DRAW
ISOL
ABCD

Description:

Found in the **PLOT** menu.

Found in the **SOLV** menu.

A user-created name. If you created a variable by this name, it could be found in either the **USER** menu or the **SOLVR** menu. If you created a program by this name, it would be found in the **USER** menu.

Menus typically include more menu labels than can be displayed above the six redefinable menu keys. Press **NEXT** and **PREV** to roll through the menu options. For simplicity, **NEXT** and **PREV** are NOT shown in the examples.

To solve for a user variable within **SOLVR**, press the shift key, followed by the appropriate user-defined menu key:

ABCD.

The keys above indicate the shift key, followed by the user-defined key labeled "ABCD". Pressing these keys initiates the Solver function to seek a solution for "ABCD" in a specified equation.

The symbol **<>** indicates the cursor-menu key.

Interactive Plots and the Graphics Cursor: Coordinate values you obtain from plots using the **INS** and **DEL** digitizing keys may differ from those shown, due to small differences in the positions of the graphics cursor. The values you obtain should be satisfactory for the Solver root-finding that follows.

Display Formats and Numeric Input: Negative numbers, displayed as

-5
-12345.678
[[-1, -2, -3 [-4, -5, -6 [...

are created using the **CHS** key.

5 **CHS**
12345.678 **CHS**
[[1 **CHS**, 2 **CHS**, ...

The examples in this book typically specify a display format for the number of decimal places. If your display is set such that numeric displays do not match exactly, you can modify your display format with the **MODE** menu and the **FIX** key within that menu. For example, to set the calculator to display two decimal places, set it to FIX2 mode by pressing **MODE** 2 **FIX**.

Programming Reminders: Before you key in the programming examples in this book, familiarize yourself with the locations of programming commands that appear as menu labels. By using the menu labels to enter commands you can speed keying in programs and avoid errors that might arise from extra spaces appearing in the programs. Remember, the calculator recognizes commands that are set off by spaces. Therefore, the arrow (→) in the command R→C (the real to complex conversion function) is interpreted differently than the arrow in the command → C (create the local variable "C").


The HP-28S automatically inserts spaces around each operator as you key it in. Therefore, using the **R**, **→**, and **C** keys to enter the R→C command will result in the expression R → C, and, ultimately in an error in your program. As you key in programs on the HP-28S, take particular care to avoid spaces inside commands, especially in commands that include an →.

The HP-28C does not automatically insert spaces around operators or commands as they are keyed in.


A Note About the Displays Used in This Book: The menus and screens that appear in this book show the HP-28S display. Most of the HP-28C and HP-28S screens are identical, but there are differences in the **MODE** menu and **SOLVR** screen that HP-28C users should be aware of.

For example, the first screen below illustrates the HP-28C **MODE** menu, and the second screen illustrates the same menu as it appears on the HP-28S.

HP-28C **MODE** display.




HP-28S **MODE** display.




Notice that the HP-28C highlights the entire active menu item, while the HP-28S display includes a small box in the active menu item.

The screens shown below illustrate the HP-28C and HP-28S versions of the  menu.

HP-28C  display.



HP-28S  display.



Both of these screens include the Solver variables , , , , and . The HP-28C displays Solver variables in gray on a black background. The HP-28S prints Solver variables in black on a gray background.

User Menus: A **PURGE** command follows most of the examples in this book. If you do not purge all of the programs and variables after working each example, or if your **USER** menu contains your own user-defined variables or programs, the **USER** menu on your calculator may differ from the displays shown in this book. Do not be concerned if the variables and programs appear in a slightly different order on your **USER** menu; this will not affect the calculator's performance.

Function Operations

The primary goals of this chapter are to write user-defined functions and introduce the root finding, plotting, and calculus capabilities of the HP-28S and HP-28C. Problems include definition and assignment of the trigonometric co-functions in the USER menu, analysis of a cubic equation, and both specific and general cases of computation of the angle between two intersecting lines.

Function Definition

This section demonstrates creation of simple user-defined functions. The use of functions of this type is basic to efficient use of the HP-28S.

Example: The HP-28S and HP-28C have three basic trigonometric functions built in – sine, cosine, and tangent. It is simple to add the remaining co-functions to the USER menu. Built-in functions of the calculator can be easily combined to create new functions. The use of programs and local variables permits the newly defined functions to be used in the same manner as the built-in functions.

The inverse of the sine is the cosecant.

```
CLEAR  <>  
↔ x '1÷SIN(x)  
ENTER
```

```
4:  
3:  
2:  
1:   <→ x '1/SIN(x)' »
```

Store the user-defined function.

```
'CSC  STO
```

```
4:  
3:  
2:  
1:
```

The inverse of the cosine is the secant.

```
↔ x '1÷COS(x)  
ENTER
```

```
4:  
3:  
2:  
1:   <→ x '1/COS(x)' »
```

Store the user-defined function.

```
'SEC  STO
```

```
4:  
3:  
2:  
1:
```

The inverse of the tangent is the cotangent.

```
↔ x '1÷TAN(x)  
ENTER
```

```
4:  
3:  
2:  
1:   <→ x '1/TAN(x)' »
```

Store the user-defined function.

' COT [STO]

4:
3:
2:
1:

Example: Evaluate, in radians, $\text{COT}(X)$ and $\text{CSC}^2(X) - \text{COT}^2(X)$, where $X = .2$.

First, store the value of X and select radians and standard display modes.

.2 [ENTER]
' X [STO]
MODE RAD STD

3:
2:
1:

STD FIX SCI ENG DEG RAD

Now enter the expression for $\text{COT}(X)$ and evaluate it.

' COT (X [ENTER]
[EVAL]

3:
2:
1:

4.93315487558
STD FIX SCI ENG DEG RAD

Enter the second expression and evaluate it.

' SQ(CSC(X))-SQ(COT(X))
[ENTER] [EVAL]

3:
2:
1:

4.93315487558
1
STD FIX SCI ENG DEG RAD

As expected, this identity returns the value 1.

Purge the variable X created in this section. You may also purge the user-defined functions if you wish.

' X' [PURGE]

Function Composition

This section demonstrates additional utility of user-defined functions. Arguments of the functions may be both numeric and symbolic.

Example: Form the compositions $F(G(x))$ and $G(F(x))$ given

$$F(x) = x^2 + 1 \text{ and } G(x) = 5x + 2.$$

Create F and G as user-defined functions.

First, create F .

CLEAR **<>**
 $\leftarrow x 'x^2+1$ **ENTER**

4:
3:
2:
1: $\leftarrow x 'x^2+1'$ *

Store in the variable F .

'F **STO**

4:
3:
2:
1:

Now create G .

$\leftarrow x '5 \times x+2$ **ENTER**

4:
3:
2:
1: $\leftarrow x '5*x+2'$ *

Store in the variable G .

'G **STO**

4:
3:
2:
1:

To form the composition $G(F(x))$, enter F as an argument of G .

'G(F(X **ENTER**

4:
3:
2:
1: $'G(F(X))'$

Evaluate the composite function.

EVAL

4:
3:
2:
1: $'5*(X^2+1)+2'$

This expression can be simplified using EXPAN and COLCT.

ALGEBRA EXPAN

```
3:  
2:  
1:      '5*X^2+5*X+2'  
COLCT EXPAN SIZE FORM DESUB ENSUB
```

COLCT

```
3:  
2:  
1:      '7+5*X^2'  
COLCT EXPAN SIZE FORM DESUB ENSUB
```

Repeat the process using G as an argument of F .

'F(G(X)) ENTER

```
3:  
2:      '7+5*X^2'  
1:      'F(G(X))'  
COLCT EXPAN SIZE FORM DESUB ENSUB
```

Evaluate the composite function.

EVAL

```
3:  
2:      '7+5*X^2'  
1:      '(5*X+2)^2+1'  
COLCT EXPAN SIZE FORM DESUB ENSUB
```

Simplify the expression.

EXPAN

```
2:      '7+5*X^2'  
1:      '(5*X)^2+2*(5*X)*2+2  
COLCT EXPAN SIZE FORM DESUB ENSUB
```

EXPAN

```
2:      '7+5*X^2'  
1:      '5*X*(5*X)+2*(5*X)*2  
COLCT EXPAN SIZE FORM DESUB ENSUB
```

COLCT

```
3:  
2:      '7+5*X^2'  
1:      '5+25*X^2+20*X'  
COLCT EXPAN SIZE FORM DESUB ENSUB
```

Purge the variables created in this problem section.

{'F','G'} PURGE

Function Analysis

The ability to locate extreme values and other key features of functions is critical to the solution of many problems in science and engineering. This section demonstrates the use of calculus to locate such features.

Example: Locate the roots, local maximum, minimum, and inflection points of

$$F(x) = x^3 + 6x^2 + 11x + 6.$$

Enter and name the given function.

CLEAR **<>**
' $X^3+6 \times X^2+11 \times X+6$ '
ENTER

4:
3:
2:
1: 'X^3+6*X^2+11*X+6'

'**FN**' **STO**

4:
3:
2:
1:

Recall the function, enter the PLOT menu, and store it for plotting.

USER **FN**


3:
2:
1: 'X^3+6*X^2+11*X+6'
FN

PLOT **STEQ**

3:
2:
1:
STEQ RCEG FMIN FMAX INDEF DRAW

Clear the plot parameters and plot the function.

'**PPAR**' **PURGE**
DRAW



Digitize all the roots.

3:	(-9,0)				
2:	(-19,0)				
1:	(-31,0)				
STEP	RCEQ	FMIN	FMAX	INDEX	DRAW

Note: Differences from the displayed results may appear due to slightly different digitizing locations.

Now enter the **SOLVR** menu and compute the three roots.

SOLV

3:	$(-9, 0)$			
2:	$(-1, 9, 0)$			
1:	$(-3, 1, 0)$			
X	EXPR=			

Enter a guess from the stack and compute the root. Remember, to calculate the value of a Solver variable, press the shift key followed by the appropriate variable key.

三 X 三

X: -3
Zero
1: -3

After obtaining the exact root, make note of it and prepare to locate the next root. Discard the first root. Then repeat the process for the other two roots.

DROP

X: -2.111111111111111
Zero
1: -2.000000000006

Compute the last root.

DROP

```
X: -1
Zero
1: -1
x EXPRE=
```

With the three roots located, find the extrema. The extrema are located by finding the roots of the first derivative.

Recall the function

CLEAR **USER** **FN**

```
3:  
2:  
1: 'X^3+6*X^2+11*X+6'  
X FPNB FC FN
```

Purge the current value of X and differentiate with respect to X .

' X ENTER ENTER
PURGE

3:	$'X^3+6*X^2+11*X+6'$				
2:	$'X'$				
1:					
DR1	FFHR	EQ	FN		

d/dx

3:					
2:					
1:	$'3*X^2+6*(2*X)+11'$				
DR1	FFHR	EQ	FN		

Store the first derivative.

' DR1 STO

3:					
2:					
1:					
DR1	FFHR	EQ	FN		

Plot the function and its first derivative.


DR1
FN

3:					
2:	$'3*X^2+6*(2*X)+11'$				
1:	$'X^3+6*X^2+11*X+6'$				
DR1	FFHR	EQ	FN		

= ENTER

2:					
1:	$'3*X^2+6*(2*X)+11=X^3+6*X^2+11*X+6'$				
DR1	FFHR	EQ	FN		

PLOT STEQ DRAW



Observe that the derivative is positive in regions where the function is increasing and negative in regions where the function is decreasing.

Digitize both roots of the derivative.

< ... < INS
< ... < INS
ATTN

3:					
2:					
1:					
STEQ	RCED	PMIN	PMAX	INDEF	DRAW

Note: Differences from the displayed results may appear due to differences in digitizing locations.

Recall the derivative and enter **SOLVR** to pinpoint the roots as above. The computed values may differ slightly depending on the seed provided as an input to the Solver.

USER **DR1**
SOLV **STEQ**
SOLVR

3:	(-1.4, 0)			
2:	(-2.6, 0)			
1:	(-2.57735026917)			
x	EXPRE			

X
 X

X: -2.57735026917				
Zero				
1:	-2.57735026917			
x	EXPRE			

This is one of the roots. Recall the function and evaluate to get the functional value.

USER **FN**
EVAL

3:	(-1.4, 0)			
2:	-2.57735026917			
1:	.38490017949			
x	DR1	FFRR	EQ	FN

Now repeat the process for the other root. First discard the root and function value.

DROP **DROP**
SOLV **SOLVR**

3:	(-1.4, 0)			
2:	(-1.42264973081)			
1:	(-1.42264973081)			
x	EXPRE			

X
 X

X: -1.42264973081				
Zero				
1:	-1.42264973081			
x	EXPRE			

USER **FN**
EVAL

3:	(-1.4, 0)			
2:	-1.42264973081			
1:	-.38490017949			
x	DR1	FFRR	EQ	FN

The extreme values of the function have been located. Clear the stack and find the inflection point. The inflection point, located at the root of the second derivative, is the point or points at which the function changes concavity. That is, it changes from concave up to concave down. The second derivative of a cubic is linear and has only one root. Therefore a cubic has only one point of inflection.

Clear the value of X to obtain symbolic results.

CLEAR
'X PURGE

3:					
2:					
1:					
DR1	PPHR	EQ	FN		

Recall the first derivative.

DR1
'X ENTER

3:					
2:	'3*X^2+6*(2*X)+11'	'X'			
1:					
DR1	PPHR	EQ	FN		

Differentiate it with respect to X .

d/dx

3:					
2:					
1:	'3*(2*X)+12'				
DR1	PPHR	EQ	FN		

Store the second derivative.

'DR2 **STO**

3:					
2:					
1:					
DR2	DR1	PPHR	EQ	FN	

Plot the function and its second derivative. Observe the location of the root and how the function behaves at that point. It is coincidental that a function root is located at the point of inflection. It remains only to repeat the root finding procedure.

DR2
FN

3:					
2:					
1:	'3*(2*X)+12'	'X^3+6*X^2+11*X+6'			
DR2	DR1	PPHR	EQ	FN	


Set them equal for plotting.

= **ENTER**

2:					
1:	'3*(2*X)+12=X^3+6*X^2+11*X+6'				
DR2	DR1	PPHR	EQ	FN	

Store and plot the equation.

PLOT **STEQ**
DRAW



Digitize the root.

. . .

3:
2:
1:
(-2, 1, 0)
STEQ | RCEQ | FMIN | FMAX | INDEF | DRW4

Recall the second derivative and solve for the root.

3:
2:
1: '3*(2*x)+12'
DR2 | DR1 | PPAR | EQ | FN

3:
2:
1:
(-2, 1, 0)
x | EXPRE |

Enter the digitized initial guess and solve for the root.

x = -2
Zero
1: -2
x | EXPRE |

This completes the analysis. We have found roots at $x = -1, -2, -3$, extrema at $x = -2.58, -1.42$, and an inflection point at $x = -2$.

Exit from the Solver menu and purge the user variables created in this section.

Note to HP-28S owners: If you do not exit from before attempting to delete the current equation, the calculator will display the message NO CURRENT EQUATION.

{ 'FN' 'X' 'DR1' 'DR2' 'PPAR'

Angle Between Two Lines

This section develops a user function to compute the angle of intersection of two lines. The slopes of the intersecting lines are supplied as arguments. The user function is used in the subsequent section in computing the angle of intersection of two general functions.

Example: Compute the angle between the lines

$$Y = 3x + 1 \text{ and } Y = -2x + 5.$$

The angle between two curves is the angle formed by the tangent lines at the point of intersection.

$$\theta = \tan^{-1} \frac{m_2 - m_1}{1 + m_1 m_2}.$$

Form a function that, given the slopes, computes the angle between two functions at a point of intersection.

CLEAR MODE DEG
«→ a b 'ATAN((b-a)÷
(1+a*b))' **ENTER**

2:
1: «→ a b 'ATAN((b-a)÷
(1+a*b))' »
STO FIX SCI ENG DEG RND

'ANG **STO**

3:
2:
1:
STO FIX SCI ENG DEG RND

Lines have a constant slope. Read the slope for each directly from the given formula.

3 **ENTER**
-2 **ENTER**

3:
2:
1:
STO FIX SCI ENG DEG RND -2

Now compute the angle.

USER ANG

3:
2:
1:
ANG 45
STO FIX SCI ENG DEG RND

The lines intersect at an angle of 45°.

ANG is used in the next problem section.

Angle Between Two Curves

The angle of intersection for two curves is defined to be the angle formed by the tangent lines at the point of intersection. When an intersection point is located, the slopes of the functions at that point can be found. The problem is then that of two intersecting lines.

Example: Find the angle formed by the tangent lines at the points of intersection of the following functions.

$$F = 3x + 1$$

$$Y = 2x^2$$

Enter and save the given functions.

CLEAR **<>**
'3*X+1 **ENTER**

4:	
3:	
2:	
1:	'3*X+1'

'F **STO**

4:	
3:	
2:	
1:	

'2*X^2 **ENTER**

4:	
3:	
2:	
1:	'2*X^2'

'Y **STO**

4:	
3:	
2:	
1:	

Plot the two functions to obtain initial guesses at the points of intersection.

First, set the two functions equal to each other.

USER **Y** **F**
= **ENTER**

3:				
2:				
1:	'2*X^2=3*X+1'			
Y	F	ANG		


Store the equation.

PLOT **STEQ**

3:
2:
1:
STEQ|RCEQ|FMIN|FMAX|INDEF|DRAW


Clear the plot parameters and draw the equation with the two functions.

'PPAR **PURGE**
DRAW



Expand the height to see both intersection points.

ATTN 10 ***H**
DRAW



Digitize both intersection points. Enter the Solver to refine the guesses.

< ... **<** **INS**
> ... **^** **INS**
ATTN

3:
2:
1:
(-.3,0)
(1.9,7)
STEQ|RCEQ|FMIN|FMAX|INDEF|DRAW

SOLV **SOLVR**

3:
2:
1:
(-.3,0)
(1.9,7)
X **LEFT** **RTE**

Use the displayed value as an initial guess.

X

X: (1.9,7)
2:
1:
(-.3,0)
X **LEFT** **RTE**

Calculate a solution to the equation by pressing the shift key followed by the Solver variable that you wish to solve.

SWAP **X**

X: 1.7807764064
Sign Reversal
1: 1.7807764064
X **LEFT** **RTE**

Repeat the procedure for the other point of intersection.

SWAP
X

X: (-.3,0)
2:
1: 1.7807764064
X **LEFT** **RTE**

$\equiv X \equiv$

X: - .280776406405
Sign Reversal
1: - .280776406405
X LEFT= RT=

Recall Y to compute the slope at an intersection point.

USER $\equiv Y \equiv$

3: 1.7807764064
2: -.280776406405
1: 2*x^2
X FPRMR EQ Y F ANG

Take the derivative with respect to x.

' X ENTER
d/dx

3: 1.7807764064
2: -.280776406405
1: -1.12310562562
X FPRMR EQ Y F ANG

Evaluate at one intersection point.

The last root computed remains assigned to x. The slope of the line can be read from the given expression.

3 ENTER

3: -280776406405
2: -1.12310562562
1: 3
X FPRMR EQ Y F ANG

Use the ANG function to compute the angle.

\equiv ANG \equiv

3: 1.7807764064
2: -.280776406405
1: -60.1164404136
X FPRMR EQ Y F ANG

This is in degrees.

Ready the stack to operate on the second intersection point.

DROP
DROP

3:
2:
1: 1.7807764064
X FPRMR EQ Y F ANG

Compute the derivative of Y .

Assigning a numeric value to x at this point will mean a numeric value for the derivative when it is computed.

' X **STO**

3:
2:
1:
X FPRM EQ Y F ANG

Y

3:
2:
1:
'2*X^2'
X FPRM EQ Y F ANG

The derivative is computed with respect to x .

' X **ENTER**

d/dx

3:
2:
1:
7.1231056256
X FPRM EQ Y F ANG

Enter the slope of the line.

3 **ENTER**

3:
2:
1:
7.1231056256
3
X FPRM EQ Y F ANG

Again use the ANG function to compute the intersection angle.

ANG

3:
2:
1:
-10.443524758
X FPRM EQ Y F ANG

Purge the variables created in the last two sections.

{ 'F''Y''X''ANG' **PURGE**

Differential Calculus

This chapter includes problems of differential calculus, including minimizing functions, calculating tangent lines, and several methods of implicitly differentiating functions. Several important features of the calculator are highlighted, including creating user-defined derivatives, the use of keyboard algebra for solving complex problems, and effective use of flag 35 for symbolic evaluation of constants. For HP-28C users, this chapter also describes use of user flag 59 (the infinite result flag).

Minimize Perimeter

Science, engineering, and business share the need to find the minimum values of given functions as some parameter changes. In this section, the function represents area, and the parameter is the area's perimeter.

Example: To minimize material expense, find the minimum amount of fencing required to enclose a rectangular plot measuring 200 square feet if one side is next to a building and needs no fence.

Let the sides be called x and y with y parallel to the building. The perimeter to be minimized is

$$P = 2x + y.$$

The area of the plot

$$x \cdot y = 200$$

gives the relationship between x and y .

Clear the display and make certain variables X and Y have no assigned values.

CLEAR **MODE** 2 **FIX** **<>**
'**X**' **PURGE**
'**Y**' **PURGE**

4:
3:
2:
1:

Note: HP-28C users must clear flag 59 to ignore "Infinite Result" errors that may occur while plotting. Before proceeding, press the following keys to clear flag 59.

59 **CF** **ENTER**

Enter the perimeter.

'**2*X+Y**' **ENTER**

4:
3:
2:
1: '2*X+Y'

Enter the area.

'**X*Y=200**' **ENTER**

4:
3:
2:
1: '2*X+Y'
'X*Y=200'

Isolate X .

'X ALGEBRA ISOL

3:	
2:	'2*X+Y'
1:	'200/Y'
TAYLR ISOL CURD SHOW DEGET ENSET	

Store the equation for X .

'X STO

3:	
2:	
1:	'2*X+Y'
TAYLR ISOL CURD SHOW DEGET ENSET	

Evaluate the expression for the perimeter.

EVAL

3:	
2:	
1:	'2*(200/Y)+Y'
TAYLR ISOL CURD SHOW DEGET ENSET	

This expresses the perimeter in terms of one variable.

Collect terms.

COLCT

3:	
2:	
1:	'400/Y+Y'
COLCT EXPAN SIZE FORM DESUB ENSUB	

Compute the derivative. Roots of this will yield the minimum value of Y .

'Y ENTER

d/dx

3:	
2:	
1:	'-(400/Y^2)+1'
COLCT EXPAN SIZE FORM DESUB ENSUB	

Plot the derivative to obtain a guess at the root.

PLOT STEQ


'PPAR PURGE

'Y INDEP

3:	
2:	
1:	
STEQ RECD FMIN FMAX INDEP DRAW	

The steps below expand the plotting area and draw the graph. If you have no prior knowledge of the appearance of the graph, you may first wish to plot the graph, modify the plotting area accordingly, and then plot the graph a second time. (Press **DRAW** followed by **ATTN**, and then proceed with the steps below).

2 ***H**
4 ***W**
DRAW



Digitize a seed for Y . Pick the guess near the positive root.

> ... **>** **INS**
ATTN

3:
2:
1: **(19.60, 0.00)**
STEP|RCED|FMIN|FMAX|INDEF|DRAW

Use the digitized value as a seed to compute Y .

SOLV **SOLVR**
Y
Y

Y: 20.00
Zero
1: **20.00**
Y EXPRE

The side parallel to the building must be 20 feet long.

Recall and evaluate the expression for X .

X **ENTER**
EVAL

3:
2:
1: **20.00**
10.00
Y EXPRE

Forty feet of fencing is required (two ends ten feet long and one side 20 feet long).

Purge the variables created in the example.

{ 'X' 'Y' **PURGE**

Mimimize Surface Area

This section uses differential calculus to minimize surface area. An application of this solution is in manufacturing, where minimization can reduce wasted raw material and increase profit. Other problem specifications may, of course, add constraints or considerations to the final real-world solution.

Example: Find the dimensions of a one liter can that has the minimum surface area.

Note: In this problem, user flag 35 is set to maintain symbolic constants until the end of the solution.

The surface area of a can (a right circular cylinder) is

$$A = 2\pi R^2 + 2\pi R H .$$

The volume is

$$V = \pi R^2 H$$

where R is the radius and H is the height of the can. To minimize the surface area, the area is expressed in terms of either R or H , and that expression is then differentiated with respect to that variable. Proceed by isolating H in the volume equation and finding the root of the derivative of the area taken with respect to R .

Clear the variables R , V , and H , and set flag 35.

CLEAR **<>**
{ **R** **V** **H** **ENTER** **PURGE**
35 **SF** **ENTER**

4:
3:
2:
1:

Factor out $2\pi R$ and key in the expression for the surface area.

' 2 π R(R+H) **ENTER**

4:
3:
2:
1: ' 2 π R(R+H) '

Duplicate the expression and store a copy for later use.

ENTER ' **A** **STO**

4:
3:
2:
1: ' 2 π R(R+H) '

Enter the volume.

'V=π×R^2×H **ENTER**

```
4:  
3:  
2:      '2*π*R*(R+H)'  
1:      'V=π*R^2*H'
```

Isolate H .

'H **ENTER**

```
4:  
3:  
2:      '2*π*R*(R+H)'  
1:      'H'
```

ALGEBRA **ISOL**

```
3:  
2:      '2*π*R*(R+H)'  
1:      'V/(π*R^2)'  
TAYLR ISOL QUDR SHOW DEGET EGET
```

Store it as H .

'H **STO**

```
3:  
2:  
1:      '2*π*R*(R+H)'  
TAYLR ISOL QUDR SHOW DEGET EGET
```

Now substitute for H in the area equation.

EVAL

```
2:  
1:      '2*π*R*(R+V/(π*R^2))'  
TAYLR ISOL QUDR SHOW DEGET EGET
```

Take the derivative with respect to R .

'R **ENTER**

d/dx

```
1:      '2*(R+V/(π*R^2))+2*  
      *π*R*(1-V*(π*(2*R))/  
      (π*R^2)^2)'  
TAYLR ISOL QUDR SHOW DEGET EGET
```

Collect terms.

COLCT

```
1:      '2*(1-2*(R^2*π)^(-2))  
      *R*V*π)*R*π+2*(R^(-2)  
      )*V/π+R)*π'  
COLCT EXPAN SIZE FORM DESUB ERSUB
```

Prepare to plot the derivative to obtain a guess for the root.

PLOT **STEQ**

```
3:  
2:  
1:  
STEQ RREC FMIN FMAX INDEF DRAW
```

One liter is the same as 1000 cubic centimeters. Enter the volume as 1000; the answer will be in centimeters.

1000 **ENTER**
'V **STO**

3:
2:
1:
STO RCL FMIN FMAX INDEF DRAW

Purge the existing plot parameters and expand the plotting area.

'PPAR **PURGE**
100 **■*H■**
5 **■*W■**

3:
2:
1:
PPAR RES AXES CENTR XW YW

Note: HP-28C users must clear flag 59 to ignore "Infinite Result" errors that may occur while plotting. Before proceeding, press the following keys to clear flag 59.

59 **CF** **ENTER**


To find the radius that minimizes the area, specify R as the independent plotting variable.

R **■INDEP■**

3:
2:
1:
INDEF RCL FMIN FMAX INDEF DRAW

Draw the graph and digitize an initial guess for the Solver.

DRAW **■** **>** **...** **>** **INS**



Now store the initial guess and compute the root.

ATTN **SOLV** **■SOLVR■**
R **■** **R**

R: 5.42
Zero
1: 5.42
R V EXPRE

This is the radius. Now find the height.

'H **ENTER**
EVAL

3:
2:
1: 5.42
'V/(π*R^2)

EVAL

3:	5.42				
2:					
1:	'1000/(\pi*29.37)'				
R	V	EXPRE			

Compute the area.

A EVAL

2:	'1000/(\pi*29.37)'				
1:	'2*\pi*5.42*(5.42+V/(\pi*R^2))'				
R	V	EXPRE			

Evaluate to a numerical result.

EVAL

2:	'1000/(\pi*29.37)'				
1:	'2*\pi*5.42*(5.42+1000/(\pi*29.37))'				
R	V	EXPRE			

Reduce the expression to a real number.

→NUM

3:	5.42				
2:	'1000/(\pi*29.37)'				
1:	553.58				
R	V	EXPRE			

To check that this is a minimum, compute the second derivative.

SOLV **'R' ENTER**

3:	553.58				
2:	'2*(1-2*(R^2*\pi)^(-2))'				
1:	'R'				
STO	RCEQ	SOLVE	ISOL	GUARD	SHOW

d/dx
→NUM

3:	'1000/(\pi*29.37)'				
2:	553.58				
1:	37.70				
STO	RCEQ	SOLVE	ISOL	GUARD	SHOW

The second derivative is positive; therefore, the curve is concave up. The root is a local minimum.

Purge the variables created in this problem section.

{ 'A' 'H' 'R' 'V' PURGE

Lines Tangent to a Circle

This section demonstrates manipulation of equations using the algebraic capabilities of the HP-28S and HP-28C. It is often necessary to calculate the derivative of a function that cannot easily be expressed in terms of one variable. In this case we use implicit differentiation. This is the first of three methods for implicit differentiation shown in this book. Problem sections "Implicit Differentiation With User-Defined Derivative" and "Implicit Functions" show two other methods.

Example: Find the two points on a circle of radius 1 that have tangent lines passing through the point (2,2).

There are two expressions for the slope of the tangent lines – one from the circle itself and the other from the point exterior to the circle.

Clear the working variables to ensure a symbolic answer. This problem also demonstrates a simple error recovery procedure. To ensure that the recovery works, turn on UNDO.

Note: The UNDO mode is set differently on the HP-28S and HP-28C. Both sets of instructions are provided below.

Also note that the **MODE** display is quite different on the two calculators. The displays used below depict the HP-28S **MODE** display. As shown below, on the HP-28S a small box appears in the **UNDO** menu item to indicate that it is on. On the HP-28C the **+UND** menu item is highlighted when the UNDO mode is active.

CLEAR
{ 'Y''R''B''A''EQ''X'
PURGE

3:
2:
1:
CMD UNDO LAST ML RDX PRMD

HP-28S Keystrokes:

MODE **UNDO**

HP-28C Keystrokes:

MODE **NEXT** **+UND**

The general equation for a circle is $x^2 + y^2 - r^2 = 0$, where r is the radius. Implicitly differentiate this equation.

Enter this equation for step-by-step differentiation. The " ∂ " character is obtained by pressing the **d/dx** key while entering an equation that begins with the **1** key.

' $\partial X(X^2 + Y^2 - R^2)$ **ENTER**

3:
2:
1: ' $\partial X(X^2 + Y^2 - R^2)$ '
CMD UNDO LAST ML RDX PRMD

EVAL

2:
1: ' $\partial X(X^2 + Y^2) - \partial X(R^2)$ '
CMD UNDO LAST ML RDX PRMD

Step through the derivative, watching for the term representing the dy/dx term.

EVAL

2:
1: ' $\partial X(X^2) + \partial X(Y^2) - \partial X(R^2) * 2 * R^{(2-1)}$ '
CMD UNDO LAST ML RDX PRMD

One more step-by-step differentiation will generate the dy/dx term from the $\partial X(Y^2)$ term in the expression.

EVAL

2:
1: ' $\partial X(X) * 2 * X^{(2-1)} + \partial X(Y) * 2 * Y^{(2-1)}$ '
CMD UNDO LAST ML RDX PRMD

Now collect terms to shorten the expression.

ALGEBRA COLCT

2:
1: ' $\partial X(X) * 2 * X + \partial X(Y) * 2 * Y$ '
COLCT EXPAN SIZE FORM DESUB EXSUB

This is a critical step. Replace the derivative sub-expression with a variable that can be isolated. Count all characters, except parentheses and quotes, up to and including the second partial derivative symbol (∂). The derivative symbol is the ninth item. Therefore "9" is used for making the substitution.

9 **ENTER**
' DY **EXSUB**

3:
2:
1: ' $\partial X(X) * 2 * X + D Y * 2 * Y'$
COLCT EXPAN SIZE FORM DESUB EXSUB

Evaluate once more to clear the last derivative.

EVAL

```
3:  
2:  
1: '2*X+DY*2*Y'  
COLCT EXPAN SIZE FORM DSUB ENSUB
```

Solve for $\frac{dy}{dx}$.

'DY ISOL

```
3:  
2:  
1: '-(2*X/Y/2)'  
TAYLR ISOL QUAD SHOW DERGET ENGET
```

Collect the 2's.

COLCT

```
3:  
2:  
1: '-(X/Y)'  
COLCT EXPAN SIZE FORM DSUB ENSUB
```

This is the slope of any line tangent to the circle. Tangent lines that pass through a point (A, B) exterior to the circle have slope $(y - B)/(x - A)$, where the point (x, y) is on the circle.

'(Y-B)/(X-A) ENTER

```
3:  
2:  
1: '-(X/Y)'  
COLCT EXPAN SIZE FORM DSUB ENSUB
```

This line must be a tangent to the circle; that is, the expressions for the slope must be equal.

= ENTER

```
3:  
2:  
1: '-(X/Y)=(Y-B)/(X-A)'  
COLCT EXPAN SIZE FORM DSUB ENSUB
```

Use algebra to solve for y .

Y

```
2:  
1: '-(X/Y*Y)=(Y-B)/(X-A)'  
COLCT EXPAN SIZE FORM DSUB ENSUB
```

Clear the denominators by collecting terms and multiplying through by denominator terms.

COLCT

```
2:  
1: '-X=INV(-A+X)*(-B+Y)'  
COLCT EXPAN SIZE FORM DSUB ENSUB
```

Extract the denominator term.

7 

```
3:  
2:  
1:      '-A+X'  
TAYLR ISOL QUDR SHOW DEGET EXGET
```


Since EXGET "consumes" the original expression, a copy should have been made first. It is easy to recover from the error.



```
2:  
1: '-X=INV(-A+X)*(-B+Y)  
*Y'  
TAYLR ISOL QUDR SHOW DEGET EXGET
```

Make a copy and re-execute EXGET.



7 

```
3:  
2: '-X=INV(-A+X)*(-B+Y...  
1: '-A+X'  
TAYLR ISOL QUDR SHOW DEGET EXGET
```

Multiply through by the extracted term.



```
2:  
1: '-(X*(-A+X))=INV(-A+  
X)*(-B+Y)*Y*(-A+X)'  
TAYLR ISOL QUDR SHOW DEGET EXGET
```

The denominator is now cleared.



```
2:  
1: '-((-A+X)*X)=(-B+Y)*  
Y'  
COLCT EXPAN SIZE FORM DESUB EXSUB
```

The following expansions distribute the x and y terms.



```
3:  
2:  
1: '-(-A+X)*X=-B*Y+Y*Y'  
COLCT EXPAN SIZE FORM DESUB EXSUB
```



```
3:  
2:  
1: '(A-X)*X=-B*Y+Y*Y'  
COLCT EXPAN SIZE FORM DESUB EXSUB
```



```
3:  
2:  
1: 'A*X-X*X=-B*Y+Y*Y'  
COLCT EXPAN SIZE FORM DESUB EXSUB
```

Now collect terms.

COLCT

```
3:  
2:  
1:  '-X^2+A*X=Y^2-B*Y'  
COLCT EXPAN SIZE FORM DESUB EXSUB
```

Gather like powers.

First gather powers of 2.

ENTER

1 EXGET

```
3:  
2:  
1:  '-X^2+A*X=Y^2-B*Y'  
     '-X^2'  
TAYLR ISOL QUDL SHOW DEGET EXGET
```

-

```
2:  
1:  '-X^2+A*X+X^2=Y^2-B*Y'  
     'Y+X^2'  
TAYLR ISOL QUDL SHOW DEGET EXGET
```

COLCT

```
3:  
2:  
1:  'A*X=-(B*Y)+X^2+Y^2'  
COLCT EXPAN SIZE FORM DESUB EXSUB
```

Now gather powers of 1.

ENTER

7 EXGET

```
3:  
2:  
1:  'A*X=-(B*Y)+X^2+Y^2'  
     'B*Y'  
TAYLR ISOL QUDL SHOW DEGET EXGET
```

+

```
2:  
1:  'A*X+B*Y=-(B*Y)+X^2+Y^2+B*Y'  
TAYLR ISOL QUDL SHOW DEGET EXGET
```

COLCT

```
3:  
2:  
1:  'A*X+B*Y=X^2+Y^2'  
COLCT EXPAN SIZE FORM DESUB EXSUB
```

The right-hand side of this equation is r^2 . Make a substitution for the right-hand side.

12 ENTER

'R^2 EXSUB

```
3:  
2:  
1:  'A*X+B*Y=R^2'  
COLCT EXPAN SIZE FORM DESUB EXSUB
```

This linear equation can now be solved for y .

Y **ISOL**

3:
2:
1: $'(R^2-A*x)/B'$
TAYLR ISOL GUARD SHOW DEGET ENGET

Save this for later use.

'Y **STO**

3:
2:
1:
TAYLR ISOL GUARD SHOW DEGET ENGET

Enter the equation for the circle.

'X^2+Y^2-R^2 **ENTER**

3:
2:
1: $'X^2+Y^2-R^2'$
TAYLR ISOL GUARD SHOW DEGET ENGET

Substitute in the expression for y .

EVAL

2:
1: $'X^2+((R^2-A*x)/B)^2-R^2'$
TAYLR ISOL GUARD SHOW DEGET ENGET

This is a quadratic equation for x , and is easy to solve.

'X **QUAD**

1: $'(A/B*2*(R^2/B)+s1*sqrt(-(A/B*2*(R^2/B)))*2-4*((2-A/B*2*(-(A/B))))'$
TAYLR ISOL GUARD SHOW DEGET ENGET

Shorten it by collecting terms.

COLCT

1: $'((sqrt(-(2*(2+2*A^2*B^(-2)))*((INV(B)*R^2)^2-R^2))+(-2*A*B^(-2))$
COLCT EXPAN SIZE FORM DESUB ENGET

Duplicate and store this expression for x .

ENTER **'X** **STO**

1: $'((sqrt(-(2*(2+2*A^2*B^(-2)))*((INV(B)*R^2)^2-R^2))+(-2*A*B^(-2))$
COLCT EXPAN SIZE FORM DESUB ENGET


In the Solver, you can assign the numbers needed to complete the given problem.

SOLV **STEQ**
SOLVR


3:
2:
1:
A B R S1 EXPRE

The exterior point is (2,2).

2 




2 




The radius of the circle is 1.

1 




There are two roots, one for each point on the circle.

1 




Solve for the x coordinate.






Now solve for the y coordinate.








Repeat the process for the other point.



-1 \equiv S1

S1:	-1.00
2:	0.91
1:	-0.41

Solve for the x coordinate.

\equiv EXPR =

EXPR=-0.41	
2:	-0.41
1:	-0.41

Now compute the y coordinate.

USER \equiv Y

3:	-0.41
2:	-0.41
1:	' $(R^2-A*x)/B'$

\rightarrow NUM

3:	-0.41
2:	-0.41
1:	0.91

The points of tangency are $(0.91, -0.41)$ and $(-0.41, 0.91)$.

The general solution approach solves the problem for any circle and any exterior point.

Purge the variables created in this problem section.

{ 'X'' Y'' A'' B'' R'' S1' PURGE

Implicit Differentiation With User-Defined Derivative

This section uses a user-defined derivative for implicit differentiation of a function. Refer to the reference manual for additional information.

Example: Given the equation $\sqrt{x} + \sqrt{y} = 3$, express $\frac{dy}{dx}$ in terms of x and y .

Create a user-defined derivative for the function $y(x)$. User-defined derivatives must take two inputs from the stack; the definition below simply discards them and returns the variable DY , which can be isolated.

CLEAR **<>**
<> x dx 'DY **ENTER**

4:
3:
2:
1: **<> x dx 'DY' >**

Store it in the variable *derY*.

'derY **STO**

4:
3:
2:
1:

Enter the *Y* variable as a function of *X*.

'√X + √Y(X) - 3 **ENTER**

4:
3:
2:
1: **'√X+√Y(X)-3'**

Differentiate with respect to *X*.

'X **ENTER** **d/dx**

3:
2:
1: **'INV(2*√X)+DY/(2*√Y(X))'**

Solve for *DY*. Remember that *DY* represents $\frac{dy}{dx}$.

'DY **ALGEBRA** **ISOL**

2:
1: **'-(INV(2*√X)*(2*√Y(X))'**
TAYLR ISOL QRD SHOW DEGET ENGET

Simplify to get the solution.

COLCT

```
3:  
2:  
1:      '-(JY(X)/JX)'  
COLCT EXPAN SIZE FORM RESUB ENSUB
```

Purge the user-defined derivative created in this example.

'derY PURGE

Taylor Series Error Term

Many physics and engineering problems are made solvable by expanding non-linear terms in a Taylor series. Ignoring the quadratic and higher degree terms leads to an approximate solution that is good for small displacement. This problem shows how to find the range for which the error in a Taylor series expansion stays small.

Example: Find the range of x for which the error in the 3rd degree approximation of $\sin(x)$ is less than .1.

The Taylor Series error term is

$$R_n(x) = f^{(n+1)}(c) \frac{x^{n+1}}{(n+1)!}$$

The exponent of f indicates the order of differentiation.

It is important to recognize that the error is the next term in the expansion. Since the sine function contains only odd-powered terms, look at the difference in the 5th and 3rd degree approximations. For the sine function the $n+1$ derivative has a maximum of 1.

$$\text{Thus } R_{(n+1)} < \frac{x^{n+1}}{(n+1)!}.$$

Compute the 5th degree expansion.

Set the angle mode. Key in the function and the variable name.

CLEAR **MODE** **RAD**
'SIN(X **ENTER**
X **ENTER**

3: **'SIN(X)**
2: **'X**
1:
STD **FIX** **SCI** **ENG** **DEG** **RAD**

Key in the order and find the Taylor Series.

5 **ALGEBRA** **TAYLR**

2: **'X-0.17*X^3+0.01*X^5**
1:
TAYLR **ISOL** **GUARD** **SHOW** **DEGET** **EGET**

Now compute the 3rd degree approximation.

'SIN(X **ENTER**
X **ENTER**
3 **TAYLR**

3: **'X-0.17*X^3+0.01*X^5**
2: **'X-0.17*X^3**
1:
TAYLR **ISOL** **GUARD** **SHOW** **DEGET** **EGET**

Make a copy and store this result for later use.

ENTER 'APS **STO**

```
3: 'X-0.17*X^3+0.01*X^5
2: 'X-0.17*X^3'
1: TAYLR ISOL QUAD SHOW DEGET EGET
```

Subtract the two approximations.

-

```
2: 'X-0.17*X^3+0.01*X^5
1: -(X-0.17*X^3)'
TAYLR ISOL QUAD SHOW DEGET EGET
```

Collect terms. The remaining expression is the 3rd degree error term.

COLCT

```
3:
2:
1: '0.01*X^5'
COLCT EXPAN SIZE FORM DEGET EGETSUB
```

Set it equal to .1 and then solve for x .

.1 **ENTER** = **ENTER**

```
3:
2:
1: '0.01*X^5=0.10'
COLCT EXPAN SIZE FORM DEGET EGETSUB
```

There are several ways to solve for x . The ISOL command will isolate x in the displayed equation and result in a generalized expression for x . A second approach is to use Solver to compute x . A third approach would be to use the laws of algebra and the capabilities of the calculator to solve for x "long-hand." All three methods are shown below; the third approach is included to illustrate the power of FORM in the ALBEGRA menu.

Choose any one of the three methods which follow, then proceed to the "Conclusion" portion of this problem.

Method 1: Using ISOL. Find the generalized expression for x . The status of flags 34 and 35 will affect the next display. The expression below is the result with both flags 34 and 35 clear. (To clear these flags, press 34 CF 35 CF.) Refer to the reference manual for a discussion on alternate settings for these flags. With flag 34 set, you would immediately obtain the result 1.64 found after the next several steps.

'X 

2:
1: 'EXP((0,00,6.28)*n1/
5)*1.64
TAYLR ISOL QRD SHOW DEGET EXGET

Assign a value of zero to the arbitrary integer $n1$ introduced into the isolation of the variable x .

0 
'n1 

2:
1: 'EXP((0,00,6.28)*n1/
5)*1.64
TAYLR ISOL QRD SHOW DEGET EXGET

Evaluate the expression.



3:
2:
1: (1.64,0.00)
TAYLR ISOL QRD SHOW DEGET EXGET

Extract the real component of the complex result.

3:
2:
1: 1.64
RES SIGN MANT EXPON

Now skip to the discussion and keystrokes labeled "Conclusion" to complete this problem.

Method 2: Using Solver. This method illustrates a simple approach to solve for x with the Solver.

Proceed to the Solver menu and store the equation.

SOLV STEQ
SOLVR

3:
2:
1:
X LEFT= RT=

Solve for the variable x .

 X

X: 1.64
Sign Reversal
1: 1.64
X LEFT= RT=

Now skip to the discussion and keystrokes labeled "Conclusion" to complete this problem.

Method 3: Using FORM and Algebraic Manipulation. This method illustrates the use of FORM and the keyboard capabilities of the calculator to manipulate algebraic expressions. While the two methods above are more direct, this alternative follows a traditional "paper-and-pencil" approach toward the solution.

First, compute the fifth root of the equation.

' 1/5 [ENTER] ^

2:
1: ' (0.01*X^5)^(1/5)=
0.10^(1/5)
COLCT EXPAN LEVEL EXGET [+/-]

Enter FORM, distribute the left hand exponential, and then associate the 5 and 1/5. Finally, collect terms in the expression.

FORM

((0.01*(X^5))^(1/5))=(
0.10^(1/5))
COLCT EXPAN LEVEL EXGET [+/-]

Move to the exponentiation sign.

[→] ... [→]

((0.01*(X^5))^(1/5))=(
0.10^(1/5))
COLCT EXPAN LEVEL EXGET [+/-]

Distribute the left-hand exponential.

↔D

$((0.01^{(1/5)}) * ((X^5)^{1/5})) = (0.10^{(1/5)})$

1/0 ← 0 0 0 ← ← ←

Move to the second exponentiation sign.

↔[→] ... ↔[→]

$((0.01^{(1/5)}) * ((X^5)^{1/5})) = (0.10^{(1/5)})$

COLCT EXPAN LEVEL EGET ← ←

Now associate the 5 and 1/5 in the expression.

A→

$((0.01^{(1/5)}) * (X^{(5*(1/5))})) = (0.10^{(1/5)})$

1/0 ← 0 0 0 ← ← ←

Exit FORM and collect terms.

ATTN COLCT

3:
2:
1: '0.38*X=0.63'
COLCT EXPAN SIZE FORM DESUB EXSUB

Solve for x .

'X ISOL

3:
2:
1: 1.64
TAYLR ISOL GUAD SHOW DEGET EGET

Conclusion: The variable x has now been isolated by one of the three methods described above. Proceed with the remainder of this problem solution.

The sine is symmetric, so $R^3 < .1$ for $-1.64 < x < 1.64$. Check the result in Solver.

USER APS

3:
2:
1: 'X-0.17*X^3'
APS

Compare the approximation to $\sin(x)$.

'SIN(X) ENTER

3: 1.64
2: 'X-0.17*X^3'
1: 'SIN(X)'
APS

= **ENTER**

3:			
2:			
1:	'X-0.17*X^3=SIN(X)'		
APS			

SOLV **STEQ**
SOLVR

3:			
2:			
1:	1.64		
X	LEFT=	RT=	

X

X:	1.64		
2:			
1:			
X	LEFT=	RT=	

LEFT=


LEFT=	0.90		
2:			
1:	0.90		
X	LEFT=	RT=	

RT=

RIGHT=	1.00		
2:		0.90	
1:	1.00		
X	LEFT=	RT=	

Clearly the difference is .1. Now plot the two equations. Purge the current plot parameters and draw the function.

PLOT 'PPAR' **PURGE**
DRAW



If the Taylor series approximation is needed for values of x that differ significantly from 0, the center of the expansion should be shifted, as demonstrated in the tangent line problem in the next section.

Exit from the **PLOT** screen and purge the variables created in this problem section.

ATTN { 'X' 'APS' 'EQ' **PURGE**

Tangent Lines and Taylor Series

This section demonstrates how to use the first order Taylor series to generate a tangent line equation. The example problem expands about a point other than the origin.

Example: Find the equation of the line tangent to the sine curve at $X = 1$.

Clear the stack. The first degree polynomial Taylor series expansion is the tangent line at the point of expansion.

Enter the function to be expanded.

CLEAR **<>**
'SIN(X) **ENTER**

4:
3:
2:
1:
'SIN(X)'

Change the variable to correspond with the new center. That is, $Y = 0$ corresponds to $X = 1$.

'Y+1 **ENTER**

4:
3:
2:
1:
'SIN(X)'
'Y+1'

'X **STO**

4:
3:
2:
1:
'SIN(X)'

This is the function to be expanded.

EVAL

4:
3:
2:
1:
'SIN(Y+1)'

Enter the variable and the degree of the polynomial.

'Y **ENTER**
1 **ENTER**

4:
3:
2:
1:
'SIN(Y+1)'
Y:
1.00

Find the Taylor expansion.

ALGEBRA TAYLR

3:
2:
1: '0.84+0.54*Y'
TAYLR ISOL GURO SHOW DEGET EXGET

This is the equation in Y .

USER X

3:
2:
1: '0.84+0.54*Y'
'Y+1'
X

Recall the change of variable equation.

'X ENTER
= ENTER

3:
2:
1: '0.84+0.54*Y'
'Y+1=X'
X

Clear the original variable change equation and solve for Y .

'X PURGE
'Y ENTER
ALGEBRA ISOL

3:
2:
1: '0.84+0.54*Y'
'X-1'
TAYLR ISOL GURO SHOW DEGET EXGET

Save the expression for Y .

'Y STO

3:
2:
1: '0.84+0.54*Y'
TAYLR ISOL GURO SHOW DEGET EXGET

Change back to the original variable and simplify the resulting expression.

EVAL

3:
2:
1: '0.84+0.54*(X-1)'
TAYLR ISOL GURO SHOW DEGET EXGET

EXPAN

2:
1: '0.84+(0.54*X-0.54*1)
)'
COLCT EXPAN SIZE FORM DESUB EXSUB

COLCT

3:
2:
1: '0.30+0.54*X'
COLCT EXPAN SIZE FORM DESUB EXSUB

Save a copy of this expression for the next problem section.

ENTER
'STN **STO**


3:
2:
1: '0.30+0.54*X'
COLCT EXPAN SIZE FORM OBSUB ENSUB

Plot the two equations for a quick check.

'SIN(X) **ENTER**
= **ENTER**

3:
2:
1: '0.30+0.54*X=SIN(X)'
COLCT EXPAN SIZE FORM OBSUB ENSUB

PLOT STEQ
'PPAR **PURGE**
'X INDEP
DRAW



Exit from the **PLOT** screen and purge variables X and Y for the next problem section.

ATTN { 'X' 'Y' **PURGE**

Normal Line

In the previous problem section, the equation for the line resulted from a Taylor series expansion. This problem section continues by manually assembling the expression for the normal line.

Example: Calculate the equation of the line normal (perpendicular) to the sine curve at $x = 1$.

First recall the equation for the tangent line.

CLEAR
USER STN
ENTER ENTER

3:	'0.30+0.54*x'
2:	'0.30+0.54*x'
1:	'0.30+0.54*x'
FFPR EQ STN	

Find the value of the function at $x = 1$. Evaluate the expression.

1 ENTER
'X STO
EVAL

3:	'0.30+0.54*x'
2:	'0.30+0.54*x'
1:	0.84
X FFPR EQ STN	

This is Y_0 .

To determine the symbolic solutions, purge the value of x .

'X PURGE

3:	'0.30+0.54*x'
2:	'0.30+0.54*x'
1:	0.84
FFPR EQ STN	

The general point slope formula for a line is

$$Y - Y_0 = m(X - X_0)$$

Y_0 is on the stack. Form the left-hand side of the relationship above.

Y ENTER
SWAP
-

3:	'0.30+0.54*x'
2:	'0.30+0.54*x'
1:	'Y-0.84'
FFPR EQ STN	

Now form the right-hand side. Bring the original line in position to find the slope.

SWAP
'X ENTER

3:	'Y-0.84'
2:	'0.30+0.54*x'
1:	'X'
FFPR EQ STN	

Find the slope by taking the derivative.

d/dx

3:	'0.30+0.54*X'			
2:	'Y-0.84'			
1:	0.54			
PFMR	EQ	STN		

This is the slope of the tangent line. The slope of the normal line is

$$m_n = -\frac{1}{m_t}.$$

Compute m_n .

CHS

1/x

3:	'0.30+0.54*X'			
2:	'Y-0.84'			
1:	-1.85			
PFMR	EQ	STN		

Now compute the right-hand side.

'X-1 **ENTER**

3:	'Y-0.84'			
2:	-1.85			
1:	'X-1'			
PFMR	EQ	STN		

×

3:	'0.30+0.54*X'			
2:	'Y-0.84'			
1:	'-(1.85*(X-1))'			
PFMR	EQ	STN		

Form the entire equation.

= **ENTER**

2:	'0.30+0.54*X'			
1:	'Y-0.84=-(1.85*(X-1))'			
PFMR	EQ	STN		

Solve for Y .

'Y **ENTER**

ALGEBRA **ISOL**

3:	'0.30+0.54*X'				
2:	'Y-0.84=-(1.85*(X-1))+0.84'				
1:	'Y=-(1.85*(X-1))+0.84'				
TAYLR	ISOL	QUAD	SHOW	DEGET	EXGET

Simplify the expression.

EXPN

3:	'0.30+0.54*X'				
2:	'Y-0.84=-(1.85*(X-1))+0.84'				
1:	'Y=-(1.85*(X-1))+0.84'				
COLCT	EXPAN	SIMPL	FORM	DESIMP	EXSUB

EXPN

2: '0.30+0.54*x'
1: '-1.85*x--1.85*x+'
0.84'
COLCT EXPAN SIZE FORM RESUB ENSUB

COLCT


3:
2: '0.30+0.54*x'
1: '2.69-1.85*x'
COLCT EXPAN SIZE FORM RESUB ENSUB

Plot the resulting function.

'SIN(X) ENTER
= ENTER

3:
2: '0.30+0.54*x'
1: '2.69-1.85*x=SIN(X)'
COLCT EXPAN SIZE FORM RESUB ENSUB

PLOT STEQ
'PPAR PURGE
'X INDEP
DRAW



Exit from the PLOT screen and purge the following variables.

ATTN { 'STN' 'EQ' 'PPAR' PURGE

Implicit Functions

The Implicit Function Theorem is, perhaps, the most elegant of three methods shown for implicit differentiation. This problem section demonstrates a more general method for finding the equation of a line than the previous problems sections.

Example: Find the equation of the line tangent to the function $x^2 + xy - 3 = 0$ at $x = 1$.

Begin by defining a function to compute the derivative of a general function $F(x, y)$. The formula, a result of the implicit function theorem, can be used as long as $\frac{\partial F}{\partial y} \neq 0$ holds.

Purge the variables that will be used in this example to ensure symbolic solutions.

CLEAR
{ 'X'' Y'' y'' x' **PURGE** <>

4:
3:
2:
1:

Enter the function for computing implicit derivatives.

«→ a' -∂X(a)÷∂Y(a)
ENTER

3:
2:
1: «→ a' -∂X(a)÷∂Y(a)'
»

Store the implicit derivatives function.

'IMP **STO**

4:
3:
2:
1:

Enter and store the general formula for a line.

'y=m*(x-X)+Y **ENTER**

4:
3:
2:
1: 'y=m*(x-X)+Y'

'LINE **STO**

4:
3:
2:
1:

The function must be expressed in terms of X and Y due to the use of those variables in the function IMP.

' $X^2+X*Y-3$ **ENTER**

4:	
3:	
2:	
1:	' $X^2+X*Y-3$ '

' F **STO**

4:	
3:	
2:	
1:	

Now find $\frac{dy}{dx}$ in terms of X and Y .

USER 

3:				
2:				
1:	' $X^2+X*Y-3$ '			
F	LINE	IMP		



2:				
1:	' $-(\partial X(X^2+X*Y)/\partial Y(X^2+X*Y))'$			
F	LINE	IMP		

Evaluate the expression until all the partial derivative symbols are gone.

EVAL

2:				
1:	' $-(\partial X(X^2)+\partial X(X*Y))/(\partial Y(X^2)+\partial Y(X*Y))$ '			
F	LINE	IMP		

EVAL

1:	' $-(\partial X(X)*2*X^(2-1)+(\partial X(X)*Y+X*\partial X(Y)))/(\partial Y(X)*2*X^(2-1)+\partial Y(X))$ '			
F	LINE	IMP		

EVAL

3:				
2:				
1:	' $-(2*X+Y)/X$ '			
F	LINE	IMP		

This expression for the slope of $F(x,y)$ at any point on the curve must be the slope of the tangent line.

' m **STO**

3:				
2:				
1:				
M	F	LINE	IMP	

Now determine the value of Y that corresponds to $x = 2$.

F

3:
2:
1:
'X^2+X*Y-3'
M F LINE IMP

SOLV **STEQ**
SOLVR

3:
2:
1:
X Y EXPRE

2 X

X: 2.00
2:
1:
X Y EXPRE

EXPR=

EXPR='4+2*Y-3'
2:
1:
'4+2*Y-3'
X Y EXPRE

Solve for Y .

'Y ALGEBRA ISOL

3:
2:
1:
-0.50
TAYLR ISOL GUARD SHOW OBJGET ENGET

'Y STO

3:
2:
1:
TAYLR ISOL GUARD SHOW OBJGET ENGET

With the coordinates of the point at the tangent line and the slope of the line in terms of those coordinates, evaluate and simplify the formula for the line.

USER LINE

3:
2:
1:
'y=m*(x-X)+Y'
Y X EQ M F LINE

EVAL

2:
1:
'y=-((2*X+Y)/X*(x-2))
>-0.50'
Y X EQ M F LINE

EVAL

2:
1: 'y=-1.75*(x-2)-
0.50'
Y X EQ M F LINE

Use EXPAN to distribute the constant.

ALGEBRA

EXPAN

3:
2:
1: 'y=-1.75*(x-2)-0.50'
COLCT EXPAN SIZE FORM DESUB EXSUB

EXPAN

2:
1: 'y=-1.75*x--1.75*2-
0.50'
COLCT EXPAN SIZE FORM DESUB EXSUB

Finally, simplify the equation for the tangent line.

COLCT

3:
2:
1: 'y=3-1.75*x'
COLCT EXPAN SIZE FORM DESUB EXSUB

Purge the variables created in this problem section.

{ 'Y'' X'' EQ'' M'' F'' LINE'' IMP' PURGE

Integral Calculus

This chapter solves a number of problems of integral calculus, including integration of simple differential equations and computation of arc lengths, surfaces, and volumes. Both symbolic and numerical solutions are demonstrated with appropriate use of system flags.

Integration and Free Falling Body

This problem section demonstrates derivation of standard equations of motion through simple integration. The example illustrates the importance of the constant of integration and shows how that constant is incorporated into the solution provided by the HP-28S and HP-28C.

Example: A stone is dropped from a bridge 100 feet above the water. Calculate how long it takes to reach the water and its final velocity.

From Newton's 2nd law

$$F = m\ddot{x} .$$

The only force acting on a falling body is that of gravity.

$$F = -mg$$

Combining these,

$$\ddot{x} = -g .$$

This is the equation of motion for a freely falling body. A well-posed problem requires two initial conditions, the starting position and velocity. The problem then may be solved by integration.

This solution approach plots the final equation to facilitate root finding. Start by configuring the plot parameters.

```
CLEAR PLOT
'PPAR PURGE
100  ■*H■
(0, -70 ■PMIN■
```

```
3:
2:
1:
STEQ RCED PMIN FMAX INDEF DRAW
```

Plot the displacement as a function of time. Let TM represent the time.

```
'TM ■INDEF■
```

```
3:
2:
1:
STEQ RCED PMIN FMAX INDEF DRAW
```

Start by integrating the above equation. Let GRV be the acceleration due to gravity. Since the expression to be integrated includes no TM terms, the specified degree of the polynomial is zero.

'-GRV [ENTER]
'TM [ENTER]
0 [ENTER]

3:
2:
1:
'-GRV'
'TM'
0.00
STO RCL PMIN PMAX INDEF DRAW

[I]

3:
2:
1:
'- (GRV*TM)'
STO RCL PMIN PMAX INDEF DRAW

This is an expression for the velocity. At $TM = 0$ the initial velocity is $V0$.

V0 [+]

3:
2:
1:
'- (GRV*TM)+V0'
STO RCL PMIN PMAX INDEF DRAW

Store this for future use.

'VEL [STO]

3:
2:
1:
STO RCL PMIN PMAX INDEF DRAW

Now recall the velocity and prepare for a second integration. The integrand includes TM to the first degree, so a 1 is specified for the last parameter to the integration.

USER [] VEL []
'TM [ENTER]
1 [ENTER]

3:
2:
1:
'- (GRV*TM)+V0'
'TM'
1.00
VEL PPAR

[I]

3:
2:
1:
'V0*TM-GRV/2*TM^2'
VEL PPAR

This is an expression for the displacement. At $TM = 0$, $x = X0$.

X0 [+]

2:
1:
'V0*TM-GRV/2*TM^2+X0'
VEL PPAR

To put this in the standard form, use the expression manipulation capabilities in FORM.

ALGEBRA FORM

$((V0*TM) - ((GRV/2)*(TM^2))) + X0$

COLCT EXPAN LEVEL EXGET (+) (-)

Move the cursor to the minus sign.

→ → → →

$((V0*TM) - ((GRV/2)*(TM^2))) + X0$

COLCT EXPAN LEVEL EXGET (+) (-)

Commute the expressions about the minus sign.

↔ ↔ ↔

$((-((GRV/2)*(TM^2)) + V0 * TM) + X0)$

(-) (+) (M) (M+) (H) (H+)

Exit FORM, make a copy, and save the expression for distance.

ATTN ENTER
'DST' STO

2:
1: ' $-(GRV/2*TM^2) + V0*TM + X0$ '

COLCT EXPAN SIZE FORM DEBUG EXSUB

Store the expression for use in the Solver menu.

SOLV STEQ
SOLVR

3:
2:
1:
GRV TM V0 X0 EXPRE

In English units the acceleration due to gravity is 32 ft/sec/sec.

32 GRV

GRV: 32.00
2:
1:
GRV TM V0 X0 EXPRE

The bridge is 100 feet high.

100 X0

X0: 100.00
2:
1:
GRV TM V0 X0 EXPRE

Since the stone is dropped, the initial velocity is zero.

0 V0

V0: 0.00
2:
1:
GRV TM V0 X0 EXPRE


Evaluate the expression EQ.

EXPR =

1: $-(16*TM^2)+100$
2:
1: $-(16*TM^2)+100$
GRV TM V0 X0 EXPR=

To find the time required to hit the water, find a root of this equation.
Digitize an initial guess from a plot of the equation.

PLOT DRAW
 INS



Assign the seed to TM .

ATTN SOLV SOLVR
 TM

1: $TM: (2.50, -3.28)$
2:
1: $-(16*TM^2)+100$
GRV TM V0 X0 EXPR=

Solve for TM .

TM

1: $TM: 2.50$
Zero
1: 2.50
GRV TM V0 X0 EXPR=

The stone hits the water after 2.5 seconds. To find the velocity, recall VEL and evaluate it.

USER VEL
 EVAL

1: $-(16*TM^2)+100$
2: 2.50
1: -80.00
VEL FPHR

The stone is falling at 80 feet per second.

By changing the initial conditions, the equations of motion developed in the previous example can be applied to a rock thrown straight up.

Example: A stone is thrown straight up from ground level with an initial velocity of 70 feet per second.

Compute its peak, the time elapsed until it hits the ground, and its final velocity.

Recall the general equation for distance traveled.

CLEAR					
USER	DST				
2: 1: $1 - (GRV/2 \cdot TM^2) + V0 \cdot TM + X0$					
TM	V0	X0	GRV	EQ	DST

Enter the **SOLV** menu and store the equation for analysis, then enter the Solver.

SOLV	STEQ
SOLVR	
3: 2: 1: GRV TM V0 X0 EXPRE	

The initial position is ground level or $x = 0$.

0 X0	
2: 1: GRV TM V0 X0 EXPRE	

The initial velocity is 70 feet per second upward, and, therefore, positive.


70 V0	
2: 1: GRV TM V0 X0 EXPRE	

The plot parameters were set in the previous problem. Plot both the velocity and the distance equations.

USER	DST
VEL	
=	ENTER
2: 1: $1 - (GRV/2 \cdot TM^2) + V0 \cdot TM + X0 = - (GRV \cdot TM) + V0$	
VEL	PPHR

Store the equation for plotting.

PLOT	STEQ
DRAW	



The velocity is the first derivative of the distance; therefore, the root of the velocity equation corresponds to a maximum of the distance equation. Digitize the roots of the velocity (where the straight line crosses the x-axis) and the distance (where the curve crosses the x-axis for the second time).

...

 ...

ATTN

3:		(2.30, -3.23)					
2:		(4.45, -3.23)					
1:		' - (GRV*TM)+V0'					
STEQ		FREQ		FMIN		FMAX	
INDEF		DRAW					

Recall the equation for velocity and save the equation for analysis.

USER

3:		(2.30, -3.23)					
2:		(4.45, -3.23)					
1:		' - (GRV*TM)+V0'					
VEL		FPHR					

SOLV

SOLVR

3:		(2.30, -3.23)					
2:		(4.45, -3.23)					
1:		GRV TM V0 EXPRE					

Enter the initial guess for the root and solve for TM .

SWAP

TM: 2.19		Zero					
1:		2.19					
GRV TM V0 EXPRE							

After 2.19 seconds, the stone reaches a maximum height. Recall the distance equation from the User menu and evaluate to find this height.

USER

2:		2.19					
1:		' - (GRV/2*TM^2)+V0*TM +X0'					
TM V0 X0 GRV EQ DST							

EVAL

3:		(4.45, -3.23)					
2:		2.19					
1:		76.56					
TM V0 X0 GRV EQ DST							

The rock reaches a height of 76.56 feet.

Now drop two numbers from the stack and recall the distance equation for analysis.

DROP
DROP
DST

2:		(4.45, -3.23)					
1:		' - (GRV/2*TM^2)+V0*TM +X0'					
TM V0 X0 GRV EQ DST							

SOLV **STEQ**
SOLVR

3:
2:
1: $(4.45, -3.23)$
GRV TM V0 X0 EXPRE

Enter the guess and solve for the root.

TM **TM**

1: 4.38
Zero
1: 4.38
GRV TM V0 X0 EXPRE

The rock hits the ground after 4.38 seconds. Note that this is exactly twice the time required to reach the maximum height. Therefore the time spent going up is equal to the time spent falling back to the ground. To find the final velocity recall the velocity equation and evaluate.

USER **VEL**

3:
2:
1: $-(GRV*TM)+V0$
VEL FFHR

EVAL

3:
2:
1: 4.38
-70.00
VEL FFHR

Note that this number differs from the initial velocity in sign only. The rock's final speed is the same as its initial speed, but it is traveling in the opposite direction.

Purge the variables created in this problem section.

{ 'TM' 'EQ' 'VEL' 'DST' 'GRV' 'X0' 'V0' 'PPAR'
PURGE

Double Integration

This problem section uses both symbolic and numerical integration to solve common problems of integral calculus.

Example: Compute the area between the line

$$Y = x$$

and the parabola

$$Y = x^2.$$

The area may be found by computing the double integral $\int_1^0 \int_{x^2}^x dy \, dx$.

To insure a symbolic answer, purge the constant and the variable of integration.

CLEAR
{ 'C' 'Y' **PURGE** <>

4:
3:
2:
1:

The next four displays show the calculator steps to compute $\int c \, dy$ where

$c = 1$. Because the result is simply y , you can choose to skip directly to the evaluation of the integral at its limits if you wish. If so, simply enter Y , and proceed to the steps below beginning with "Enter the upper limit."

Otherwise, prepare the stack for a symbolic integration with a first degree result. Start by integrating a constant.

'C **ENTER**
'Y **ENTER**
1 **ENTER**

4:
3:
2:
1: 'C'
1: 'Y'
1: 1.00

Execute the integral.

I

4:
3:
2:
1: 'C*Y'

Eliminate the constant by equating it to 1.

1 **ENTER**
'C **STO**

4:	
3:	
2:	
1:	'C*Y'

EVAL

4:	
3:	
2:	
1:	'Y'

Enter the upper limit.

'X **ENTER**
'Y **STO**

4:	
3:	
2:	
1:	'Y'

Save a copy of the integrand for later use and evaluate the integral at the limit.

ENTER
EVAL

4:	
3:	
2:	
1:	'Y' 'X'

Repeat the process for the lower limit.

'X^2 **ENTER**
'Y **STO**

4:	
3:	
2:	
1:	'Y' 'X'

Place a copy of the integrand in position for evaluation at the lower limit.

SWAP
EVAL

4:	
3:	
2:	
1:	'X' 'X^2'

The difference is the integrand for the second integration.

-

4:	
3:	
2:	
1:	'X-X^2'

Key in the parameters for the integration.

{ X 0 1 [ENTER]

4:	
3:	
2:	
1:	'X-X^2' (X 0.00 1.00)

Key in the error bound.

.005 [ENTER]

4:	
3:	
2:	
1:	'X-X^2' (X 0.00 1.00) 0.01

Evaluate the second integral. The error bound provides accuracy to the number of displayed digits (assuming 2 \equiv FIX \equiv).

[∫]

4:	
3:	
2:	
1:	0.17 8.37E-4

The area is 0.17.

Purge the variables created in this problem section.

{ 'Y' 'C' [PURGE]

Area Between Two Curves

This problem section provides a general approach for finding the area between any two intersecting curves.

Example: Find the area enclosed by the parabola $f(x) = x^2$ and the line $y(x) = x + 3$.

The area between two curves can be found by computing the integral $\int_a^b |f(x) - y(x)| dx$. In this problem the limits will be the intersection points of the curves.

Enter and store the integrand.

CLEAR **<>**
'ABS(F-Y) **ENTER**

4:	'ABS(F-Y)'
3:	
2:	
1:	

'AREA **STO**

4:	
3:	
2:	
1:	

Enter and store the functions.

'X^2 **ENTER**

4:	'X^2'
3:	
2:	
1:	

'F **STO**

4:	
3:	
2:	
1:	

'X+3 **ENTER**

4:	'X+3'
3:	
2:	
1:	

'Y **STO**

4:	
3:	
2:	
1:	

Plot both curves to find the intersection points.

' F=Y [ENTER]

4:
3:
2:
1:
'F=Y'

[EVAL]

4:
3:
2:
1:
'X^2=X+3'


Store the equation and set the plot parameters. If you have no prior knowledge of the graph of the curves, you can first draw the graph, exit and modify the plot parameters as shown below, then proceed with a second graph.

[PLOT] [STEQ]
' PPAR [PURGE]
5 [H]

3:
2:
1:
PFMR RES AXES CENTR SW [H]

The rightmost intersection point will become the upper limit. The leftmost intersection point is the lower limit. Draw the equation and digitize the rightmost point first, followed by the leftmost point.

[DRAW]



[>] ... [↑] [INS]
[↓] ... [←] [INS]
ATTN

3:
2:
1:
(2.30, 5.50)
(-1.40, 2.00)
STEQ RCEQ FMIN FMAX INDEF DRAW

Use the Solver to refine the initial guess.

[SOLV] [SOLVR]

3:
2:
1:
(2.30, 5.50)
(-1.40, 2.00)
X LEFT= RT=

[X]
[] [X]

X: -1.30
Sign Reversal
1:
X LEFT= RT= -1.30

Repeat the process for the upper limit.

SWAP
X
X

X	2.30				
Sign Reversal					
1:	2.30				
X	LEFT	RT			

The limits are in the correct order for integration, but the variable is missing. Manipulate the stack to put it in place.

'X ENTER
3 STACK ROLLD

3:	'X'				
2:	-1.30				
1:	2.30				
ROLLD	PICK	QOPEN	OPEN	DEPTH	LIST

Now convert the three elements to a list.

3 LIST →LIST

3:				
2:	{ X -1.30 2.30 }			
1:	LIST			
LIST	PUT	GET	PUTI	GETI

Recall the integrand.

USER AREA

3:	{ X -1.30 2.30 }
2:	ABS(F-Y)
1:	X FPAR EQ Y F AREA

Put them in the necessary order.

SWAP

3:	'ABS(F-Y)'
2:	{ X -1.30 2.30 }
1:	X FPAR EQ Y F AREA

Enter the error and integrate.

.005 ENTER

3:	'ABS(F-Y)'				
2:	{ X -1.30 2.30 }				
1:	0.01				
X	FFAR	EQ	Y	F	AREA

I

3:	7.81
2:	0.04
1:	
X PPAR EQ Y F AREA	

The area is 7.81.

Purge the variables created in this problem section.

{ 'AREA' 'F' 'Y' 'EQ' 'X' 'PPAR' PURGE

Arc Length

This section demonstrates keystroke and programming examples for computing arc lengths of rectifiable functions. The program ARC created in the second example is used in a later section entitled "Surface Area."

Example: Find the length of the curve

$$F(x) = \frac{(\sqrt{x^2+2})^3}{3}$$

from $x = 0$ to $x = 3$.

The arc length of a function is found by evaluating the integral

$$\int_a^b \sqrt{1+f'(x)^2} dx.$$

First form the integrand. Enter the given function in terms of x .

CLEAR **<>**
' ((X^2+2)^(3/2))÷3 **ENTER**

4:
3:
2:
1: ' (X^2+2)^(3/2)/3'

Specify the variable of differentiation.

' X **ENTER**

4:
3:
2:
1: ' (X^2+2)^(3/2)/3' X

Take the derivative and simplify.

d/dx

3:
2:
1: ' 2*X*1.50*(X^2+2)^0.50/3'

Collect terms.

ALGEBRA **COLCT**

3:
2:
1: ' (2+X^2)^0.50*X'
COLCT EXPAND SIZE FORM DESUB ENSUB

Square the derivative, add one, and take the square root.

x² 1 **+** **✓**

2:
1: ' ((SQ((2+X^2)^0.50*X)+1)^2'
COLCT EXPAND SIZE FORM DESUB ENSUB

This is the differential of arc length.

Place the list containing the variable and limits of integration on the stack.

{ X 0 3 [ENTER]

```
3:  
2: 'SQ((2+X^2)^0.50*...  
1: { X 0.00 3.00 }  
COLCT EXPAN SIZE FORM DESUB EXSUB
```

Specify the accuracy and perform the integration.

.005 [ENTER]

[ʃ]

```
3: 12.00  
2:  
1: 0.06  
COLCT EXPAN SIZE FORM DESUB EXSUB
```

The arc length is 12.00.

Example: Compute the arc length of $f(x) = x^2$ for $x = 0$ to $x = 2$.

For repeated problems, a simple program facilitates the computation of arc length. The program below differentiates the function with respect to X . This means that functions must be entered in terms of X .

The partial derivative symbol " ∂ " is obtained by pressing the **[d/dx]** key.

CLEAR
« → X '√(1+∂X(X)^2)
ENTER

2:
1: « → X '√(1+∂X(X)^2)'
»
COLCT EXPAN SIZE FORM ABSUB EXSUB

Examine this function to see that it is equivalent to the integrand in the previous example.

Store the program in the variable ARC.

'ARC **STO**

3:
2:
1:
COLCT EXPAN SIZE FORM ABSUB EXSUB

The program below first stores the error in the variable ER , then converts the next three levels of the stack to the list required for integration. The function is then brought to level 1 and operated on by the ARC function. Finally the function is returned to its position, and the error is recalled. The integration completes the process.

« 'ER' STO 3 →LIST
SWAP ARC SWAP ER
∫ **ENTER**

1: « 'ER' STO 3.00
→LIST SWAP ARC SWAP
ER 'J' »
+LIST LIST+ PUT GET PUTI GETI

Store the program ARCP.

'ARCP **STO**

3:
2:
1:
+LIST LIST+ PUT GET PUTI GETI

Computing the arc length of any function now only requires placing the correct information on the stack. This program requires the function on level 5, the variable of integration on level 4, the upper limit on level 3, the lower integration limit on level 2, and the error bound on level 1.

'X^2' 'X' 0 2 .005
ENTER

3: 0.00
2: 2.00
1: 0.01
+LIST LIST+ PUT GET PUTI GETI

Compute the arc length.

USER 

3:	4.65
2:	0.02
1:	
ER ARCP MRC	

Purge the variable *ER* and, if you wish, the program ARCP. Program ARC is used in the next problem section.

{ 'ARCP' 'ER' PURGE

Surface Area

The function created to compute arc lengths can be extended to computing surface areas.

Example: Compute the surface area of the solid formed by revolving the section of $f(x) = x^2$ between 0 and 1 about the x axis.

In this problem the integrand is expressed in terms of a function of x . The surface area can be computed from

$$S = \int_a^b 2\pi f(x) \sqrt{1+f'(x)^2} \, dx.$$

The square root factor in the integrand is identical to the ARC function used in the problem section entitled "Arc Length." If you have not already done so, key in the ARC function from the previous section. Enter the integrand using ARC as a function.

CLEAR **< >**
'2*xxFxARC(F) **ENTER**

4:	
3:	
2:	
1:	'2*π*F*ARC(F)'

Enter the function to be integrated.

'X^2 **ENTER**

4:	
3:	
2:	'2*π*F*ARC(F)'
1:	'X^2'

Store the function by the corresponding name appearing in the integrand.

'F **STO**

4:	
3:	
2:	
1:	'2*π*F*ARC(F)'

Purge the variable of integration to ensure that the name is not in use.

'X **PURGE**

4:	
3:	
2:	
1:	'2*π*F*ARC(F)'

Enter the variable of integration and the limits.

{X 0 1 **ENTER**

4:	
3:	
2:	
1:	'2*π*F*ARC(F)'
	{ X 0.00 1.00 }

Enter the error bound and compute the surface area.

.005 **ENTER**
[f]

4:	
3:	
2:	
1:	3.81 0.02

The surface area is 3.81.

Purge the variables created in this problem section.

{ 'F' 'ARC' **PURGE**

Arc Length of Parametric Equations

It is often necessary to work with equations expressed in terms of a parameter. The coordinates of a particle moving in a plane as a function of time is a common example.

Example: Compute the length of the curve corresponding to the equations

$$x(t) = \frac{t^2}{2} \quad \text{and} \quad y(t) = \frac{(2t+1)^{\frac{3}{2}}}{3}$$

for $t = 0$ to $t = 4$.

In parametric form the arc length is

$$L = \int_a^b \sqrt{dx^2 + dy^2}.$$

Enter the integrand in terms of the differentials of x and y . This general relationship can be used for any set of parametric equations with T as the parameter.

CLEAR **<>**
' $\sqrt{(\text{SQ}(\partial T(X)) + \text{SQ}(\partial T(Y))}$
ENTER

3:
2:
1: 'sqrt(SQ(∂T(X))+SQ(∂T(Y)))'

Save the parametric arc length in PARC.

' PARC **STO**

4:
3:
2:
1:

Enter the parametric equations. Store them under the names X and Y as expected by the PARC function.

' T^2/2 **ENTER**
' (2*T+1)^(3/2)/3 **ENTER**

4:
3:
2:
1: 'T^2/2'
1: '(2*T+1)^(3/2)/3'

'Y **STO**
'X **STO**

4:	
3:	
2:	
1:	

Now integrate with respect to T from 0 to 4.

First recall the integrand.

USER **PARC**

2:	
1:	' $\int(SQ(\partial T(X))+SQ(\partial T(Y))$
) 2

Key in the variable of integration and the limits.

{T 0 4 **ENTER**

3:	
2:	' $\int(SQ(\partial T(X))+SQ(\partial T(Y))$
1:	{T 0.00 4.00}

Enter the desired error bound.

.005 **ENTER**

3:	' $\int(SQ(\partial T(X))+SQ(\partial T(Y))$
2:	{T 0.00 4.00}
1:	0.01

Now perform the integration.

J

3:	
2:	
1:	12.00
	0.06

The arc length is 12.00.

Program PARC is used in the next program section, and X and Y are replaced by new functions.

Surface Area of Parametric Equations

The function created to compute arc lengths can be extended to compute surface areas. The surface area can be found by computing the integral

$$S = \int_a^b 2\pi Y \sqrt{dx^2 + dy^2}.$$

Example: Determine the surface area of the sphere formed by revolving a circle about the x axis.

$$x(t) = 2\cos(t) \quad y(t) = 2\sin(t)$$

These are the parametric equations for a circle of radius 2.

Note that the integrand includes the parametric arc length as a factor. Use the function defined in the previous section in the integrand. Clear user flag 35 for numeric evaluation of π when it is supplied as a limit to the integration.

CLEAR 35 CF **ENTER**
'2×π×Y×PARC **ENTER**

3:		
2:		
1:	'2*π*Y*PARC'	
	X	Y
	PARC	

Now enter the X and Y equations.

'2×SIN(T) **ENTER**

3:		
2:		
1:	'2*π*Y*PARC'	
	X	Y
	PARC	

'Y **STO**

3:		
2:		
1:	'2*π*Y*PARC'	
	X	Y
	PARC	


'2×COS(T) **ENTER**

3:		
2:		
1:	'2*π*Y*PARC'	
	X	Y
	PARC	

'X **STO**

3:		
2:		
1:	'2*π*Y*PARC'	
	X	Y
	PARC	

Key in the variable and limits of integration. With flag 35 cleared, π is evaluated to its numeric representation. The integration that follows requires a non-symbolic representation. Convert the parameters into a list.

T 0 π **ENTER**
3 **LIST** 

3:	
2:	'2* π *Y*PARC'
1:	{ T 0.00 3.14 }
+LIST LIST+ PUT GET PUTI GETI	

Key in the error bound and perform the integration.

.005 **ENTER**
J

3:	
2:	
1:	0.04 1.89E-4
+LIST LIST+ PUT GET PUTI GETI	

Note that 50.27 is $4\pi r^2$.

Purge the programs and variables created in this problem section.

{ 'X''Y''PARC' **PURGE**

Volume of Solid of Revolution: Method of Shells

This section demonstrates computation of the volume of a solid of revolution by the method of shells.

The method of shells requires evaluation of the integral

$$\int_a^b 2\pi x F(x) dx .$$

Example: Find the volume of the solid formed by revolving the curve

$$F(x) = e^{-x^2}$$

from $x = 0$ to $x = 3$ about the Y axis. Consider the behavior of the integral as the region of integration is extended.

Form an algebraic expression for the integrand including a general function $F(x)$.

CLEAR **<>**
'2*x*π*x*F' **ENTER**

4:	
3:	
2:	
1:	'2*π*x*F'

Store the integrand.

'SHEL' **STO**

4:	
3:	
2:	
1:	

Now enter the function. This must be a function of X as specified in the volume integrand.

'EXP(-X^2' **ENTER**

4:	
3:	
2:	
1:	'EXP(-X^2)'

Store the function by the name used in the SHEL program.

' F [STO]

4:
3:
2:
1:

Recall the expression to be integrated.

[USER] [SHEL]

3:
2:
1: '2*π*X*F'
F SHEL

Place the variable of integration and the limits on the stack.

{ X 0 3 [ENTER]

3:
2:
1: '2*π*X*F'
(X 0.00 3.00)
F SHEL

Specify the error bound of the integration.

.005 [ENTER]

3:
2:
1: '2*π*X*F'
(X 0.00 3.00)
0.01
F SHEL

Now integrate the function.

[ʃ]

3:
2:
1: 3.14
0.02
F SHEL

The result corresponds to π within the error specified.

Reset the display to show four digits.

[MODE] 4 [FIX]

3:
2:
1: 3.1403
0.0158
STD FIX SCI ENG DEG RAD

As expected, the accuracy is limited by the specification of two digits.

Perform the integration again, increasing the accuracy to produce four digits to the right of the decimal.

[USER] [SHEL]
{ X 0 3 [ENTER]
.00005 [ENTER]

[ʃ]

3: 0.0158
2: 3.1412
1: 0.0002
F SHEL

The desired accuracy was not achieved. By extending the region of integration, it may be possible to generate more digits of accuracy.

■ SHEL ■
{ X 0 4 [ENTER]
.00005 [ENTER]
[f]

3:	0.0002
2:	3.1416
1:	0.0002
F	SHEL

This is indeed π to four digits. This process does not prove that the integral, taken to infinity, converges to π . That proof requires an explicit solution to the integral. The curve that was specified is, of course, the "bell curve" used frequently in statistical analysis.

Purge the programs and variables used in this problem section.

{ 'SHEL' 'F' [PURGE]

Volume of Solids of Revolution: Method of Disks

This problem section computes volume of solids of revolution by the method of disks.

The method of disks requires evaluation of the integral

$$\int_a^b \pi f(x)^2 dx .$$

In general, for a given integral, the smaller the error bound the longer the integration will take. The appropriate choice of error bound depends on the problem being solved, but the method to reach a solution remains constant.

Example: Compute the volume of the solid formed by revolving the function $f(x) = x^2$ from 0 to 1 about the x axis.

Key in the first program for the general form of the integrand.

CLEAR **<>**
« → **X** 'π×**X**²' **ENTER**

4:
3:
2:
1: « → **X** 'π×**X**²' »

Store the program in DSK.

'**DSK**' **STO**

4:
3:
2:
1:

Key in the second program. This program puts the function and integration parameters in the appropriate form on the stack and calls DSK for the general form of the integrand. It then performs the volume computation.

« 'ER' **STO** 3.00 →**LIST**
SWAP **DSK** **SWAP** **ER** /
ENTER

2:
1: « 'ER' **STO** 3.0000
→**LIST** **SWAP** **DSK** **SWAP**
ER / »

Store the second program by the name DSKP.

'**DSKP**' **STO**

4:
3:
2:
1:

Now enter the function and integration data.

'X^2''X' 0 1 .005 **ENTER**

4:	'X'
3:	0.0000
2:	1.0000
1:	0.0050

Execute the program.

USER **DSKP**

3:	0.6283
2:	
1:	0.0031

The computed volume is .6283. The explicit solution to the integral is $\pi/5$.

For greater accuracy, increase the error bound as appropriate.

Purge the programs and variables created in this section.

{ 'DSK' 'DSKP' 'ER' **PURGE**

More Step-by-Step Solutions for Your HP-28S or HP-28C Calculator

These additional books offer a variety of examples and keystroke procedures to help set up your calculations the way *you* need them.

Practical routines show you how to use the built-in menus to solve problems more effectively, while easy-to-follow instructions help you create personalized menus.

Algebra and College Math (00028-90101)

- Solve algebraic problems: polynomial long division, function evaluation, simultaneous linear equations, quadratic equations, logarithms, polynomial equations, matrix determinants, and infinite series.
- Perform trigonometric calculations: graphs of functions, relations and identities, inverse functions, equations, and complex numbers.
- Solve analytic geometry problems: rectangular and polar coordinates, straight line, circle, parabola, ellipse, hyperbola, and parametric equations.

Vectors and Matrices (00028-90105)

- Perform general matrix operations: summation, multiplication, determinant, inverse, transpose, conjugate, and minor rank.
- Solve a system of linear equations.
- Calculate several important vector operations.
- Learn methods for calculating eigenvalues and eigenvectors.
- Perform the method of least squares and Markov Chain calculations.

Probability and Statistics (00028-90104)

- Set up a statistical matrix.
- Calculate basic statistics: mean, standard deviation, variance, covariance, correlation coefficient, sums of products, normalization, delta percent on paired data, moments, skewness, and kurtosis.

- Perform regression techniques: curve fitting, multiple linear, and polynomial regression.
- Compute several test statistics.

And Specifically for Your HP-28S...

Mathematical Applications (00028-90111)

- Find the area and all sides and angles of any plane triangle.
- Perform synthetic division on polynomials of arbitrary order.
- Calculate all the roots of a first, second, third and fourth degree polynomial, with real or complex coefficients.
- Solve first- and second-order differential equations.
- Convert the coordinates of two- or three-dimensional vectors between two coordinate systems, where one system is translated and/or rotated with respect to the other.
- Collect statistical data points, and fit curves to the data.

How to Order...

For the location and number of the U.S. dealer nearest you, call toll-free 1-800-752-0900. To order a book your dealer does not carry, call toll-free 1-800-538-8787 and refer to call code P270. Master Card, Visa, and American Express cards are welcome. For countries outside the U.S., contact your local Hewlett-Packard sales office.

Step-by-Step Solutions for Your HP-28S or HP-28C Calculator

Calculus contains a variety of examples and solutions to show how you can easily solve your technical problems.

■ Function Operations

Function Definition • Function Composition • Function Analysis
• Angle Between Two Lines • Angle Between Two Curves

■ Differential Calculus

Minimize Perimeter • Minimize Surface Area • Lines Tangent to a Circle • Implicit Differentiation With User-Defined Derivative
• Taylor Series Error Term • Tangent Lines and Taylor Series
• Normal Line • Implicit Functions

■ Integral Calculus

Integration and Free Falling Body • Double Integration • Area Between Two Curves • Arc Length • Surface Area • Arc Length of Parametric Equations • Surface Area of Parametric Equations
• Volume of A Solid of Revolution: Method of Shells and Method of Disks




**HEWLETT
PACKARD**

Reorder Number

00028-90102

00028-90134 English
Printed in Canada 6/88



0 88698 00022 9

Scan Copyright ©
The Museum of HP Calculators
www.hpmuseum.org

Original content used with permission.

Thank you for supporting the Museum of HP
Calculators by purchasing this Scan!

Please do not make copies of this scan or
make it available on file sharing services.