



HEWLETT  PACKARD

HP-55 mathematics programs



a comprehensive
guidebook:
74 common
programs
in such areas as

complex
arithmetic
and functions,
linear algebra,
trigonometry,
geometry,
business,
and others



\$10.00


Domestic
U.S.A. price
2/75

The program material contained herein is supplied without representation or warranty of any kind. Hewlett-Packard Company therefore assumes no responsibility and shall have no liability, consequential or otherwise, of any kind arising from the use of this program material or any part thereof.

Malcolm Lockwood Photography
A Standard in Quality Photographic Images
P.O. Box 2251 Fairbanks, AK 99707
(907) 474-0333

HEWLETT  PACKARD

HP-55
mathematics programs



Shown actual size.

INTRODUCTION

Material in *HP-55 Mathematics Programs* has been selected from the areas of complex variables, business, linear algebra, integration, interpolation, number theory, algebra, trigonometry, and analytical geometry.

Each program includes a general description, formulas used in the program solution, numerical examples, user instructions, program listings, and register allocations. The body of the book is arranged logically according to subject matter.

Some related individual programs were combined into one program when it seemed they might be useful together. In this way more programs could be included in the book. For individual program use it is possible for you to use only the portion of the combined program needed for your particular application.

In most cases the programs do not destroy stored data. Therefore, in order to run a different example only the data that is changed for the new example need be reentered.

We suggest that you first read the material explaining the Format of User Instructions, then use the programs. An understanding of the HP-55 Owner's Handbook is also required if, in addition, you wish to track the changes in the storage registers and stack registers on a step-by-step basis.

We hope you find *HP-55 Mathematics Programs* a useful tool for your mathematical work, and welcome your comments, requests and suggestions—these are our most important source of future user-oriented programs.

TABLE OF CONTENTS

Introduction

Format of User Instructions 5

COMPLEX VARIABLES

Complex Arithmetic $+$, $-$, $\times$, $\div$ 6

Complex Functions

$\ln z$, $\log_z w$, $\log_c z$ 9

$|z|$, z^2 , $1/z$, e^z , $\sqrt{z}$ 12

z^n , $z^{1/n}$ 14

z^w , $z^{1/w}$, c^z 16

Complex Trigonometric

$\sin z$, $\csc z$ 18

$\cos z$, $\sec z$ 20

$\tan z$, $\cot z$ 22

Complex Hyperbolic

$\sinh z$, $\operatorname{csch} z$ 24

$\cosh z$, $\operatorname{sech} z$ 26

$\tanh z$, $\operatorname{coth} z$ 28

Complex Inverse Trigonometric

$\sin^{-1} z$, $\csc^{-1} z$ 30

$\cos^{-1} z$, $\sec^{-1} z$ 32

$\tan^{-1} z$, $\cot^{-1} z$ 34

Complex Inverse Hyperbolic

$\sinh^{-1} z$, $\operatorname{csch}^{-1} z$ 36

$\cosh^{-1} z$, $\operatorname{sech}^{-1} z$ 38

$\tanh^{-1} z$, $\operatorname{coth}^{-1} z$ 40

Complex Polynomial Evaluation 42

BUSINESS

Compounded Amount 44

Direct Reduction Loan

 Interest Rate 48

 Payment, Present Value, Number of Time Periods 51

 Accumulated Interest, Remaining Balance 54

 Amortization Schedule 57

Sinking Fund

 Interest Rate 60

 Payment, Future Value, Number of Time Periods 63

Discounted Cash Flow Analysis	66
Depreciation Schedules	
Straight Line	68
Sum-of-the-Year's Digits	70
Variable Rate Declining Balance	72
Calendar Routines	
Day of the Week, Days between Two Dates	75
LINEAR ALGEBRA	
Determinant and Inverse of a 2×2 Matrix	78
Determinant of a 3×3 Matrix	80
3×3 Matrix Inversion	82
Vector Cross Product	86
Simultaneous Equations in Two Unknowns	88
Simultaneous Equations in Three Unknowns	91
2×2 Matrix Multiplication	94
Angle Between, Norm, and Dot Product of Vectors	96
INTEGRATION AND INTERPOLATION	
Sine Integral	100
Cosine Integral	102
Exponential Integral	104
Numerical Integration, Trapezoidal Rule	106
Numerical Integration, Simpson's Rule	108
Numerical Solution to Differential Equations	110
Linear Interpolation	112
NUMBER THEORY AND ALGEBRA	
Quadratic Equations	114
Synthetic Division	116
Factoring Integers and Determining Primes	118
Polynomial Evaluation	120
Number in Base b to a Number in Base 10	122
Number in Base 10 to Number in Base b	124
Newton's Method Solution to $f(x) = 0$	126
TRIGONOMETRY/ANALYTICAL GEOMETRY	
Trigonometric Functions ($\cot$, $\sec$, $\csc$, $\cot^{-1}$, $\sec^{-1}$, $\csc^{-1}$)	130
Versine, Coversine, Haversine, Exsecant	132
Hyperbolic Functions	134
Inverse Hyperbolic Functions	136
Polygons Inscribed in a Circle	138
Polygons Circumscribed about a Circle	140
Circle Determined by Three Points	142
Equally Spaced Points on a Circle	144

4 Table of Contents

Triangle Solutions

B, b, c	147
a, b, c	150
a, A, C	152
a, b, C	154
a, B, C	156

Area of a Triangle

a, b, c	158
a, b, C	160
a, B, C	162
$[(x_1, y_1), (x_2, y_2), (x_3, y_3)]$	164

Area of a Polygon 166

Spherical Triangle Solution

A, b, c	168
a, b, c	170
A, B, C	172

Translation and/or Rotation of Coordinate Axis 174

FORMAT OF USER INSTRUCTIONS

The completed User Instructions form is your guide to operating the programs in this book.

The form is composed of five columns. Reading from left to right, the first column, labeled STEP, gives the instruction step number.

The INSTRUCTIONS column gives instructions and comments concerning the operations to be performed. Steps are executed in sequential order except where the INSTRUCTIONS column directs otherwise.

Normally, the first instruction is "Enter program", which means to store the keystrokes of the program into memory (press **BSI** in RUN mode, switch to PRGM mode, then key in the program, switch back to RUN mode).

Repeated processes—used in most cases for a long string of input/output data—are outlined with a bold border together with a "Perform" instruction.

The INPUT DATA/UNITS column specifies the input data to be supplied, and the units of data if applicable.

The KEYS column specifies the keys to be pressed.  is the symbol used to denote the **ENTER** key. All other key designations are identical to those appearing on the HP-55. Ignore any blank positions in the KEYS column.

Some programs are sufficiently complex that they cannot be done in the 49 programming steps. However, they were sufficiently important to be included in this pac. In these cases the users must press additional keystrokes (other than program control keys) in order to get the answer. Those keys will also be shown in the KEYS column.

COMPLEX ARITHMETIC, +, -, ×, ÷

Let $a_1 + ib_1$ and $a_2 + ib_2$ be two complex numbers. The arithmetic operations +, -, ×, ÷ are defined as follows:

1. +, addition

$$(a_1 + ib_1) + (a_2 + ib_2) = (a_1 + a_2) + (b_1 + b_2)i$$

2. -, subtraction

$$(a_1 + ib_1) - (a_2 + ib_2) = (a_1 - a_2) + (b_1 - b_2)i$$

3. ×, multiplication

$$(a_1 + ib_1) \times (a_2 + ib_2) = r_1 r_2 e^{i(\theta_1 + \theta_2)}$$

4. ÷, division

$$\frac{(a_1 + ib_1)}{(a_2 + ib_2)} = \frac{r_1}{r_2} e^{i(\theta_1 - \theta_2)}, \quad a_2 + ib_2 \neq 0$$

where $r_1 e^{i\theta_1}$ is the polar representation of $a_1 + ib_1$ and $r_2 e^{i\theta_2}$ is the polar representation of $a_2 + ib_2$. In each case let the answer be $x + iy$.

After a calculation is finished x is stored in R_1 as well as the X-register and y is stored in R_2 as well as the Y-register. In this way arithmetic operations can be chained together.

DISPLAY			DISPLAY			REGISTERS
LINE	CODE	KEY ENTRY	LINE	CODE	KEY ENTRY	
00.			25.	34	RCL	R ₀
01.	42	CHS	26.	02	2	R ₁ a ₁ , x
02.	22	x $\overleftrightarrow{z}$ y	27.	34	RCL	R ₂ b ₁ , y
03.	42	CHS	28.	01	1	R ₃ Used
04.	22	x $\overleftrightarrow{z}$ y	29.	32	g	R ₄
05.	34	RCL	30.	00	R→P	R ₅
06.	01	1	31.	34	RCL	R ₆
07.	61	+	32.	03	3	R ₇
08.	22	x $\overleftrightarrow{z}$ y	33.	71	x	R ₈
09.	34	RCL	34.	33	STO	R ₉
10.	02	2	35.	03	3	R ₀₀
11.	61	+	36.	23	R↓	R ₀₁
12.	-43	GTO 43	37.	61	+	R ₀₂
13.	32	g	38.	34	RCL	R ₀₃
14.	00	R→P	39.	03	3	R ₀₄
15.	13	1/x	40.	31	f	R ₀₅
16.	22	x $\overleftrightarrow{z}$ y	41.	00	R←P	R ₀₆
17.	42	CHS	42.	22	x $\overleftrightarrow{z}$ y	R ₀₇
18.	22	x $\overleftrightarrow{z}$ y	43.	33	STO	R ₀₈
19.	-22	GTO 22	44.	02	2	R ₀₉
20.	32	g	45.	22	x $\overleftrightarrow{z}$ y	
21.	00	R→P	46.	33	STO	
22.	33	STO	47.	01	1	
23.	03	3	48.	-00	GTO 00	
24.	23	R↓	49.			

8 Complex Arithmetic, +, -, ×, ÷

Examples:

$$1. (3 + 4i) + (7.4 - 5.6i) = 10.40 - 1.60i$$

$$2. (3 + 4i) - (7.4 - 5.6i) = -4.40 + 9.60i$$

$$3. (3.1 + 4.6i) \times (5 - 12i) = 70.70 - 14.20i$$

$$4. \frac{(3 + 4i)}{7 - 2i} = .25 + .64i$$

$$5. \left[\frac{(3 + 4i) + (7.4 - 5.6i)}{7 - 2i} \right] [3.1 + 4.6i] = 3.61 + 7.16i$$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	For first calculation in chain						
	enter $a_1 + ib_1$	b_1	STO	2			
		a_1	STO	1			
3	Enter $a_2 + ib_2$	b_2	↑				
		a_2					
4	For Addition		GTO	0	5	R/S	x
	or						
	Subtraction		BST	R/S			x
	or						
	Multiplication		GTO	2	0	R/S	x
	or						
	Division		GTO	1	3	R/S	x
5	For imaginary part		$x \div y$				y
6	For next calculation in chain go						
	to step 3						
	or						
	For a new calculation, go to						
	step 2						

COMPLEX FUNCTIONS $\ln z$, $\log_z w$, $\log_c z$

Let $z = a_1 + ib_1$ and $w = a_2 + ib_2$ be complex numbers with polar representations $r_1 e^{i\theta_1}$ and $r_2 e^{i\theta_2}$ respectively. Also, let c be a positive real number. The formulas used to evaluate $\ln z$, $\log_z w$, and $\log_c z$ are as follows:

$$1. \quad \ln z = \ln r_1 + i\theta_1$$

$$2. \quad \log_z w = \frac{\ln w}{\ln z} = \frac{r_3}{r_4} e^{i(\theta_3 - \theta_4)} \quad z \neq 0$$

$$\text{where } \ln r_2 + i\theta_2 = r_3 e^{i\theta_3}$$

$$\text{and } \ln r_1 + i\theta_1 = r_4 e^{i\theta_4}$$

$$3. \quad \log_c z = \frac{\ln z}{\ln c} = \frac{\ln r_1}{\ln c} + \frac{\theta_1}{\ln c} i \quad c > 0$$

In each case let the solution be $x + iy$. The calculator must be in RADIAN mode for all three functions.

10 Complex Functions $\ln z$, $\log_z w$, $\log_c z$

DISPLAY			KEY ENTRY	DISPLAY		KEY ENTRY	REGISTERS
LINE	CODE			LINE	CODE		
00.				25.	33	STO	R_0
01.	32	g		26.	03	3	$R_1 \ a_1, c$
02.	00	$R \rightarrow P$		27.	23	$R \downarrow$	$R_2 \ b_1$
03.	31	f		28.	51	—	$R_3 \ r_3/r_4$
04.	22	$\ln$		29.	42	CHS	R_4
05.	-00	GTO 00		30.	34	RCL	R_5
06.	32	g		31.	03	3	R_6
07.	00	$R \rightarrow P$		32.	31	f	R_7
08.	31	f		33.	00	$R \leftarrow P$	R_8
09.	22	$\ln$		34.	-00	GTO 00	R_9
10.	32	g		35.	32	g	R_{10}
11.	00	$R \rightarrow P$		36.	00	$R \rightarrow P$	R_{11}
12.	34	RCL		37.	31	f	R_{12}
13.	02	2		38.	22	$\ln$	R_{13}
14.	34	RCL		39.	34	RCL	R_{14}
15.	01	1		40.	01	1	R_{15}
16.	32	g		41.	31	f	R_{16}
17.	00	$R \rightarrow P$		42.	22	$\ln$	R_{17}
18.	31	f		43.	81	$\div$	R_{18}
19.	22	$\ln$		44.	22	$x \leftrightarrow y$	R_{19}
20.	32	g		45.	31	f	
21.	00	$R \rightarrow P$		46.	34	LAST X	
22.	22	$x \leftrightarrow y$		47.	81	$\div$	
23.	23	$R \downarrow$		48.	22	$x \leftrightarrow y$	
24.	81	$\div$		49.	-00	GTO 00	

Examples:

- $\ln(1 + i) = .35 + .79i$
- $\log_{(1+i)}(1.49 + 4.13i) = 2.00 - 1.00i$
- $\log_2(-7.46 + 2.89i) = 3.00 + 4.00i$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Set in radian mode		f	RAD			
3	For $\ln z$, enter z	b_1	$\uparrow$				
		a_1	BST	R/S			x
			$x \div y$				y
	or						
3	For $\log_z w$, store z	b_1	STO	2			
		a_1	STO	1			
4	Enter w	b_2	$\uparrow$				
		a_2	GTO	0	6	R/S	x
			$x \div y$				y
	or						
3	For $\log_c z$, store c	c	STO	1			
4	Enter z^c	b_1	$\uparrow$				
		a_1	GTO	3	5	R/S	x
			$x \div y$				y

COMPLEX FUNCTIONS $|z|$, z^2 , $1/z$, e^z , $\sqrt{z}$

A complex number $z = a + ib$ has polar representation $re^{i\theta}$. The formulas used to evaluate the given functions are as follows:

- $|z| = r$
- $z^2 = r^2 e^{i2\theta}$
- $1/z = \frac{1}{r} e^{-i\theta}$, $z \neq 0$
- $e^z = e^a e^{ib}$
- $\sqrt{z} = \pm (\sqrt{r} e^{i\theta/2}) = \pm(x + iy)$

The answer is represented by $x + iy$. For e^z the calculator must be in RADIAN mode.

DISPLAY		KEY ENTRY	DISPLAY		KEY ENTRY	REGISTERS
LINE	CODE		LINE	CODE		
00.			25.	22	e^x	R_0
01.	32	g	26.	31	f	R_1
02.	00	R→P	27.	00	R←P	R_2
03.	-00	GTO 00	28.	-00	GTO 00	R_3
04.	32	g	29.	32	g	R_4
05.	00	R→P	30.	00	R→P	R_5
06.	41	↑	31.	31	f	R_6
07.	71	x	32.	42	$\sqrt{x}$	R_7
08.	22	$x \rightarrow y$	33.	22	$x \rightarrow y$	R_8
09.	41	↑	34.	02	2	R_9
10.	61	+	35.	81	÷	R_{00}
11.	22	$x \rightarrow y$	36.	22	$x \rightarrow y$	R_{01}
12.	31	f	37.	31	f	R_{02}
13.	00	R←P	38.	00	R←P	R_{03}
14.	-00	GTO 00	39.	-00	GTO 00	R_{04}
15.	32	g	40.			R_{05}
16.	00	R→P	41.			R_{06}
17.	13	$1/x$	42.			R_{07}
18.	22	$x \rightarrow y$	43.			R_{08}
19.	42	CHS	44.			R_{09}
20.	22	$x \rightarrow y$	45.			
21.	31	f	46.			
22.	00	R←P	47.			
23.	-00	GTO 00	48.			
24.	32	g	49.			

Examples:

- $|3 + 4i| = 5.00$
- $(7 - 2i)^2 = 45.00 - 28.00i$
- $\frac{1}{2 + 3i} = .15 - .23i$
- $e^{(3+4i)} = -13.13 - 15.20i$
- $\sqrt{7 + 6i} = \pm(2.85 + 1.05i)$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Set to radian mode (only necessary for e^z)		f	RAD			
3	Enter z	b	↑				
		a					
4	For $ z $		BST	R/S			$ z $
	or						
	z^2		GTO	0	4	R/S	x
			$x \rightarrow y$				y
	or						
	$1/z$		GTO	1	5	R/S	x
			$x \rightarrow y$				y
	or						
	e^z		GTO	2	4	R/S	x
			$x \rightarrow y$				y
	or						
	$\sqrt{z}$ (one root only)		GTO	2	9	R/S	x
			$x \rightarrow y$				y

COMPLEX FUNCTIONS $z^n, z^{1/n}$

A complex number $z = a + ib$ has polar representation $re^{i\theta}$. The formulas used to evaluate z^n and $z^{1/n}$ where n is a positive integer are:

$$1. z^n = r^n e^{in\theta}$$

$$2. z^{1/n} = r^{1/n} \left[\cos\left(\frac{\theta + ck}{n}\right) + i \sin\left(\frac{\theta + ck}{n}\right) \right]$$

where $c = 4 \sin^{-1} 1 = 360^\circ = 2\pi$ radians = 400 grads

and $k = 0, 1, \dots, n-1$

The solution to z^n is represented by $x + iy$ and the n solutions to $z^{1/n}$ are represented by $x_k + iy_k$.

DISPLAY			KEY ENTRY			REGISTERS
LINE	CODE		LINE	CODE		
00.			25.	71	x	R ₀
01.	34	RCL	26.	33	STO	R ₁ $n, 1/n$
02.	01	1	27.	04	4	R ₂ c/n
03.	13	$1/x$	28.	22	$x \leftrightarrow y$	R ₃ $R^{1/n}$
04.	33	STO	29.	31	f	R ₄ θ/n
05.	01	1	30.	00	R←P	R ₅
06.	01	1	31.	84	R/S	R ₆
07.	32	g	32.	34	RCL	R ₇
08.	12	$\sin^{-1}$	33.	04	4	R ₈
09.	71	x	34.	34	RCL	R ₉
10.	04	4	35.	02	2	R ₀₀
11.	71	x	36.	61	+	R ₀₁
12.	33	STO	37.	33	STO	R ₀₂
13.	02	2	38.	04	4	R ₀₃
14.	23	R↓	39.	34	RCL	R ₀₄
15.	32	g	40.	03	3	R ₀₅
16.	00	R→P	41.	31	f	R ₀₆
17.	34	RCL	42.	00	R←P	R ₀₇
18.	01	1	43.	-31	GTO 31	R ₀₈
19.	12	y^x	44.			R ₀₉
20.	33	STO	45.			
21.	03	3	46.			
22.	22	$x \leftrightarrow y$	47.			
23.	34	RCL	48.			
24.	01	1	49.			

Examples:

1. $(3 + 4.5i)^5 = 926.44 - 4533.47i$

2. $(5 + 3i)^{1/3} = \begin{cases} 1.77 + .32i \\ -1.16 + 1.37i \\ -.61 - 1.69i \end{cases}$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Store n	n	STO	1			
3	Enter z	b	↑				
		a					
4	For z^n		GTO	1	5	R/S	x
			$x \leftrightarrow y$				y
	or						
	$(a + ib)^{1/n}$ (primary root)		BST	R/S			x_0
			$x \leftrightarrow y$				y_0
5	Then for other n^{th} roots		R/S				x_i
			$x \leftrightarrow y$				y_i

COMPLEX FUNCTIONS $z^w, z^{1/w}, c^z$

Let $z = a_1 + ib_1$ and $w = a_2 + ib_2$ be complex numbers with polar representations $r_1 e^{i\theta_1}$ and $r_2 e^{i\theta_2}$ respectively. Also, let c be a positive real number. The formulas used to evaluate $z^w, z^{1/w}$, and c^z are as follows:

$$1. \quad z^w = e^{w \ln z} = e^{(r_2 e^{i\theta_2}) (\ln r_1 + i\theta_1)} = e^{r_2 r_3 e^{i(\theta_2 + \theta_3)}} = e^{a_4} e^{ib_4}$$

$$\text{where } \ln z = \ln r_1 + i\theta_1 = r_3 e^{i\theta_3}$$

$$\text{and } a_4 + ib_4 = r_2 r_3 e^{i(\theta_2 + \theta_3)}$$

$$2. \quad z^{1/w} = z^{w'}$$

$$\text{where } w' = \frac{1}{r_2} e^{-i\theta_2}$$

$$3. \quad c^z = e^{z \ln c} = e^{a_1 \ln c} e^{ib_1 \ln c}$$

Let the solution in each case be $x + iy$. The calculator must be set in RADIAN mode for all three cases.

DISPLAY			KEY ENTRY			REGISTERS
LINE	CODE		LINE	CODE		
00.			25.	23	R↓	R ₀
01.	32	g	26.	61	+	R ₁ a ₁ , c
02.	00	R→P	27.	34	RCL	R ₂ b ₁
03.	13	1/x	28.	03	3	R ₃ r ₁ r ₃
04.	22	x↔y	29.	31	f	R ₄
05.	42	CHS	30.	00	R←P	R ₅
06.	22	x↔y	31.	32	g	R ₆
07.	-10	GTO 10	32.	22	e ^x	R ₇
08.	32	g	33.	31	f	R ₈
09.	00	R→P	34.	00	R←P	R ₉
10.	34	RCL	35.	-00	GTO 00	R ₀₀
11.	02	2	36.	22	x↔y	R ₀₁
12.	34	RCL	37.	34	RCL	R ₀₂
13.	01	1	38.	01	1	R ₀₃
14.	32	g	39.	31	f	R ₀₄
15.	00	R→P	40.	22	ln	R ₀₅
16.	31	f	41.	71	x	R ₀₆
17.	22	ln	42.	22	x↔y	R ₀₇
18.	32	g	43.	31	f	R ₀₈
19.	00	R→P	44.	34	LAST X	R ₀₉
20.	22	x↔y	45.	71	x	
21.	23	R↓	46.	32	g	
22.	71	x	47.	22	e ^x	
23.	33	STO	48.	31	f	
24.	03	3	49.	00	R←P	

Examples:

- $(1 + i)^{(2-i)} = 1.49 + 4.13i$
- $(1.49 + 4.13i)^{1/(2-i)} = 1.00 + 1.00i$
- $2^{(3+4i)} = -7.46 + 2.89i$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Set to radian mode		f	RAD			
3	For z^w , store z	b_1	STO	2			
		a_1	STO	1			
4	Enter w	b_2	↑				
		a_2	GTO	0	8	R/S	x
			$x \leftrightarrow y$				y
	or						
3	For $z^{1/w}$, store z	b_1	STO	2			
		a_1	STO	1			
4	Enter w	b_2	↑				
		a_2	BST	R/S			x
			$x \leftrightarrow y$				y
	or						
3	For c^z , store c	c	STO	1			
4	Enter z	b_1	↑				
		a_1	GTO	3	6	R/S	x
			$x \leftrightarrow y$				y

COMPLEX TRIGONOMETRIC $\sin z$, $\csc z$

Let $z = a + ib$ be a complex number. The functions $\sin z$ and $\csc z$ are evaluated by the following formulas:

$$1. \sin z = \sin a \cosh b + i \cos a \sinh b$$

$$2. \csc z = \frac{1}{\sin z} \quad z \neq 0, \pm \pi, \pm 2\pi, \dots$$

Let the solutions be $x_1 + iy_1$ and $x_2 + iy_2$ respectively.

The calculator must be set in RADIAN mode.

DISPLAY			KEY ENTRY			REGISTERS
LINE	CODE	KEY ENTRY	LINE	CODE	KEY ENTRY	
00.			25.	13	$1/x$	R_0
01.	33	STO	26.	51	—	R_1 a
02.	01	1	27.	02	2	R_2 e^b
03.	31	f	28.	81	$\div$	R_3 x_1
04.	12	sin	29.	71	x	R_4
05.	22	$x \rightarrow y$	30.	34	RCL	R_5
06.	32	g	31.	03	3	R_6
07.	22	e^x	32.	84	R/S	R_7
08.	33	STO	33.	32	g	R_8
09.	02	2	34.	00	$R \rightarrow P$	R_9
10.	41	$\uparrow$	35.	13	$1/x$	R_{10}
11.	13	$1/x$	36.	22	$x \rightarrow y$	R_{11}
12.	61	+	37.	42	CHS	R_{12}
13.	02	2	38.	22	$x \rightarrow y$	R_{13}
14.	81	$\div$	39.	31	f	R_{14}
15.	71	x	40.	00	$R \rightarrow P$	R_{15}
16.	33	STO	41.	-00	GTO 00	R_{16}
17.	03	3	42.			R_{17}
18.	34	RCL	43.			R_{18}
19.	01	1	44.			R_{19}
20.	31	f	45.			
21.	13	cos	46.			
22.	34	RCL	47.			
23.	02	2	48.			
24.	41	$\uparrow$	49.			

Examples:

- 1. $\sin (2 + 3i) = 9.15 - 4.17i$
- 2. $\csc (2 + 3i) = .09 + .04i$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Set to radian mode		f	RAD			
3	For $\sin z$, enter z	b	↑				
		a	BST	R/S			x_1
			$x \leftrightarrow y$				y_1
	or						
3	For $\csc z$, enter z	b	↑				
		a	BST	R/S			x_1
4	(Do not change the contents of the X and Y registers at this point.)						
			R/S				x_2
			$x \leftrightarrow y$				y_2

COMPLEX TRIGONOMETRIC $\cos z$, $\sec z$

Let $z = a + ib$ be a complex number. The functions $\cos z$ and $\sec z$ are evaluated by the following formulas:

$$1. \cos z = \cos a \cosh b - i \sin a \sinh b$$

$$2. \sec z = \frac{1}{\cos z} \quad z \neq \pm \frac{\pi}{2}, \pm \frac{3\pi}{2}, \pm \frac{5\pi}{2}, \dots$$

Let the solutions be $x_1 + iy_1$ and $x_2 + iy_2$ respectively.

The calculator must be set in RADIAN mode.

DISPLAY		KEY ENTRY	DISPLAY		KEY ENTRY	REGISTERS
LINE	CODE		LINE	CODE		
00.			25.	13	$1/x$	R_0
01.	33	STO	26.	51	—	R_1 a
02.	01	1	27.	02	2	R_2 e^b
03.	31	f	28.	81	$\div$	R_3 x_1
04.	13	cos	29.	71	x	R_4
05.	22	$x \rightarrow y$	30.	42	CHS	R_5
06.	32	g	31.	34	RCL	R_6
07.	22	e^x	32.	03	3	R_7
08.	33	STO	33.	84	R/S	R_8
09.	02	2	34.	32	g	R_9
10.	41	$\uparrow$	35.	00	$R \rightarrow P$	R_{00}
11.	13	$1/x$	36.	13	$1/x$	R_{01}
12.	61	+	37.	22	$x \rightarrow y$	R_{02}
13.	02	2	38.	42	CHS	R_{03}
14.	81	$\div$	39.	22	$x \rightarrow y$	R_{04}
15.	71	x	40.	31	f	R_{05}
16.	33	STO	41.	00	$R \leftarrow P$	R_{06}
17.	03	3	42.	-00	GTO 00	R_{07}
18.	34	RCL	43.			R_{08}
19.	01	1	44.			R_{09}
20.	31	f	45.			
21.	12	sin	46.			
22.	34	RCL	47.			
23.	02	2	48.			
24.	41	$\uparrow$	49.			

Examples:

1. $\cos(2 + 3i) = -4.19 - 9.11i$

2. $\sec(2 + 3i) = -.04 + .09i$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Set to radian mode		f	RAD			
3	For $\cos z$, enter z	b	↑				
		a	BST	R/S			x_1
			$x \leftrightarrow y$				y_1
	or						
3	For $\sec z$, enter z	b	↑				
		a	BST	R/S			x_1
	(Do not change the contents of the X and Y registers at this point.)						
			R/S				x_2
			$x \leftrightarrow y$				y_2

COMPLEX TRIGONOMETRIC $\tan z$, $\cot z$

Let $z = a + ib$ be a complex number. The functions $\tan z$ and $\cot z$ are evaluated by the following formulas:

$$1. \tan z = \frac{\sin 2a + i \sinh 2b}{\cos 2a + \cosh 2b} \quad z \neq \pm \frac{\pi}{2}, \pm \frac{3\pi}{2}, \dots$$

$$2. \cot z = \frac{1}{\tan z} \quad z \neq 0, \pm \frac{\pi}{2}, \pm \pi, \pm \frac{3\pi}{2}, \dots$$

Let the solutions be $x_1 + iy_1$ and $x_2 + iy_2$ respectively.

The calculator must be set in RADIAN mode.

If $z = \pm \frac{\pi}{2}, \pm \frac{3\pi}{2}, \dots$ then $\cot z = 0$.

DISPLAY		KEY ENTRY	DISPLAY		KEY ENTRY	REGISTERS
LINE	CODE		LINE	CODE		
00.			25.	13	$1/x$	R_0
01.	02	2	26.	51	—	$R_1 \ 2a$
02.	71	x	27.	02	2	$R_2 \ e^{2b}$
03.	33	STO	28.	81	$\div$	$R_3 \ \cos 2a + \cosh 2b$
04.	01	1	29.	22	$x \rightarrow y$	R_4
05.	31	f	30.	81	$\div$	R_5
06.	13	cos	31.	34	RCL	R_6
07.	22	$x \rightarrow y$	32.	01	1	R_7
08.	02	2	33.	31	f	R_8
09.	71	x	34.	12	sin	R_9
10.	32	g	35.	34	RCL	R_{10}
11.	22	e^x	36.	03	3	R_{11}
12.	33	STO	37.	81	$\div$	R_{12}
13.	02	2	38.	84	R/S	R_{13}
14.	41	$\uparrow$	39.	32	g	R_{14}
15.	13	$1/x$	40.	00	$R \rightarrow P$	R_{15}
16.	61	+	41.	13	$1/x$	R_{16}
17.	02	2	42.	22	$x \rightarrow y$	R_{17}
18.	81	$\div$	43.	42	CHS	R_{18}
19.	61	+	44.	22	$x \rightarrow y$	R_{19}
20.	33	STO	45.	31	f	
21.	03	3	46.	00	$R \leftarrow P$	
22.	34	RCL	47.	-00	GTO 00	
23.	02	2	48.			
24.	41	$\uparrow$	49.			

Examples:

- $\tan(4 + .01i) = 1.16 + .02i$
- $\cot(4 + .01i) = .86 - .02i$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Set to radian mode		f	RAD			
3	For $\tan z$, enter z	b	↑				
		a	BST	R/S			x_1
			$x \leftrightarrow y$				y_1
	or						
3	For $\cot z$, enter z	b	↑				
		a	BST	R/S			x_1
	(Do not change the contents of the X and Y registers at this point.)						
			R/S				x_2
			$x \leftrightarrow y$				y_2

COMPLEX HYPERBOLIC $\sinh z$, $\cosh z$

Let $z = a + ib$ be a complex number. The functions $\sinh z$ and $\cosh z$ are evaluated by the following formulas:

$$1. \sinh z = -i \sin i z = \cos b \sinh a + i \sin b \cosh a$$

$$2. \cosh z = \frac{1}{\sinh z} \quad z \neq 0, \pm i\pi, \pm 2i\pi, \dots$$

Let the solutions be $x_1 + iy_1$ and $x_2 + iy_2$ respectively.

The calculator must be set in RADIAN mode.

DISPLAY			KEY ENTRY	DISPLAY			KEY ENTRY	REGISTERS
LINE	CODE			LINE	CODE			
00.				25.	31	f		R ₀
01.	32	g		26.	13	cos		R ₁ e ^a
02.	22	e ^x		27.	71	x		R ₂ b
03.	41	↑		28.	84	R/S		R ₃
04.	33	STO		29.	32	g		R ₄
05.	01	1		30.	00	R→P		R ₅
06.	13	1/x		31.	13	1/x		R ₆
07.	61	+		32.	22	x↔y		R ₇
08.	02	2		33.	42	CHS		R ₈
09.	81	÷		34.	22	x↔y		R ₉
10.	22	x↔y		35.	31	f		R ₀₀
11.	33	STO		36.	00	R←P		R ₀₁
12.	02	2		37.	-00	GTO 00		R ₀₂
13.	31	f		38.				R ₀₃
14.	12	sin		39.				R ₀₄
15.	71	x		40.				R ₀₅
16.	34	RCL		41.				R ₀₆
17.	01	1		42.				R ₀₇
18.	41	↑		43.				R ₀₈
19.	13	1/x		44.				R ₀₉
20.	51	-		45.				
21.	02	2		46.				
22.	81	÷		47.				
23.	34	RCL		48.				
24.	02	2		49.				

Examples:

1. $\sinh (3 - 2i) = -4.17 - 9.15i$

2. $\cosh (1 + 2i) = -.22 - .64i$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Set to radian mode		f	RAD			
3	For $\sinh z$, enter z	b	↑				
		a	BST	R/S			x_1
			$x \div y$				y_1
	or						
3	For $\cosh z$, enter z	b	↑				
		a	BST	R/S			x_1
	(Do not change the contents of the X and Y registers at this point.)						
			R/S				x_2
			$x \div y$				y_2

COMPLEX HYPERBOLIC cosh z, sech z

Let $z = a + ib$ be a complex number. The functions $\cosh z$ and $\operatorname{sech} z$ are evaluated by the following formulas:

$$1. \cosh z = \cos iz = \cos b \cosh a + i \sin b \sinh a$$

$$2. \operatorname{sech} z = \frac{1}{\cosh z} \quad z \neq \pm \frac{i\pi}{2}, \pm \frac{3i\pi}{2}, \pm \frac{5i\pi}{2}, \dots$$

Let the solutions be $x_1 + iy_1$ and $x_2 + iy_2$ respectively.

The calculator must be set in RADIAN mode.

DISPLAY			KEY ENTRY			REGISTERS
LINE	CODE		LINE	CODE		
00.			25.	31	f	R ₀
01.	32	g	26.	13	cos	R ₁ e ^a
02.	22	e ^x	27.	71	x	R ₂ b
03.	33	STO	28.	84	R/S	R ₃
04.	01	1	29.	32	g	R ₄
05.	41	↑	30.	00	R→P	R ₅
06.	13	1/x	31.	13	1/x	R ₆
07.	51	—	32.	22	x↔y	R ₇
08.	02	2	33.	42	CHS	R ₈
09.	81	÷	34.	22	x↔y	R ₉
10.	22	x↔y	35.	31	f	R ₁₀
11.	33	STO	36.	00	R←P	R ₁₁
12.	02	2	37.	-00	GTO 00	R ₁₂
13.	31	f	38.			R ₁₃
14.	12	sin	39.			R ₁₄
15.	71	x	40.			R ₁₅
16.	34	RCL	41.			R ₁₆
17.	01	1	42.			R ₁₇
18.	41	↑	43.			R ₁₈
19.	13	1/x	44.			R ₁₉
20.	61	+	45.			
21.	02	2	46.			
22.	81	÷	47.			
23.	34	RCL	48.			
24.	02	2	49.			

Examples:

- $\cosh (1 + 2i) = -.64 + 1.07i$
- $\operatorname{sech} (1 + 2i) = -.41 - .69i$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Set to radian mode		f	RAD			
3	For cosh z, enter z	b	↑				
		a	BST	R/S			x_1
			$x \leftrightarrow y$				y_1
	or						
3	For sech z, enter z	b	↑				
		a	BST	R/S			x_1
	(Do not change the contents of the X and Y registers at this point.)						
			R/S				x_2
			$x \leftrightarrow y$				y_2

COMPLEX HYPERBOLIC $\tanh z$, $\coth z$

Let $z = a + ib$ be a complex number. The functions $\tanh z$ and $\coth z$ are evaluated by the following formulas:

$$1. \quad \tanh z = -i \tan iz = \frac{\sinh 2a + i \sin 2b}{\cosh 2a + \cos 2b} \quad z \neq \pm \frac{i\pi}{2}, \pm \frac{3i\pi}{2}, \dots$$

$$2. \quad \coth z = \frac{1}{\tanh z} \quad z \neq \pm \frac{i\pi}{2}, \pm \frac{3i\pi}{2}, \dots \text{ or } z \neq 0, \pm i\pi, \pm 2i\pi$$

Let the solutions be $x_1 + iy_1$ and $x_2 + iy_2$ respectively.

The calculator must be set in RADIAN mode.

$$\text{If } z = \frac{i\pi}{2}, \frac{3i\pi}{2}, \dots \coth z = 0$$

DISPLAY			KEY ENTRY	DISPLAY			KEY ENTRY	REGISTERS
LINE	CODE			LINE	CODE			
00.				25.	12	sin		R ₀
01.	02	2		26.	22	$x \leftrightarrow y$		R ₁ e^{2a}
02.	71	x		27.	81	$\div$		R ₂ 2b
03.	32	g		28.	34	RCL		R ₃ $\cosh 2a + \cos 2b$
04.	22	e^x		29.	01	1		R ₄
05.	41	$\uparrow$		30.	41	$\uparrow$		R ₅
06.	33	STO		31.	13	$1/x$		R ₆
07.	01	1		32.	51	—		R ₇
08.	13	$1/x$		33.	02	2		R ₈
09.	61	+		34.	81	$\div$		R ₉
10.	02	2		35.	34	RCL		R ₀₀
11.	81	$\div$		36.	03	3		R ₀₁
12.	22	$x \leftrightarrow y$		37.	81	$\div$		R ₀₂
13.	02	2		38.	84	R/S		R ₀₃
14.	71	x		39.	32	g		R ₀₄
15.	33	STO		40.	00	R $\rightarrow$ P		R ₀₅
16.	02	2		41.	13	$1/x$		R ₀₆
17.	31	f		42.	22	$x \leftrightarrow y$		R ₀₇
18.	13	cos		43.	42	CHS		R ₀₈
19.	61	+		44.	22	$x \leftrightarrow y$		R ₀₉
20.	33	STO		45.	31	f		
21.	03	3		46.	00	R $\leftarrow$ P		
22.	34	RCL		47.	-00	GTO 00		
23.	02	2		48.				
24.	31	f		49.				

Examples:

1. $\tanh(1 + 2i) = 1.17 - .24i$

2. $\coth(1 + 2i) = .82 + .17i$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Set to radian mode		f	RAD			
3	For $\tanh z$, enter z	b	↑				
		a	BST	R/S			x_1
			$x \leftrightarrow y$				y_1
	or						
3	For $\coth z$, enter z	b	↑				
		a	BST	R/S			x_1
	(Do not change the contents of the X and Y registers at this point.)						
			R/S				x_2
			$x \leftrightarrow y$				y_2

COMPLEX INVERSE TRIGONOMETRIC

$\sin^{-1} z$, $\csc^{-1} z$

Let $z = a + ib$ be a complex number. The functions $\sin^{-1} z$ (arc sin) and $\csc^{-1} z$ (arc csc) are evaluated by the following formulas:

$$1. \sin^{-1} z = k\pi + (-1)^k \sin^{-1} \beta + (-1)^k i \operatorname{sgn}(b) \ln [\alpha + (\alpha^2 - 1)^{1/2}]$$

$$\text{where } \alpha = \frac{1}{2} \sqrt{(a+1)^2 + b^2} + \frac{1}{2} \sqrt{(a-1)^2 + b^2}$$

$$\beta = \frac{1}{2} \sqrt{(a+1)^2 + b^2} - \frac{1}{2} \sqrt{(a-1)^2 + b^2}$$

$$\operatorname{sgn}(b) = b / \sqrt{b^2} = \begin{cases} 1 & \text{if } b > 0 \\ -1 & \text{if } b < 0 \end{cases}$$

$$k = 0, \pm 1, \pm 2, \dots$$

$$2. \csc^{-1} z = \sin^{-1} \left[\frac{1}{z} \right] \quad z \neq 0$$

k is assumed to be zero for this program.

Program does not work for $b = 0$ but the calculator functions can be used instead.

Let the solutions be $x_1 + iy_1$ and $x_2 + iy_2$ respectively.

The calculator must be in RADIAN mode and all angles are assumed to be in radians.

DISPLAY			KEY ENTRY			REGISTERS
LINE	CODE	KEY ENTRY	LINE	CODE	KEY ENTRY	
00.			25.	71	x	R ₀
01.	22	$x \leftrightarrow y$	26.	01	1	R ₁
02.	33	STO	27.	51	—	R ₂ b
03.	02	2	28.	31	f	R ₃ $\sqrt{(a-1)^2 + b^2}$
04.	22	$x \leftrightarrow y$	29.	42	$\sqrt{x}$	R ₄
05.	01	1	30.	61	+	R ₅
06.	51	—	31.	31	f	R ₆
07.	32	g	32.	22	ln	R ₇
08.	00	R→P	33.	34	RCL	R ₈
09.	33	STO	34.	02	2	R ₉
10.	03	3	35.	41	↑	R ₀₀
11.	31	f	36.	32	g	R ₀₁
12.	00	R←P	37.	42	x^2	R ₀₂
13.	02	2	38.	31	f	R ₀₃
14.	61	+	39.	42	$\sqrt{x}$	R ₀₄
15.	32	g	40.	81	÷	R ₀₅
16.	00	R→P	41.	71	x	R ₀₆
17.	34	RCL	42.	22	$x \leftrightarrow y$	R ₀₇
18.	03	3	43.	34	RCL	R ₀₈
19.	61	+	44.	03	3	R ₀₉
20.	02	2	45.	51	—	
21.	81	÷	46.	32	g	
22.	41	↑	47.	13	$\sin^{-1}$	
23.	41	↑	48.	—00	GTO 00	
24.	41	↑	49.			

Examples:

- $\sin^{-1} (5 + 8i) = .56 + 2.94i$
- $\csc^{-1} (5 + 8i) = .06 - .09i$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Set to radian mode		f	RAD			
3	For $\sin^{-1} z$, enter z	b	↑				
		a	BST	R/S			x ₁
			$x \leftrightarrow y$				y ₁
	or						
3	For $\csc^{-1} z$, enter z						
	and calculate 1/z	b	↑				
		a	g	R→P	1/x	$x \leftrightarrow y$	
			CHS	$x \leftrightarrow y$	f	R←P	
			BST	R/S			x ₂
			$x \leftrightarrow y$				y ₂

COMPLEX INVERSE TRIGONOMETRIC $\cos^{-1} z$, $\sec^{-1} z$

Let $z = a + ib$ be a complex number. The functions $\cos^{-1} z$ (arc cos) and $\sec^{-1} z$ (arc sec) are evaluated by the following formulas:

$$1. \quad \cos^{-1} z = 2k\pi \pm \left\{ \cos^{-1} \beta - i \operatorname{sgn}(b) \ln [\alpha + (\alpha^2 - 1)^{1/2}] \right\}$$

$$\text{where } \alpha = \frac{1}{2} \sqrt{(a+1)^2 + b^2} + \frac{1}{2} \sqrt{(a-1)^2 + b^2}$$

$$\beta = \frac{1}{2} \sqrt{(a+1)^2 + b^2} - \frac{1}{2} \sqrt{(a-1)^2 + b^2}$$

$$\operatorname{sgn}(b) = b/\sqrt{b^2} = \begin{cases} 1 & \text{if } b > 0 \\ -1 & \text{if } b < 0 \end{cases}$$

$$k = 0, \pm 1, \pm 2, \dots$$

$$2. \quad \sec^{-1} z = \cos^{-1} \left(\frac{1}{z} \right), \quad z \neq 0$$

k is assumed to be zero for this program.

Program does not work for $b = 0$ but the inbuilt calculator functions can be used instead.

Let the solutions be $x_1 + iy_1$ and $x_2 + iy_2$ respectively.

The calculator must be in RADIAN mode and all angles are assumed to be in radians.

DISPLAY		KEY ENTRY	DISPLAY		KEY ENTRY	REGISTERS
LINE	CODE		LINE	CODE		
00.			25.	71	x	R_0
01.	22	$x \leftrightarrow y$	26.	01	1	R_1
02.	33	STO	27.	51	—	R_2 b
03.	02	2	28.	31	f	$R_3 \sqrt{(a-1)^2 + b^2}$
04.	22	$x \leftrightarrow y$	29.	42	$\sqrt{x}$	R_4
05.	01	1	30.	61	+	R_5
06.	51	—	31.	31	f	R_6
07.	32	g	32.	22	ln	R_7
08.	00	R→P	33.	42	CHS	R_8
09.	33	STO	34.	34	RCL	R_9
10.	03	3	35.	02	2	R_{10}
11.	31	f	36.	41	↑	R_{11}
12.	00	R←P	37.	32	g	R_{12}
13.	02	2	38.	42	x^2	R_{13}
14.	61	+	39.	31	f	R_{14}
15.	32	g	40.	42	$\sqrt{x}$	R_{15}
16.	00	R→P	41.	81	÷	R_{16}
17.	34	RCL	42.	71	x	R_{17}
18.	03	3	43.	22	$x \leftrightarrow y$	R_{18}
19.	61	+	44.	34	RCL	R_{19}
20.	02	2	45.	03	3	
21.	81	÷	46.	51	—	
22.	41	↑	47.	32	g	
23.	41	↑	48.	13	$\cos^{-1}$	
24.	41	↑	49.	—00	GTO 00	

Examples:

1. $\cos^{-1} (0.9 + 3i) = 1.29 - 1.86i$

2. $\sec^{-1} (0.9 + 3i) = 1.48 + .30i$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Set to radian mode		f	RAD			
3	For $\cos^{-1} z$, enter z	b	↑				
		a	BST	R/S			x_1
			$x \leftrightarrow y$				y_1
	or						
3	For $\sec^{-1} z$, enter z						
	and calculate 1/z	b	↑				
		a	g	R→P	1/x	$x \leftrightarrow y$	
			CHS	$x \leftrightarrow y$	f	R←P	
			BST	R/S			x_2
			$x \leftrightarrow y$				y_2

COMPLEX INVERSE TRIGONOMETRIC

$\tan^{-1} z$, $\cot^{-1} z$

Let $z = a + ib$ be a complex number. The functions $\tan^{-1}$ (arc tan) and $\cot^{-1} z$ (arc cot) are evaluated by the following formulas:

$$1. \quad \tan^{-1} z = \frac{1}{2} \left[(2k+1)\pi - \tan^{-1} \left(\frac{1+b}{a} \right) - \tan^{-1} \left(\frac{1-b}{a} \right) \right] \\ + \frac{i}{2} \ln \left(\frac{[(1+b)^2 + a^2]^{\frac{1}{2}}}{[(1-b)^2 + a^2]^{\frac{1}{2}}} \right)$$

where $k = 0, \pm 1, \pm 2, \dots$ ($z^2 \neq -1$)

$$2. \quad \cot^{-1} z = \tan^{-1} \left(\frac{1}{z} \right) \quad (z^2 \neq -1 \text{ and } z \neq 0)$$

The rectangular to polar routine is used so the division by zero problem is avoided in the evaluation of $\tan^{-1} \left(\frac{1+b}{a} \right)$ and $\tan^{-1} \left(\frac{1-b}{a} \right)$.

k is assumed to be zero.

Let the solutions be $x_1 + iy_1$ and $x_2 + iy_2$ respectively.

The calculator must be in RADIAN mode and all angles are assumed to be in radians.

DISPLAY		KEY ENTRY	DISPLAY		KEY ENTRY	REGISTERS
LINE	CODE		LINE	CODE		
00.			25.	81	$\div$	R_0
01.	33	STO	26.	31	f	R_1 a
02.	01	1	27.	83	π	R_2 b
03.	22	$x \leftrightarrow y$	28.	23	$R \downarrow$	R_3
04.	33	STO	29.	23	$R \downarrow$	R_4
05.	02	2	30.	61	+	R_5
06.	01	1	31.	51	-	R_6
07.	61	+	32.	02	2	R_7
08.	22	$x \leftrightarrow y$	33.	81	$\div$	R_8
09.	32	g	34.	-00	GTO 00	R_9
10.	00	$R \rightarrow P$	35.	32	g	R_{00}
11.	01	1	36.	00	$R \rightarrow P$	R_{01}
12.	34	RCL	37.	13	$1/x$	R_{02}
13.	02	2	38.	22	$x \leftrightarrow y$	R_{03}
14.	51	-	39.	42	CHS	R_{04}
15.	34	RCL	40.	22	$x \leftrightarrow y$	R_{05}
16.	01	1	41.	31	f	R_{06}
17.	32	g	42.	00	$R \leftarrow P$	R_{07}
18.	00	$R \rightarrow P$	43.	-01	GTO 01	R_{08}
19.	22	$x \leftrightarrow y$	44.			R_{09}
20.	23	$R \downarrow$	45.			
21.	81	$\div$	46.			
22.	31	f	47.			
23.	22	ln	48.			
24.	02	2	49.			

Examples:

- $\tan^{-1} (5 + 8i) = 1.51 + .09i$
- $\cot^{-1} (5 + 8i) = .06 - .09i$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Set to radian mode		f	RAD			
3	For $\tan^{-1} z$, enter z	b	$\uparrow$				
		a	BST	R/S			x_1
			$x \leftrightarrow y$				y_1
	or						
3	For $\cot^{-1} z$, enter z	b	$\uparrow$				
		a	GTO	3	5	R/S	x_2
			$x \leftrightarrow y$				y_2

COMPLEX INVERSE HYPERBOLIC $\sinh^{-1} z$, $\cosh^{-1} z$

Let $z = a + ib$ be a complex number. The functions $\sinh^{-1} z$ (arc hyperbolic sine) and $\cosh^{-1} z$ (arc hyperbolic cosecant) are evaluated by the following formulas:

$$1. \sinh^{-1} z = -i \sin^{-1} iz = (-1)^k \operatorname{sgn}(a) \ln [\alpha + (\alpha^2 - 1)^{1/2}] \\ + [(k\pi + (-1)^k \sin^{-1}(-\beta))] i$$

$$\text{where } \alpha = \frac{1}{2} \sqrt{(1-b)^2 + a^2} + \frac{1}{2} \sqrt{(1+b)^2 + a^2}$$

$$\beta = \frac{1}{2} \sqrt{(1-b)^2 + a^2} - \frac{1}{2} \sqrt{(1+b)^2 + a^2}$$

$$\operatorname{sgn}(a) = a/\sqrt{a^2}$$

$$k = 0, \pm 1, \pm 2, \dots$$

$$2. \cosh^{-1} z = \sinh^{-1} \left(\frac{1}{z} \right) \quad z \neq 0$$

k is assumed to be zero.

Let the solutions be $x_1 + iy_1$ and $x_2 + iy_2$ respectively.

The calculator must be in RADIAN mode and all angles are assumed to be in radians.

Note:

If the calculator flashes zero because $a = 0$ in the division at line 39, the result can still be found by rolling down twice, **R↓**, **R↓**, switching to PRGM, single stepping twice, **SST**, **SST**, switching to RUN, and pressing **R/S**.

DISPLAY		KEY ENTRY	DISPLAY		KEY ENTRY	REGISTERS
LINE	CODE		LINE	CODE		
00.			25.	01	1	R_0
01.	33	STO	26.	51	—	R_1
02.	02	2	27.	31	f	R_2 a
03.	22	$x \leftrightarrow y$	28.	42	$\sqrt{x}$	$R_3 \sqrt{(1-b)^2 + a^2}$
04.	01	1	29.	61	+	R_4
05.	51	—	30.	31	f	R_5
06.	32	g	31.	22	ln	R_6
07.	00	$R \rightarrow P$	32.	34	RCL	R_7
08.	33	STO	33.	02	2	R_8
09.	03	3	34.	41	$\uparrow$	R_9
10.	31	f	35.	32	g	R_{10}
11.	00	$R \leftarrow P$	36.	42	x^2	R_{11}
12.	02	2	37.	31	f	R_{12}
13.	61	+	38.	42	$\sqrt{x}$	R_{13}
14.	32	g	39.	81	$\div$	R_{14}
15.	00	$R \rightarrow P$	40.	71	x	R_{15}
16.	34	RCL	41.	22	$x \leftrightarrow y$	R_{16}
17.	03	3	42.	34	RCL	R_{17}
18.	61	+	43.	03	3	R_{18}
19.	02	2	44.	51	—	R_{19}
20.	81	$\div$	45.	32	g	
21.	41	$\uparrow$	46.	12	$\sin^{-1}$	
22.	41	$\uparrow$	47.	22	$x \leftrightarrow y$	
23.	41	$\uparrow$	48.	-00	GTO 00	
24.	71	x	49.			

Examples:

- $\sinh^{-1} (3.14 + 10.3i) = 3.07 + 1.27i$
- $\cosh^{-1} (3.14 + 10.3i) = .03 - .09i$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Set to radian mode		f	RAD			
3	For $\sinh^{-1} z$, enter z	b	$\uparrow$				
		a	BST	R/S			x_1
			$x \leftrightarrow y$				y_1
	or						
3	For $\cosh^{-1} z$, enter z	b	$\uparrow$				
	and calculate $1/z$	a	g	$R \rightarrow P$	$1/x$	$x \leftrightarrow y$	
			CHS	$x \leftrightarrow y$	f	$R \leftarrow P$	
			BST	R/S			x_2
			$x \leftrightarrow y$				y_2

COMPLEX INVERSE HYPERBOLIC $\cosh^{-1} z$, $\operatorname{sech}^{-1} z$

Let $z = a + ib$ be a complex number. The functions $\cosh^{-1} z$ (arc hyperbolic cosine) and $\operatorname{sech}^{-1} z$ (arc hyperbolic secant) are evaluated by the following formulas:

$$1. \quad \cosh^{-1} z = 2k\pi i \pm \left\{ \operatorname{sgn}(b) \ln [\alpha + (\alpha^2 - 1)^{1/2}] + i \cos^{-1} \beta \right\}$$

$$\text{where } \alpha = \frac{1}{2} \sqrt{(a+1)^2 + b^2} + \frac{1}{2} \sqrt{(a-1)^2 + b^2}$$

$$\beta = \frac{1}{2} \sqrt{(a+1)^2 + b^2} - \frac{1}{2} \sqrt{(a-1)^2 + b^2}$$

$$\operatorname{sgn}(b) = b / \sqrt{b^2}$$

$$k = 0, \pm 1, \pm 2, \dots$$

$$2. \quad \operatorname{sech}^{-1} z = \cosh^{-1} \left(\frac{1}{z} \right) \quad z \neq 0$$

Let the solutions be $x_1 + iy_1$ and $x_2 + iy_2$ respectively.

The calculator must be in RADIAN mode.

If the calculator flashes zero because $b = 0$ in the division at line 40, the result can still be found by rolling down twice, **R↓**, **R↓**, switching to PRGM, single stepping twice, **SST**, **SST**, switching to RUN, and pressing **R/S**.

DISPLAY		KEY ENTRY	DISPLAY		KEY ENTRY	REGISTERS
LINE	CODE		LINE	CODE		
00.			25.	71	x	R_0
01.	22	$x \leftrightarrow y$	26.	01	1	R_1
02.	33	STO	27.	51	—	R_2 b
03.	02	2	28.	31	f	$R_3 \sqrt{(a-1)^2 + b^2}$
04.	22	$x \leftrightarrow y$	29.	42	$\sqrt{x}$	R_4
05.	01	1	30.	61	+	R_5
06.	51	—	31.	31	f	R_6
07.	32	g	32.	22	ln	R_7
08.	00	R→P	33.	34	RCL	R_8
09.	33	STO	34.	02	2	R_9
10.	03	3	35.	41	↑	R_{10}
11.	31	f	36.	32	g	R_{11}
12.	00	R←P	37.	42	x^2	R_{12}
13.	02	2	38.	31	f	R_{13}
14.	61	+	39.	42	$\sqrt{x}$	R_{14}
15.	32	g	40.	81	÷	R_{15}
16.	00	R→P	41.	71	x	R_{16}
17.	34	RCL	42.	22	$x \leftrightarrow y$	R_{17}
18.	03	3	43.	34	RCL	R_{18}
19.	61	+	44.	03	3	R_{19}
20.	02	2	45.	51	—	
21.	81	÷	46.	32	g	
22.	41	↑	47.	13	$\cos^{-1}$	
23.	41	↑	48.	22	$x \leftrightarrow y$	
24.	41	↑	49.	-00	GTO 00	

Examples:

1. $\cosh^{-1} (5 + 8i) = 2.94 + 1.01i$

2. $\operatorname{sech}^{-1} (5 + 8i) = -.09 + 1.51i$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Set to radian mode		f	RAD			
3	For $\cosh^{-1} z$, enter z	b	↑				
		a	BST	R/S			x_1
			$x \leftrightarrow y$				y_1
	or						
3	For $\operatorname{sech}^{-1} z$, enter z	b	↑				
	and calculate $1/z$	a	g	R→P	1/x	$x \leftrightarrow y$	
			CHS	$x \leftrightarrow y$	f	R←P	
			BST	R/S			x_2
			$x \leftrightarrow y$				y_2

COMPLEX INVERSE HYPERBOLIC $\tanh^{-1} z, \coth^{-1} z$

Let $z = a + ib$ be a complex number. The functions $\tanh^{-1} z$ (arc hyperbolic tangent) and $\coth^{-1} z$ (arc hyperbolic cotangent) are evaluated by the following formulas:

1. $\tanh^{-1} z = -i \tan^{-1} i z = \frac{1}{2} \ln \left[\frac{[(1+a)^2 + b^2]^{\frac{1}{2}}}{[(1-a)^2 + b^2]^{\frac{1}{2}}} \right] + \left(\frac{i}{2} \right) \left[-(2k+1)\pi + \tan^{-1} \left(\frac{1+a}{-b} \right) + \tan^{-1} \left(\frac{1-a}{-b} \right) \right] \quad (z^2 \neq 1)$
2. $\coth^{-1} z = \tanh^{-1} \left(\frac{1}{z} \right) \quad (z^2 \neq 1, z \neq 0)$

The rectangular to polar routine is used so the division by zero problem is avoided in the evaluation of $\tan^{-1} \left(\frac{1+a}{-b} \right)$ and $\tan^{-1} \left(\frac{1-a}{-b} \right)$.

k is assumed to be zero.

Let the solutions be $x_1 + iy_1$ and $x_2 + iy_2$ respectively.

The calculator must be in RADIAN mode.

DISPLAY		KEY ENTRY	DISPLAY		KEY ENTRY	REGISTERS
LINE	CODE		LINE	CODE		
00.			25.	81	$\div$	R_0
01.	33	STO	26.	31	f	$R_1 -b$
02.	02	2	27.	83	π	$R_2 a$
03.	01	1	28.	23	$R\downarrow$	R_3
04.	61	+	29.	23	$R\downarrow$	R_4
05.	22	$x\dot{z}y$	30.	61	+	R_5
06.	42	CHS	31.	51	-	R_6
07.	33	STO	32.	02	2	R_7
08.	01	1	33.	81	$\div$	R_8
09.	32	g	34.	42	CHS	R_9
10.	00	$R\rightarrow P$	35.	22	$x\dot{z}y$	R_{00}
11.	01	1	36.	-00	GTO 00	R_{01}
12.	34	RCL	37.	32	g	R_{02}
13.	02	2	38.	00	$R\rightarrow P$	R_{03}
14.	51	-	39.	13	$1/x$	R_{04}
15.	34	RCL	40.	22	$x\dot{z}y$	R_{05}
16.	01	1	41.	42	CHS	R_{06}
17.	32	g	42.	22	$x\dot{z}y$	R_{07}
18.	00	$R\rightarrow P$	43.	31	f	R_{08}
19.	22	$x\dot{z}y$	44.	00	$R\leftarrow P$	R_{09}
20.	23	$R\downarrow$	45.	-01	GTO 01	
21.	81	$\div$	46.			
22.	31	f	47.			
23.	22	ln	48.			
24.	02	2	49.			

Examples:

- $\tanh^{-1} (8 - 5i) = .09 - 1.51i$
- $\coth^{-1} (-7i) = .14i$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Set to radian mode		f	RAD			
3	For $\tanh^{-1} z$, enter z	b	$\uparrow$				
		a	BST	R/S			x_1
			$x\dot{z}y$				y_1
	or						
3	For $\coth^{-1} z$, enter z	b	$\uparrow$				
		a	GTO	3	7	R/S	x_2
			$x\dot{z}y$				y_2

COMPLEX POLYNOMIAL EVALUATION

Given a polynomial (with complex coefficients) of the form

$$f(z) = (a_1 + ib_1) z^n + (a_2 + ib_2) z^{n-1} + \dots + (a_n + ib_n) z + (a_{n+1} + ib_{n+1})$$

This program evaluates $f(z)$ for a complex number $z_0 = c + di$. Let the solution be $x + iy$.

If a coefficient is zero it still must be entered.

The polynomial is evaluated in the form

$$(\dots ((a_1 + ib_1) z + (a_2 + ib_2)) z + \dots) + (a_{n+1} + ib_{n+1})$$

DISPLAY		KEY ENTRY	DISPLAY		KEY ENTRY	REGISTERS
LINE	CODE		LINE	CODE		
00.			25.	34	RCL	R ₀
01.	33	STO	26.	07	7	R ₁ c
02.	05	5	27.	31	f	R ₂ d
03.	23	R↓	28.	00	R←P	R ₃ a ₁
04.	33	STO	29.	34	RCL	R ₄ b ₁
05.	06	6	30.	05	5	R ₅ Used
06.	34	RCL	31.	61	+	R ₆ Used
07.	04	4	32.	33	STO	R ₇ Used
08.	34	RCL	33.	03	3	R ₈
09.	03	3	34.	22	x↔y	R ₉
10.	32	g	35.	34	RCL	R ₀₀
11.	00	R→P	36.	06	6	R ₀₁
12.	34	RCL	37.	61	+	R ₀₂
13.	02	2	38.	33	STO	R ₀₃
14.	34	RCL	39.	04	4	R ₀₄
15.	01	1	40.	22	x↔y	R ₀₅
16.	32	g	41.	-00	GTO 00	R ₀₆
17.	00	R→P	42.			R ₀₇
18.	22	x↔y	43.			R ₀₈
19.	23	R↓	44.			R ₀₉
20.	71	x	45.			
21.	33	STO	46.			
22.	07	7	47.			
23.	23	R↓	48.			
24.	61	+	49.			

Example:

Evaluate the complex polynomial $f(z) = (3 + 4i) z^4 + 18 z^3 + (-2 + i) z^2 - 10 z + (5 - 7i)$ at the complex number $z_0 = 2 + i$.

Solution:

$$f(z_0) = -106.00 + 220.00i$$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Store z_0	d	STO	2			
		c	STO	1			
3	Store $a_1 + b_1 i$	b_1	STO	4			
		a_1	STO	3	BST		
4	Repeat this step for $k = 2, 3, \dots, n$						
	Enter $a_k + b_k i$	b_k	↑				
		a_k	R/S				Temp
5	Enter $a_{n+1} + ib_{n+1}$	b_{n+1}	↑				
		a_{n+1}	R/S				x
			$x \leftrightarrow y$				y

COMPOUNDED AMOUNT

Let n = number of time periods

i = periodic interest rate expressed as a decimal, e.g., 6% is represented as .06

PV = present value or principal

FV = future value or amount

I = interest amount

Each value can be calculated from the others by the following formulas:

$$1. \quad FV = PV (1 + i)^n$$

$$2. \quad PV = FV (1 + i)^{-n}$$

$$3. \quad n = \frac{\ln (FV/PV)}{\ln (1 + i)}$$

$$4. \quad i = \left(\frac{FV}{PV} \right)^{1/n} - 1$$

$$5. \quad I = PV [(1 + i)^n - 1]$$

DISPLAY		KEY ENTRY	DISPLAY		KEY ENTRY	REGISTERS
LINE	CODE		LINE	CODE		
00.			25.	04	4	R ₀
01.	34	RCL	26.	34	RCL	R ₁ n, -n
02.	02	2	27.	01	1	R ₂ i
03.	01	1	28.	13	$1/x$	R ₃ PV, FV
04.	61	+	29.	12	y^x	R ₄ FV/PV
05.	34	RCL	30.	01	1	R ₅
06.	01	1	31.	51	-	R ₆
07.	12	y^x	32.	-00	GTO 00	R ₇
08.	34	RCL	33.	34	RCL	R ₈
09.	03	3	34.	02	2	R ₉
10.	71	x	35.	01	1	R ₀₀
11.	-00	GTO 00	36.	61	+	R ₀₁
12.	34	RCL	37.	34	RCL	R ₀₂
13.	04	4	38.	01	1	R ₀₃
14.	31	f	39.	12	y^x	R ₀₄
15.	22	ln	40.	01	1	R ₀₅
16.	34	RCL	41.	51	-	R ₀₆
17.	02	2	42.	34	RCL	R ₀₇
18.	01	1	43.	03	3	R ₀₈
19.	61	+	44.	71	x	R ₀₉
20.	31	f	45.	-00	GTO 00	
21.	22	ln	46.			
22.	81	÷	47.			
23.	-00	GTO 00	48.			
24.	34	RCL	49.			

46 Compounded Amount

Examples:

1. Find the future amount of \$1500 invested at 6% (.06) compounded annually for 5 years.
2. What sum invested today at 6% compounded annually will amount to \$2007.34 in 5 years?
3. How long does it take \$100 to double if it is invested at 6% annual interest compounded quarterly?
4. Find the rate of return on \$2000 compounded monthly if it amounts to \$3000 at the end of 5 years?
5. Find the compound interest on \$1500 for 5 years if interest at 6% is compounded annually.

Solutions:

1. \$2007.34
2. \$1500.00
3. 46.56 quarters = 11.64 years ($i = .06/4$)
4. .0068 monthly = 8.14% annually ($n = 60$)
5. \$507.34

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	For FV	n	STO	1			FV
		i	STO	2			
		PV	STO	3			
			BST	R/S			
2	For PV	n	CHS	STO	1		PV
		i	STO	2			
		FV	STO	3			
			BST	R/S			
2	For n	FV	↑				n
		PV	÷	STO	4		
		i	STO	2			
			GTO	1	2	R/S	
2	For i	FV	↑				i
		PV	÷	STO	4		
		n	STO	1			
			GTO	2	4	R/S	
2	For I	n	STO	1			I
		i	STO	2			
		PV	STO	3			
			GTO	3	3	R/S	

DIRECT REDUCTION LOAN INTEREST RATE

This program calculates the interest rate on a mortgage where payments are made at the end of period.

Let n = number of payments

i = periodic interest rate expressed as a decimal, e.g., 6% is represented as .06

PMT = payment

PV = present value or principal

This routine solves the equation $f(i)$ by an iteration for i using Newton's method:

$$i_{k+1} = i_k - \frac{f(i)}{f'(i)}$$

$$\text{where } f(i) = \frac{1 - (1 + i)^{-n}}{i} - \frac{PV}{PMT}$$

and $f'(i)$ is the first derivative of $f(i)$

First an initial guess i_0 is stored in R_2 . For i_0 either the suggested routine or a rough guess can be used. However, i_0 cannot be zero. If the suggested routine produces zero, then zero is the solution.

DISPLAY		KEY ENTRY	DISPLAY		KEY ENTRY	REGISTERS
LINE	CODE		LINE	CODE		
00.			25.	61	+	R ₀
01.	34	RCL	26.	81	÷	R ₁ n
02.	03	3	27.	01	1	R ₂ i
03.	34	RCL	28.	61	+	R ₃ PV/PMT
04.	02	2	29.	34	RCL	R ₄
05.	71	x	30.	05	5	R ₅ (1 + i) ⁻ⁿ
06.	01	1	31.	71	x	R ₆
07.	34	RCL	32.	01	1	R ₇
08.	02	2	33.	51	—	R ₈
09.	01	1	34.	34	RCL	R ₉
10.	61	+	35.	02	2	R ₀₀
11.	34	RCL	36.	81	÷	R ₀₁
12.	01	1	37.	81	÷	R ₀₂
13.	42	CHS	38.	33	STO	R ₀₃
14.	12	y ^x	39.	61	+	R ₀₄
15.	33	STO	40.	02	2	R ₀₅
16.	05	5	41.	41	↑	R ₀₆
17.	51	—	42.	71	x	R ₀₇
18.	51	—	43.	43	EEX	R ₀₈
19.	34	RCL	44.	01	1	R ₀₉
20.	01	1	45.	02	2	
21.	34	RCL	46.	42	CHS	
22.	02	2	47.	31	f	
23.	13	1/x	48.	-01	x ≤ y 01	
24.	01	1	49.	-00	GTO 00	

50 Direct Reduction Loan—Interest Rate

Example:

Find the monthly interest rate on a mortgage of \$30,000 if the loan requires 360 monthly payments of \$179.86 to be paid off.

Answer:

.0050 (0.5%) **FIX** **4** (15 seconds iteration)

The suggested routine supplies an initial guess of .0047 (.47%). The user can notice the change in iteration time if he tries different i_0 's such as .001 or .01 (25 seconds iteration).

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Store PV and PMT	PV	↑				
		PMT	÷	STO	3		
3	Store n	n	STO	1			
4	Store a rough guess	i_0	STO	2			
	or						
4	Calculate an initial guess		RCL	3	1/x	RCL	
			1	↑	x	1/x	
			RCL	3	x	-	
			STO	2			i_0
5	Iterate for i		BST	R/S			1.0000000-12
6	When iteration stops		RCL	2			i

DIRECT REDUCTION LOAN PAYMENT, PRESENT VALUE, NUMBER OF TIME PERIODS

Calculates payment, present value, and number of time periods of a mortgage given two of the three and the interest rate.

Let n = number of payment periods

PV = present value or principal

PMT = payment

i = periodic interest rate expressed as a decimal, e.g., 6% is represented as .06.

Then, PMT, PV, and n can be calculated from the other three by the following formulas:

$$1. \text{ PMT} = \text{PV} \left[\frac{i}{1 - (1 + i)^{-n}} \right]$$

$$2. \text{ PV} = \text{PMT} \left[\frac{1 - (1 + i)^{-n}}{i} \right]$$

$$3. \text{ } n = - \frac{\ln (1 - i\text{PV}/\text{PMT})}{\ln (1 + i)}$$

52 Direct Reduction Loan—Pmt, Pv, n

DISPLAY			DISPLAY			REGISTERS
LINE	CODE	KEY ENTRY	LINE	CODE	KEY ENTRY	
00.			25.	34	RCL	R ₀
01.	34	RCL	26.	02	2	R ₁ n
02.	02	2	27.	81	÷	R ₂ i
03.	01	1	28.	34	RCL	R ₃ PMT, PV, PV/PMT
04.	34	RCL	29.	03	3	R ₄
05.	02	2	30.	71	x	R ₅
06.	01	1	31.	-00	GTO 00	R ₆
07.	61	+	32.	01	1	R ₇
08.	34	RCL	33.	34	RCL	R ₈
09.	01	1	34.	03	3	R ₉
10.	42	CHS	35.	34	RCL	R ₁₀
11.	12	y ^x	36.	02	2	R ₁₁
12.	51	—	37.	71	x	R ₁₂
13.	81	÷	38.	51	—	R ₁₃
14.	-28	GTO 28	39.	31	f	R ₁₄
15.	01	1	40.	22	ln	R ₁₅
16.	34	RCL	41.	34	RCL	R ₁₆
17.	02	2	42.	02	2	R ₁₇
18.	01	1	43.	01	1	R ₁₈
19.	61	+	44.	61	+	R ₁₉
20.	34	RCL	45.	31	f	
21.	01	1	46.	22	ln	
22.	42	CHS	47.	81	÷	
23.	12	y ^x	48.	42	CHS	
24.	51	—	49.	-00	GTO 00	

Examples:

1. To pay off a loan of \$4000 at 9.5% (.095) in 30 months, what monthly payment is required?
2. A person is willing to pay \$150 per month for 30 months for a loan at 9.5%. How much can be borrowed?
3. How many monthly payments of \$100 must be made to pay off a loan of \$4000 at 9.5% annual interest?

Note:

Divide 0.095 by 12 to find the monthly interest rate expressed as a decimal.

Answers:

1. \$150.32
2. \$3991.55
3. 48.29 (4.02 years)

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	For payment	PV	STO	3			PMT
		i	STO	2			
		n	STO	1			
			BST	R/S			
	or						
2	For present value	PMT	STO	3			PV
		i	STO	2			
		n	STO	1			
			GTO	1	5	R/S	
	or						
3	For number of payments	PV	↑				n
		PMT	÷	STO	3		
		i	STO	2			
			GTO	3	2	R/S	

DIRECT REDUCTION LOAN ACCUMULATED INTEREST, REMAINING BALANCE

This program finds the accumulated interest and remaining balance of a mortgage.

Let I_{c-k} = the accumulated interest paid by payments c through k

PV_k = the remaining balance after payment k .

n = number of payments

i = periodic interest rate expressed as a decimal, e.g., 6% is expressed as .06

$j = c - 1$

Then, I_{c-k} and PV_k can be calculated by the following formulas:

$$1. \quad I_{c-k} = PMT \left[k - j - \frac{(1+i)^{k-n}}{i} (1 - (1+i)^{j-k}) \right]$$

$$2. \quad PV_k = \frac{PMT}{i} [1 - (1+i)^{k-n}]$$

DISPLAY			DISPLAY			KEY ENTRY	REGISTERS
LINE	CODE	KEY ENTRY	LINE	CODE	KEY ENTRY		
00.			25.	34	RCL		R_0
01.	34	RCL	26.	02	2		R_1 n
02.	02	2	27.	81	$\div$		R_2 i
03.	01	1	28.	34	RCL		R_3 PMT
04.	61	+	29.	06	6		R_4 k
05.	34	RCL	30.	51	—		R_5 j
06.	04	4	31.	-47	GTO 47		R_6 j-k
07.	34	RCL	32.	01	1		R_7
08.	01	1	33.	34	RCL		R_8
09.	00	0	34.	02	2		R_9
10.	61	+	35.	01	1		R_{00}
11.	51	—	36.	61	+		R_{01}
12.	12	y^x	37.	34	RCL		R_{02}
13.	22	$x \nabla y$	38.	04	4		R_{03}
14.	34	RCL	39.	34	RCL		R_{04}
15.	05	5	40.	01	1		R_{05}
16.	34	RCL	41.	51	—		R_{06}
17.	04	4	42.	12	y^x		R_{07}
18.	51	—	43.	51	—		R_{08}
19.	33	STO	44.	34	RCL		R_{09}
20.	06	6	45.	02	2		
21.	12	y^x	46.	81	$\div$		
22.	01	1	47.	34	RCL		
23.	51	—	48.	03	3		
24.	71	x	49.	71	x		

56 Direct Reduction Loan—Accumulated Interest, Remaining Balance

Examples:

- 1. A house costs \$30,000 with a mortgage life of 30 years at 8% (.08) yearly interest. Find the interest paid on the first 36 monthly payments. The payment can be calculated from the program, Direct Reduction Loan Payment but is \$220.13.
- 2. Using the above example calculate the remaining balance after the 36th payment.

Solutions:

- 1. \$7108.72
- 2. \$29,184.13

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	For accumulated interest of c th through k th payment	n	STO	1			Ic-k
		i	STO	2			
		PMT	STO	3			
		k	STO	4			
		c-i	STO	5			
			BST	R/S			
	or						
2	For remaining balance after k th payment	n	STO	1			PV _k
		i	STO	2			
		PMT	STO	3			
		k	STO	4			
			GTO	3	2	R/S	

DIRECT REDUCTION LOAN AMORTIZATION SCHEDULES

This program calculates a table of interest paid, payment to principal, and present value of a mortgage. It also can be used to find accumulated interest.

Let I_k = interest paid in k^{th} payment

PMT = payment

PP_k = payment to principal of k^{th} payment

PV_k = remaining balance after k^{th} payment

PV_0 = amount of loan

i = periodic interest rate expressed as a decimal, e.g., 6% is expressed as .06

An amortization schedule consists of the interest paid, the payment to principal, and the remaining balance for each payment $k = 1, 2, \dots$

These quantities are calculated by the following formulas:

1. $I_k = i PV_{k-1}$
2. $PP_k = PMT - I_k$
3. $PV_k = PV_{k-1} - PP_k$

58 Direct Reduction Loan—Amortization Schedules

DISPLAY			DISPLAY			REGISTERS
LINE	CODE	KEY ENTRY	LINE	CODE	KEY ENTRY	
00.			25.	34	RCL	R ₀
01.	00	0	26.	06	6	R ₁ ΣI_K
02.	33	STO	27.	51	—	R ₂ i
03.	01	1	28.	33	STO	R ₃ PMT
04.	34	RCL	29.	04	4	R ₄ PV _K
05.	04	4	30.	84	R/S	R ₅ I _K
06.	34	RCL	31.	-04	GTO 04	R ₆ PP _K
07.	02	2	32.			R ₇
08.	71	x	33.			R ₈
09.	33	STO	34.			R ₉
10.	05	5	35.			R ₀₀
11.	33	STO	36.			R ₀₁
12.	61	+	37.			R ₀₂
13.	01	1	38.			R ₀₃
14.	84	R/S	39.			R ₀₄
15.	34	RCL	40.			R ₀₅
16.	03	3	41.			R ₀₆
17.	34	RCL	42.			R ₀₇
18.	05	5	43.			R ₀₈
19.	51	—	44.			R ₀₉
20.	33	STO	45.			
21.	06	6	46.			
22.	84	R/S	47.			
23.	34	RCL	48.			
24.	04	4	49.			

Example:

Find the amortization schedule for a loan of \$30,000 at 7% (.07) annual interest with payments of \$200 monthly.

Note:

Be sure to enter monthly interest rate by dividing 0.07 by 12.

Solution:

Payment No.	I_k	PP_k	PV_k	ΣI
1	175.00	25.00	29,975.00	175.00
2	174.85	25.15	29,949.85	349.85
3	174.71	25.29	29,924.56	524.56

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Store periodic interest rate	i	STO	2			
3	Store payment	PMT	STO	3			
4	Store initial loan amount	PV_0	STO	4	BST		
5	Calculate interest		R/S				I_k
6	Calculate payment to principal		R/S				PP_k
7	Calculate remaining balance		R/S				PV_k
8	Repeat steps 5, 6, and 7						
	For $k = 2, 3, 4, \dots$						
	Note: At anytime after steps						
	5, 6, or 7 have been executed the						
	accumulated interest can be						
	found.		RCL	1			ΣI

SINKING FUND INTEREST RATE

This program calculates the interest rate on a savings program where payments are assumed to be made at the end of compounding period.

Let n = number of payments

i = periodic interest rate expressed as a decimal, e.g., 6% is expressed as .06

PMT = payment

FV = future value or amount

This routine solves the equation $f(i)$ by an iteration for i using Newton's method:

$$i_{k+1} = i_k - \frac{f(i)}{f'(i)}$$

$$\text{where } f(i) = \frac{(1+i)^n - 1}{i} - \frac{FV}{PMT}$$

and $f'(i)$ is the first derivative of $f(i)$. First an initial guess i_0 is stored in R_2 . Either the suggested routine or a rough guess can be used. However, i_0 cannot be zero. If the suggested routine produces zero, then zero is the solution.

DISPLAY		KEY ENTRY	DISPLAY		KEY ENTRY	REGISTERS
LINE	CODE		LINE	CODE		
00.			25.	81	÷	R ₀
01.	34	RCL	26.	01	1	R ₁ n
02.	03	3	27.	51	—	R ₂ i
03.	34	RCL	28.	34	RCL	R ₃ FV/PMT
04.	02	2	29.	04	4	R ₄ (1 + i) ⁿ
05.	71	x	30.	71	x	R ₅
06.	01	1	31.	01	1	R ₆
07.	61	+	32.	61	+	R ₇
08.	34	RCL	33.	34	RCL	R ₈
09.	02	2	34.	02	2	R ₉
10.	01	1	35.	81	÷	R ₀₀
11.	61	+	36.	81	÷	R ₀₁
12.	34	RCL	37.	33	STO	R ₀₂
13.	01	1	38.	61	+	R ₀₃
14.	12	y ^x	39.	02	2	R ₀₄
15.	33	STO	40.	41	↑	R ₀₅
16.	04	4	41.	71	x	R ₀₆
17.	51	—	42.	43	EEX	R ₀₇
18.	34	RCL	43.	42	CHS	R ₀₈
19.	01	1	44.	01	1	R ₀₉
20.	01	1	45.	02	2	
21.	34	RCL	46.	31	f	
22.	02	2	47.	-01	x≤y 01	
23.	13	1/x	48.	34	RCL	
24.	61	+	49.	02	2	

62 Sinking Fund—Interest Rate

Example:

What annual rate of interest must be obtained to amass a total of \$10,000 in 10 years on an annual investment of \$600.

Solution:

.1093 (10.93%) **FIX** **4** (20 seconds iteration)

The suggested routine supplies an initial guess of .1365 (13.65%). The user can try other initial guesses such as .10 or .20 and notice the change in iteration time. (15 and 25 seconds iteration time respectively.)

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Store FV and PMT	FV	↑				
		PMT	÷	STO	3		
3	Store n	n	STO	1			
4	Enter a rough guess for i	i_0	STO	2			
	or						
4	Calculate an initial guess		RCL	3	RCL	1	
			−	2	x	RCL	
			1	1	−	↑	
			x	RCL	3	+	
			÷	STO	2		i_0
5	Calculate i		BST	R/S			i

SINKING FUND PAYMENT, FUTURE VALUE, NUMBER OF TIME PERIODS

This program calculates payment, future value, or number of time periods given two of the three and the interest rate.

Let n = number of payments

i = periodic interest rate expressed as a decimal, e.g., 6% is expressed as .06

PMT = payment

FV = future value

Then, FV, PMT, or n can be calculated from the other three by the following formulas:

$$1. \quad FV = PMT \left[\frac{(1+i)^n - 1}{i} \right]$$

$$2. \quad PMT = FV \left[\frac{i}{(1+i)^n - 1} \right]$$

$$3. \quad n = \frac{\ln \left(i \frac{FV}{PMT} + 1 \right)}{\ln (1+i)}$$

64 Sinking Fund—Pmt, Fv, n

DISPLAY		KEY ENTRY	DISPLAY		KEY ENTRY	REGISTERS
LINE	CODE		LINE	CODE		
00.			25.	61	+	R ₀
01.	34	RCL	26.	34	RCL	R ₁ n
02.	02	2	27.	01	1	R ₂ i
03.	01	1	28.	12	y ^x	R ₃ PMT, FV, FV/PMT
04.	61	+	29.	01	1	R ₄
05.	34	RCL	30.	51	—	R ₅
06.	01	1	31.	81	÷	R ₆
07.	12	y ^x	32.	-00	GTO 00	R ₇
08.	01	1	33.	34	RCL	R ₈
09.	51	—	34.	03	3	R ₉
10.	34	RCL	35.	34	RCL	R ₀₀
11.	02	2	36.	02	2	R ₀₁
12.	81	÷	37.	71	x	R ₀₂
13.	34	RCL	38.	01	1	R ₀₃
14.	03	3	39.	61	+	R ₀₄
15.	71	x	40.	31	f	R ₀₅
16.	-00	GTO 00	41.	22	ln	R ₀₆
17.	34	RCL	42.	34	RCL	R ₀₇
18.	02	2	43.	02	2	R ₀₈
19.	34	RCL	44.	01	1	R ₀₉
20.	03	3	45.	61	+	
21.	71	x	46.	31	f	
22.	34	RCL	47.	22	ln	
23.	02	2	48.	81	÷	
24.	01	1	49.	-00	GTO 00	

Examples:

1. How much money will a person have if he saves \$100 a month for 5 years at 7% (.07) annual interest?
2. How big a monthly payment does a person have to make to save \$10,000 at the end of 5 years? Assume an annual interest rate of 7%.
3. How long will it take to save \$10,000 making monthly payments of \$100 at 7% annual interest?

Note:

Remember to enter interest rate as 0.07 divided by 12.

Solutions:

1. \$7,159.29
2. \$139.68
3. 79.01 months or 6.58 years

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	For FV	n	STO	1			FV
		i	STO	2			
		PMT	STO	3			
			BST	R/S			
	or						
2	For PMT	n	STO	1			PMT
		i	STO	2			
		FV	STO	3			
			GTO	1	7	R/S	
	or						
2	For n	FV	↑				n
		PMT	÷	STO	3		
		i	STO	2			
			GTO	3	3	R/S	

DISCOUNTED CASH FLOW ANALYSIS

Let PV_0 = original investment

PV_k = cash flow of k^{th} period

i = discount rate per period as a decimal, e.g., 6% is expressed as .06

C_k = net present value at period k

Then

$$C_k = -PV_0 + \sum_{k=1}^n \frac{PV_k}{(1+i)^k}$$

DISPLAY		KEY ENTRY	DISPLAY		KEY ENTRY	REGISTERS
LINE	CODE		LINE	CODE		
00.			25.	33	STO	R_0
01.	34	RCL	26.	03	3	R_1 PV_0
02.	02	2	27.	12	y^x	R_2 i
03.	01	1	28.	81	$\div$	R_3 k
04.	61	+	29.	34	RCL	R_4 C_k
05.	81	$\div$	30.	04	4	R_5
06.	01	1	31.	61	+	R_6
07.	33	STO	32.	-13	GTO 13	R_7
08.	03	3	33.			R_8
09.	23	$R\downarrow$	34.			R_9
10.	34	RCL	35.			R_{00}
11.	01	1	36.			R_{01}
12.	51	-	37.			R_{02}
13.	33	STO	38.			R_{03}
14.	04	4	39.			R_{04}
15.	84	R/S	40.			R_{05}
16.	41	$\uparrow$	41.			R_{06}
17.	01	1	42.			R_{07}
18.	34	RCL	43.			R_{08}
19.	02	2	44.			R_{09}
20.	61	+	45.			
21.	34	RCL	46.			
22.	03	3	47.			
23.	01	1	48.			
24.	61	+	49.			

Example:

You are offered an investment opportunity for \$100,000 at a capital cost of 10% after taxes. Will this investment be profitable based on the following cash flows?

Year	Cash Flow
1	\$34,000
2	\$27,500
3	\$59,700
4	\$ 7,800

Solution:

$$C_1 = \$-69,090.91$$

$$C_2 = \$-46,363.64$$

$$C_3 = \$-1,510.14$$

$$C_4 = \$3817.36$$

Since C_4 is positive the cash flow is profitable to the extent that the cost of capital is 10%.

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Store PV_0 and i	PV_0	STO	1			
		i	STO	2			
3	Enter PV_1	PV_1	BST	R/S			C_1
4	Perform for $k = 2, \dots, n$	PV_k	R/S				C_k

DEPRECIATION SCHEDULES

STRAIGHT LINE

Let PV = original value of asset (less salvage value)

n = lifetime number of periods of asset

B_k = book value at time period K

D = each year's depreciation

k = number of time period, i.e., 1, 2, 3, ..., or n

Then, B_k and D can be calculated by the following formulas:

1. $D = PV/n$
2. $B_k = PV - kD$

DISPLAY		KEY ENTRY	DISPLAY		KEY ENTRY	REGISTERS
LINE	CODE		LINE	CODE		
00.			25.			$R_0 D$
01.	34	RCL	26.			$R_1 n$
02.	03	3	27.			R_2
03.	34	RCL	28.			$R_3 PV$
04.	01	1	29.			$R_4 k$
05.	81	$\div$	30.			R_5
06.	33	STO	31.			R_6
07.	00	0	32.			R_7
08.	84	R/S	33.			R_8
09.	34	RCL	34.			R_9
10.	03	3	35.			R_{00}
11.	34	RCL	36.			R_{01}
12.	04	4	37.			R_{02}
13.	34	RCL	38.			R_{03}
14.	00	0	39.			R_{04}
15.	71	$\times$	40.			R_{05}
16.	51	—	41.			R_{06}
17.	84	R/S	42.			R_{07}
18.	01	1	43.			R_{08}
19.	33	STO	44.			R_{09}
20.	61	+	45.			
21.	04	4	46.			
22.	-01	GTO 01	47.			
23.			48.			
24.			49.			

Example:

A fleet car has a value (not including salvage value) of \$2100 and a life expectancy of six years. Using the straight line method, what is the amount of depreciation and what is the book value after two years?

Solutions:

$$D = \$350.00$$

$$B_2 = \$1400.00$$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Store n, PV, and k	n	STO	1			
		PV	STO	3			
		k	STO	4	BST		
3	To calculate D and B _k		R/S				D
			R/S				B _k
4	For the next year go to step 3						

DEPRECIATION SCHEDULES SUM-OF-THE-YEAR'S DIGITS

Let n = life time number of periods of asset

S = salvage value

D_k = depreciaton over time period k

B_k = book value at time period k

PV = original value of asset (less salvage value)

k = number of time period, i.e., 1, 2, 3, ..., or n

Then, D_k and B_k can be calculated by the following formulas:

$$1. D_k = \frac{2(n - k + 1)}{n(n + 1)} PV$$

$$2. B_k = S + \frac{(n - k) D_k}{2}$$

DISPLAY			KEY ENTRY	DISPLAY			KEY ENTRY	REGISTERS
LINE	CODE			LINE	CODE			
00.				25.	84	R/S		$R_0 D_k$
01.	34	RCL		26.	34	RCL		$R_1 n$
02.	01	1		27.	02	2		$R_2 n - k$
03.	34	RCL		28.	02	2		$R_3 PV$
04.	04	4		29.	81	÷		$R_4 k$
05.	51	—		30.	34	RCL		$R_5 S$
06.	33	STO		31.	00	0		R_6
07.	02	2		32.	71	x		R_7
08.	01	1		33.	34	RCL		R_8
09.	61	+		34.	05	5		R_9
10.	02	2		35.	61	+		R_{10}
11.	71	x		36.	84	R/S		R_{11}
12.	34	RCL		37.	01	1		R_{12}
13.	03	3		38.	33	STO		R_{13}
14.	71	x		39.	61	+		R_{14}
15.	34	RCL		40.	04	4		R_{15}
16.	01	1		41.	-01	GTO 01		R_{16}
17.	01	1		42.				R_{17}
18.	61	+		43.				R_{18}
19.	34	RCL		44.				R_{19}
20.	01	1		45.				
21.	71	x		46.				
22.	81	÷		47.				
23.	33	STO		48.				
24.	00	0		49.				

Example:

A car has a value (not including the salvage value of \$800) of \$2100 and a life expectancy of 6 years. Using the Sum-of-Year's Digits method what is the amount of depreciation and what is the book value after 2 years?

Solutions:

$$D_2 = \$500.00$$

$$B_2 = \$1800.00$$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Store n, PV, k, and S	n	STO	1			
		PV	STO	3			
		k	STO	4			
		S	STO	5	BST		
3	To calculate B_k and D_k		R/S				D_k
			R/S				B_k
4	For the next year go to step 3						

DEPRECIATION SCHEDULES

VARIABLE RATE DECLINING BALANCE

Let PV = original value of asset (less salvage value)

n = lifetime periods of asset

R = depreciation rate (given by user)

D_k = depreciation at time period k

B_k = book value at time period k

k = number of time period, i.e., 1, 2, 3, ..., or n

Then, D_k and B_k can be calculated by the following formulas:

$$1. \quad D_k = PV \frac{R}{n} \left(1 - \frac{R}{n} \right)^{k-1}$$

$$2. \quad B_k = PV \left(1 - \frac{R}{n} \right)^k$$

If $R = 2$ the program gives the double declining balance method. If $R = 1.5$ the program gives the 150% declining balance method.

DISPLAY		KEY ENTRY	DISPLAY		KEY ENTRY	REGISTERS
LINE	CODE		LINE	CODE		
00.			25.	03	3	R ₀
01.	01	1	26.	71	x	R ₁ R/n
02.	34	RCL	27.	84	R/S	R ₂
03.	01	1	28.	01	1	R ₃ PV
04.	51	—	29.	33	STO	R ₄ k
05.	34	RCL	30.	61	+	R ₅
06.	04	4	31.	04	4	R ₆
07.	01	1	32.	-01	GTO 01	R ₇
08.	51	—	33.			R ₈
09.	12	y ^x	34.			R ₉
10.	34	RCL	35.			R ₀₀
11.	01	1	36.			R ₀₁
12.	71	x	37.			R ₀₂
13.	34	RCL	38.			R ₀₃
14.	03	3	39.			R ₀₄
15.	71	x	40.			R ₀₅
16.	84	R/S	41.			R ₀₆
17.	01	1	42.			R ₀₇
18.	34	RCL	43.			R ₀₈
19.	01	1	44.			R ₀₉
20.	51	—	45.			
21.	34	RCL	46.			
22.	04	4	47.			
23.	12	y ^x	48.			
24.	34	RCL	49.			

74 Depreciation Schedules—Variable Rate Declining Balance

Example:

A fleet car has a value of \$2500 and a life expectancy of six years. Use the double declining balance method ($R = 2$) to find the amount of depreciation and book value after four years.

Solutions:

$$D_4 = \$246.91$$

$$B_4 = \$493.83$$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Store R/n, PV, k	R	↑				
		n	÷	STO	1		
		PV	STO	3			
		k	STO	4	BST		
3	Calculate D_k and B_k		R/S				D_k
			R/S				B_k
4	For next year go to step 3						

CALENDAR ROUTINES

DAY OF THE WEEK

DAYS BETWEEN TWO DATES

This program calls March 1, 1700, day 1 and gives every succeeding day a corresponding number. The program works for days to and including February 28, 2100. However, for days from March 1, 1700, to February 28, 1800, 2 days must be added to the answer and for days from March 1, 1800, to February 18, 1900, 1 day must be added.

Let M = month, D = day, Y = year, W = day of the week (0 = Sunday, 1 = Monday, etc.)

The day's number is calculated from the following formula:

$$N(M, D, Y) = [365.25 g(y, m)] + [30.6 f(m)] + D - 621049$$

where

$$g(y, m) = \begin{cases} y - 1 & \text{if } m = 1 \text{ or } 2 \\ y & \text{if } m > 2 \end{cases} \quad \text{and} \quad f(m) = \begin{cases} m + 13 & \text{if } m = 1 \text{ or } 2 \\ m + 1 & \text{if } m > 2 \end{cases}$$

[m] represents the integer part of a number, i.e., if $n = 7.2$ then $[7.2] = 7$. This must be put in by user.

76 Calendar Routines

DISPLAY			KEY ENTRY	DISPLAY			KEY ENTRY	REGISTERS
LINE	CODE			LINE	CODE			
00.				25.	84	R/S		R ₀
01.	02	2		26.	22	$x \geq y$		R ₁ Month
02.	34	RCL		27.	23	R↓		R ₂ Day
03.	01	1		28.	22	$x \geq y$		R ₃ Year
04.	31	f		29.	03	3		R ₄
05.	-11	$x \leq y$ 11		30.	00	0		R ₅
06.	01	1		31.	83	.		R ₆
07.	61	+		32.	06	6		R ₇
08.	34	RCL		33.	71	x		R ₈ Temporary
09.	03	3		34.	84	R/S		R ₉
10.	-18	GTO 18		35.	22	$x \geq y$		R ₀₀
11.	01	1		36.	23	R↓		R ₀₁
12.	03	3		37.	61	+		R ₀₂
13.	61	+		38.	34	RCL		R ₀₃
14.	34	RCL		39.	02	2		R ₀₄
15.	03	3		40.	61	+		R ₀₅
16.	01	1		41.	06	6		R ₀₆
17.	51	-		42.	02	2		R ₀₇
18.	03	3		43.	01	1		R ₀₈
19.	06	6		44.	00	0		R ₀₉
20.	05	5		45.	04	4		
21.	83	.		46.	09	9		
22.	02	2		47.	51	-		
23.	05	5		48.	-00	GTO 00		
24.	71	x		49.				

DETERMINANT AND INVERSE OF A 2 × 2 MATRIX

Let $A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}$ be a 2 × 2 matrix.

The determinant of A denoted by Det A or |A| is evaluated by the following formula:

$$\text{Det } A = a_{22} a_{11} - a_{12} a_{21}$$

Also, the program evaluates the multiplicative inverse A^{-1} of A. The following formulas are used:

$$A^{-1} = \begin{bmatrix} a_{22}/\text{Det } A & -a_{12}/\text{Det } A \\ -a_{21}/\text{Det } A & a_{11}/\text{Det } A \end{bmatrix}$$

DISPLAY			KEY ENTRY	DISPLAY			KEY ENTRY	REGISTERS
LINE	CODE			LINE	CODE			
00.				25.	09	9		R ₀ Det A
01.	34	RCL		26.	81	÷		R ₁ a ₁₁
02.	04	4		27.	33	STO		R ₂ a ₁₂
03.	34	RCL		28.	08	8		R ₃ a ₂₁
04.	01	1		29.	34	RCL		R ₄ a ₂₂
05.	71	x		30.	02	2		R ₅ a ₁₁ ⁻¹
06.	34	RCL		31.	34	RCL		R ₆ a ₁₂ ⁻¹
07.	02	2		32.	09	9		R ₇ a ₂₁ ⁻¹
08.	34	RCL		33.	81	÷		R ₈ a ₂₂ ⁻¹
09.	03	3		34.	42	CHS		R ₉ Det A
10.	71	x		35.	33	STO		R ₀₀
11.	51	—		36.	06	6		R ₀₁
12.	33	STO		37.	34	RCL		R ₀₂
13.	09	9		38.	03	3		R ₀₃
14.	84	R/S		39.	34	RCL		R ₀₄
15.	34	RCL		40.	09	9		R ₀₅
16.	04	4		41.	81	÷		R ₀₆
17.	34	RCL		42.	42	CHS		R ₀₇
18.	09	9		43.	33	STO		R ₀₈
19.	81	÷		44.	07	7		R ₀₉
20.	33	STO		45.	-00	GTO 00		
21.	05	5		46.				
22.	34	RCL		47.				
23.	01	1		48.				
24.	34	RCL		49.				

Example:

Find the determinant and inverse of the matrix

$$A = \begin{bmatrix} 1 & 4 \\ 3 & 2 \end{bmatrix}$$

Solution:

$$\text{Det } A = -10$$

$$A^{-1} = \begin{bmatrix} -.20 & .40 \\ .30 & -.10 \end{bmatrix}$$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	For Det A	a_{11}	STO	1			
		a_{12}	STO	2			
		a_{21}	STO	3			
		a_{22}	STO	4			
			BST	R/S			Det A
3	Then for A^{-1}		R/S				
			RCL	5			a_{11}^{-1}
			RCL	6			a_{12}^{-1}
			RCL	7			a_{21}^{-1}
			RCL	8			a_{22}^{-1}

DETERMINANT OF A 3 x 3 MATRIX

Let $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$ be a 3 x 3 matrix.

The determinant of A denoted by $|A|$ or $\text{Det } A$, is calculated by expanding A by minors about the first column. The formula is:

$$\begin{aligned} \text{Det } A &= a_{11} \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix} - a_{21} \begin{vmatrix} a_{12} & a_{13} \\ a_{32} & a_{33} \end{vmatrix} + a_{31} \begin{vmatrix} a_{12} & a_{13} \\ a_{22} & a_{23} \end{vmatrix} \\ &= a_{11} [a_{22} a_{33} - a_{23} a_{32}] - a_{21} [a_{33} a_{12} - a_{32} a_{13}] \\ &\quad + a_{31} [a_{23} a_{12} - a_{13} a_{22}] \end{aligned}$$

DISPLAY			KEY ENTRY			REGISTERS
LINE	CODE		LINE	CODE		
00.			25.	51	—	R ₀ Det A
01.	34	RCL	26.	34	RCL	R ₁ a ₁₁
02.	05	5	27.	04	4	R ₂ a ₁₂
03.	34	RCL	28.	71	x	R ₃ a ₁₃
04.	09	9	29.	51	—	R ₄ a ₂₁
05.	71	x	30.	34	RCL	R ₅ a ₂₂
06.	34	RCL	31.	06	6	R ₆ a ₂₃
07.	06	6	32.	34	RCL	R ₇ a ₃₁
08.	34	RCL	33.	02	2	R ₈ a ₃₂
09.	08	8	34.	71	x	R ₉ a ₃₃
10.	71	x	35.	34	RCL	R ₀₀
11.	51	—	36.	03	3	R ₀₁
12.	34	RCL	37.	34	RCL	R ₀₂
13.	01	1	38.	05	5	R ₀₃
14.	71	x	39.	71	x	R ₀₄
15.	34	RCL	40.	51	—	R ₀₅
16.	09	9	41.	34	RCL	R ₀₆
17.	34	RCL	42.	07	7	R ₀₇
18.	02	2	43.	71	x	R ₀₈
19.	71	x	44.	61	+	R ₀₉
20.	34	RCL	45.	33	STO	
21.	08	8	46.	00	0	
22.	34	RCL	47.	-00	GTO 00	
23.	03	3	48.			
24.	71	x	49.			

Example:

Find the Determinant of

$$A = \begin{bmatrix} -1 & 0 & 3 \\ 7 & 1 & -1 \\ 2 & 3 & 0 \end{bmatrix}$$

Solution:

Det A = 54

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Store A	a ₁₁	STO	1			
		a ₁₂	STO	2			
		a ₁₃	STO	3			
		a ₂₁	STO	4			
		a ₂₂	STO	5			
		a ₂₃	STO	6			
		a ₃₁	STO	7			
		a ₃₂	STO	8			
		a ₃₃	STO	9			
3	Calculate Det A		BST	R/S			Det A

3 × 3 MATRIX INVERSION

If a_{ij} indicates a number in the i^{th} row, j^{th} column then a 3×3 matrix A can be represented as

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$$

then the multiplicative inverse of A is denoted by A^{-1} and is calculated as follows:

$$A^{-1} = \begin{bmatrix} \frac{\begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix}}{\text{Det } A} & -\frac{\begin{vmatrix} a_{12} & a_{13} \\ a_{32} & a_{33} \end{vmatrix}}{\text{Det } A} & \frac{\begin{vmatrix} a_{12} & a_{13} \\ a_{22} & a_{23} \end{vmatrix}}{\text{Det } A} \\ -\frac{\begin{vmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{vmatrix}}{\text{Det } A} & \frac{\begin{vmatrix} a_{11} & a_{13} \\ a_{31} & a_{33} \end{vmatrix}}{\text{Det } A} & -\frac{\begin{vmatrix} a_{11} & a_{13} \\ a_{21} & a_{23} \end{vmatrix}}{\text{Det } A} \\ \frac{\begin{vmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{vmatrix}}{\text{Det } A} & -\frac{\begin{vmatrix} a_{11} & a_{12} \\ a_{31} & a_{32} \end{vmatrix}}{\text{Det } A} & \frac{\begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix}}{\text{Det } A} \end{bmatrix}$$

For the i^{th} , j^{th} position of A^{-1} use the minor of the j^{th} , i^{th} position of the original matrix. The minor is the two by two matrix left after crossing out the i^{th} row and j^{th} column of A .

DISPLAY		KEY ENTRY	DISPLAY		KEY ENTRY	REGISTERS
LINE	CODE		LINE	CODE		
00.			25.			R ₀ Det A
01.	23	R↓	26.			R ₁ a ₁₁
02.	71	x	27.			R ₂ a ₁₂
03.	41	↑	28.			R ₃ a ₁₃
04.	23	R↓	29.			R ₄ a ₂₁
05.	23	R↓	30.			R ₅ a ₂₂
06.	71	x	31.			R ₆ a ₂₃
07.	51	—	32.			R ₇ a ₃₁
08.	42	CHS	33.			R ₈ a ₃₂
09.	34	RCL	34.			R ₉ a ₃₃
10.	00	0	35.			R ₀₀
11.	81	÷	36.			R ₀₁
12.	-00	GTO 00	37.			R ₀₂
13.			38.			R ₀₃
14.			39.			R ₀₄
15.			40.			R ₀₅
16.			41.			R ₀₆
17.			42.			R ₀₇
18.			43.			R ₀₈
19.			44.			R ₀₉
20.			45.			
21.			46.			
22.			47.			
23.			48.			
24.			49.			

84 3 x 3 Matrix Inversion

Example:

Find the inverse of the matrix

$$A = \begin{bmatrix} -1 & 0 & 3 \\ 7 & 1 & -1 \\ 2 & 3 & 0 \end{bmatrix}$$

Solution:

$$\text{Det } A = 54$$

$$A^{-1} = \begin{bmatrix} .056 & .167 & -.056 \\ -.037 & -.111 & .370 \\ .352 & .056 & -.019 \end{bmatrix}$$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Follow instruction of Determinant of 3 x 3 Matrix program						
2	Enter this program (do not change contents of registers.)						
3	To find A^{-1} (Note: coefficients must be calculated in order shown.)		RCL	5	RCL	6	
			RCL	8	RCL	9	
			BST	R/S			a_{11}^{-1}
			RCL	8	RCL	9	
			RCL	2	RCL	3	
			R/S				a_{12}^{-1}
			RCL	2	RCL	3	
			RCL	5	RCL	6	
			R/S				a_{13}^{-1}
			RCL	7	RCL	9	
			RCL	4	RCL	6	
			R/S				a_{21}^{-1}
			RCL	1	RCL	3	
			RCL	7	RCL	9	
			R/S				a_{22}^{-1}
			RCL	4	RCL	6	
			RCL	1	RCL	3	
			R/S				a_{23}^{-1}
			RCL	4	RCL	5	
			RCL	7	RCL	8	
			R/S				a_{31}^{-1}
			RCL	7	RCL	8	
			RCL	1	RCL	2	
			R/S				a_{32}^{-1}
			RCL	1	RCL	2	
			RCL	4	RCL	5	
			R/S				a_{33}^{-1}

VECTOR CROSS PRODUCT

If $A = (a_1, a_2, a_3)$ and $B = (b_1, b_2, b_3)$ are two three dimensional vectors then the cross product of A and B is denoted by $A \times B$ and is calculated as follows:

$$A \times B = \left(\begin{vmatrix} a_2 & a_3 \\ b_2 & b_3 \end{vmatrix}, -\begin{vmatrix} a_1 & a_3 \\ b_1 & b_3 \end{vmatrix}, \begin{vmatrix} a_1 & a_2 \\ b_1 & b_2 \end{vmatrix} \right) = (a_2 b_3 - a_3 b_2, a_3 b_1 - a_1 b_3, a_1 b_2 - a_2 b_1)$$

Let the solution be represented by (c_1, c_2, c_3) .

DISPLAY			KEY ENTRY			REGISTERS
LINE	CODE	KEY ENTRY	LINE	CODE	KEY ENTRY	
00.			25.	34	RCL	R ₀
01.	34	RCL	26.	01	1	R ₁ a ₁
02.	02	2	27.	34	RCL	R ₂ a ₂
03.	34	RCL	28.	05	5	R ₃ a ₃
04.	06	6	29.	71	x	R ₄ b ₁
05.	71	x	30.	34	RCL	R ₅ b ₂
06.	34	RCL	31.	02	2	R ₆ b ₃
07.	03	3	32.	34	RCL	R ₇
08.	34	RCL	33.	04	4	R ₈
09.	05	5	34.	71	x	R ₉
10.	71	x	35.	51	—	R ₀₀
11.	51	—	36.	-00	GTO 00	R ₀₁
12.	84	R/S	37.			R ₀₂
13.	34	RCL	38.			R ₀₃
14.	03	3	39.			R ₀₄
15.	34	RCL	40.			R ₀₅
16.	04	4	41.			R ₀₆
17.	71	x	42.			R ₀₇
18.	34	RCL	43.			R ₀₈
19.	01	1	44.			R ₀₉
20.	34	RCL	45.			
21.	06	6	46.			
22.	71	x	47.			
23.	51	—	48.			
24.	84	R/S	49.			

Example:

Find the cross product of the two vectors $A = (2.34, 5.17, 7.43)$ and $B = (.072, .231, .409)$.

Solution:

$$A \times B = (.40, -.42, .17)$$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Store A	a_1	STO	1			
		a_2	STO	2			
		a_3	STO	3			
3	Store B	b_1	STO	4			
		b_2	STO	5			
		b_3	STO	6			
4	Calculate $A \times B$		BST	R/S			c_1
			R/S				c_2
			R/S				c_3

**SIMULTANEOUS EQUATIONS
IN TWO UNKNOWN**

Let $ax + by = e$

and $cx + dy = f$

be a system of two equations in two unknowns. Cramer's Rule is used to find the solution.

$$x = \frac{\begin{vmatrix} e & b \\ f & d \end{vmatrix}}{\begin{vmatrix} a & b \\ c & d \end{vmatrix}} = \frac{ed - bf}{ad - bc}$$

$$y = \frac{\begin{vmatrix} a & e \\ c & f \end{vmatrix}}{\begin{vmatrix} a & b \\ c & d \end{vmatrix}} = \frac{af - ec}{ad - bc}$$

If $ad - bc = 0$ the calculator flashes 0. In this case no solution or no unique solution exists.

DISPLAY			DISPLAY			KEY ENTRY	REGISTERS
LINE	CODE	KEY ENTRY	LINE	CODE	KEY ENTRY		
00.			25.	81	÷		R ₀ ad - bc
01.	34	RCL	26.	84	R/S		R ₁ a
02.	03	3	27.	34	RCL		R ₂ b
03.	34	RCL	28.	01	1		R ₃ e
04.	05	5	29.	34	RCL		R ₄ c
05.	71	x	30.	06	6		R ₅ d
06.	34	RCL	31.	71	x		R ₆ f
07.	02	2	32.	34	RCL		R ₇
08.	34	RCL	33.	03	3		R ₈
09.	06	6	34.	34	RCL		R ₉
10.	71	x	35.	04	4		R ₀₀
11.	51	-	36.	71	x		R ₀₁
12.	34	RCL	37.	51	-		R ₀₂
13.	01	1	38.	34	RCL		R ₀₃
14.	34	RCL	39.	00	0		R ₀₄
15.	05	5	40.	81	÷		R ₀₅
16.	71	x	41.	-00	GTO 00		R ₀₆
17.	34	RCL	42.				R ₀₇
18.	02	2	43.				R ₀₈
19.	34	RCL	44.				R ₀₉
20.	04	4	45.				
21.	71	x	46.				
22.	51	-	47.				
23.	33	STO	48.				
24.	00	0	49.				

90 Simultaneous Equations in Two Unknowns

Example:

Solve for x and y the following system of equations.

$$7.32x - 9.08y = 3.14$$

$$12.39x + 7.00y = .05$$

Solution:

$$x = .14$$

$$y = -.24$$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Store coefficients	a	STO	1			
		b	STO	2			
		e	STO	3			
		c	STO	4			
		d	STO	5			
		f	STO	6			
3	Find x and y		BST	R/S			x
			R/S				y

SIMULTANEOUS EQUATIONS IN THREE UNKNOWNNS

$$\text{Let} \quad a_1 x + b_1 y + c_1 z = d_1$$

$$a_2 x + b_2 y + c_2 z = d_2$$

$$a_3 x + b_3 y + c_3 z = d_3$$

be a system of three equations in three unknowns. Cramer's Rule is used to find the solution.

$$x = \frac{\begin{vmatrix} d_1 & b_1 & c_1 \\ d_2 & b_2 & c_2 \\ d_3 & b_3 & c_3 \end{vmatrix}}{\text{Det } A} \quad y = \frac{\begin{vmatrix} a_1 & d_1 & c_1 \\ a_2 & d_2 & c_2 \\ a_3 & d_3 & c_3 \end{vmatrix}}{\text{Det } A} \quad z = \frac{\begin{vmatrix} a_1 & b_1 & d_1 \\ a_2 & b_2 & d_2 \\ a_3 & b_3 & d_3 \end{vmatrix}}{\text{Det } A}$$

where $\begin{vmatrix} \end{vmatrix}$ and Det represent the determinant and

$$A = \begin{bmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{bmatrix}$$

If $\text{Det } A = 0$ in step 3 the system has no solution or no unique solution. Continuing the program will cause the calculator to flash zero.

92 Simultaneous Equations in Three Unknowns

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program "Determinant of 3 x 3 Matrix"						
2	Store coefficients	a_1	STO	1			
		b_1	STO	2			
		c_1	STO	3			
		a_2	STO	4			
		b_2	STO	5			
		c_2	STO	6			
		a_3	STO	7			
		b_3	STO	8			
		c_3	STO	9			
3	Calculate Det A		BST	R/S			Det A
4	Store Det A		STO	.	1		
5	Store constants	d_1	STO	1			
		d_2	STO	4			
		d_3	STO	7			
6	Calculate x		R/S	RCL	.	1	
			÷				x
7	Calculate y	a_1	STO	2			
		a_2	STO	5			
		a_3	STO	8			
			R/S	RCL	.	1	
			÷	CHS			y
8	Calculate z	b_1	STO	3			
		b_2	STO	6			
		b_3	STO	9			
			R/S	RCL	.	1	
			÷				z
	(If the coefficients are quite long it may be useful at step 2 to also store $a_1, b_1, a_2, b_2, a_3,$ b_3 in $R_{02}, R_{03}, \dots, R_{07}$ respec- tively and then recall them in steps 7 and 8 as needed.)						

Example:

Solve the following system for x , y , and z .

$$x + y + z = 6$$

$$3x + 5y + 4z = 23$$

$$6x - 7y - 2z = 2$$

Solution:

$$\text{Det } A = -3$$

$$x = 3, \quad y = 2, \quad z = 1$$

2 × 2 MATRIX MULTIPLICATION

Let

$$A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{bmatrix}$$

be two 2×2 matrices. The matrix product of A and B is calculated as follows:

$$AB = \begin{bmatrix} a_{11} b_{11} + a_{12} b_{21} & a_{11} b_{12} + a_{12} b_{22} \\ a_{21} b_{11} + a_{22} b_{21} & a_{21} b_{12} + a_{22} b_{22} \end{bmatrix}$$

Let the answer be denoted by:

$$C = \begin{bmatrix} c_{11} & c_{12} \\ c_{21} & c_{22} \end{bmatrix}$$

DISPLAY			KEY ENTRY	DISPLAY			KEY ENTRY	REGISTERS
LINE	CODE			LINE	CODE			
00.				25.	34		RCL	R ₀
01.	34		RCL	26.	03		3	R ₁ a ₁₁
02.	01		1	27.	34		RCL	R ₂ a ₁₂
03.	34		RCL	28.	05		5	R ₃ a ₂₁
04.	05		5	29.	71		x	R ₄ a ₂₂
05.	71		x	30.	34		RCL	R ₅ b ₁₁
06.	34		RCL	31.	04		4	R ₆ b ₁₂
07.	02		2	32.	34		RCL	R ₇ b ₂₁
08.	34		RCL	33.	07		7	R ₈ b ₂₂
09.	07		7	34.	71		x	R ₉
10.	71		x	35.	61		+	R ₀₀
11.	61		+	36.	84		R/S	R ₀₁
12.	84		R/S	37.	34		RCL	R ₀₂
13.	34		RCL	38.	03		3	R ₀₃
14.	01		1	39.	34		RCL	R ₀₄
15.	34		RCL	40.	06		6	R ₀₅
16.	06		6	41.	71		x	R ₀₆
17.	71		x	42.	34		RCL	R ₀₇
18.	34		RCL	43.	04		4	R ₀₈
19.	02		2	44.	34		RCL	R ₀₉
20.	34		RCL	45.	08		8	
21.	08		8	46.	71		x	
22.	71		x	47.	61		+	
23.	61		+	48.	-00		GTO 00	
24.	84		R/S	49.				

Example:

Find the product of the two matrices

$$A = \begin{bmatrix} 1 & 2 \\ -1 & 3 \end{bmatrix} \text{ and } B = \begin{bmatrix} 1 & -1 \\ 2 & 4 \end{bmatrix}$$

Solution:

$$C = \begin{bmatrix} 5 & 7 \\ 5 & 13 \end{bmatrix}$$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Store A	a_{11}	STO	1			
		a_{12}	STO	2			
		a_{21}	STO	3			
		a_{22}	STO	4			
3	Store B	b_{11}	STO	5			
		b_{12}	STO	6			
		b_{21}	STO	7			
		b_{22}	STO	8			
4	Calculate C		BST	R/S			c_{11}
			R/S				c_{12}
			R/S				c_{21}
			R/S				c_{22}

ANGLE BETWEEN, NORM, AND DOT PRODUCT OF VECTORS

Let $\vec{a} = (a_1, a_2, \dots, a_n)$ and $\vec{b} = (b_1, b_2, \dots, b_n)$ be two vectors.

The norm of $\vec{a}$ is denoted by $|\vec{a}|$ and is calculated by the following formula:

$$|\vec{a}| = \sqrt{a_1^2 + a_2^2 + \dots + a_n^2}$$

similarly,

$$|\vec{b}| = \sqrt{b_1^2 + b_2^2 + \dots + b_n^2}$$

The dot product of $\vec{a}$ and $\vec{b}$ is denoted by $\vec{a} \cdot \vec{b}$ and is calculated by the following formula:

$$\vec{a} \cdot \vec{b} = a_1 b_1 + a_2 b_2 + \dots + a_n b_n$$

The angle between a and b is denoted by θ and is calculated by the following formula:

$$\theta = \cos^{-1} \left(\frac{\vec{a} \cdot \vec{b}}{|\vec{a}| \cdot |\vec{b}|} \right)$$

The angle is calculated in any mode the calculator is set. However, if in degrees, decimal degrees are assumed.

DISPLAY			DISPLAY			KEY ENTRY	REGISTERS
LINE	CODE	KEY ENTRY	LINE	CODE	KEY ENTRY		
00.			25.	81	$\div$		R_0
01.	34	RCL	26.	32	g		R_1
02.	83	$\cdot$	27.	13	$\cos^{-1}$		R_2
03.	02	2	28.	-00	GTO 00		R_3
04.	31	f	29.				R_4
05.	42	$\sqrt{x}$	30.				R_5
06.	-00	GTO 00	31.				R_6
07.	34	RCL	32.				R_7
08.	83	$\cdot$	33.				R_8
09.	04	4	34.				R_9
10.	31	f	35.				$R_{00} n$
11.	42	$\sqrt{x}$	36.				$R_{01} \sum a_i$
12.	-00	GTO 00	37.				$R_{02} \sum a_i^2$
13.	34	RCL	38.				$R_{03} \sum b_i$
14.	83	$\cdot$	39.				$R_{04} \sum b_i^2$
15.	05	5	40.				$R_{05} \sum a_i b_i$
16.	34	RCL	41.				R_{06}
17.	83	$\cdot$	42.				R_{07}
18.	02	2	43.				R_{08}
19.	34	RCL	44.				R_{09}
20.	83	$\cdot$	45.				
21.	04	4	46.				
22.	71	x	47.				
23.	31	f	48.				
24.	42	$\sqrt{x}$	49.				

98 Angle Between, Norm, and Dot Product of Vectors

Example:

Let $\vec{a} = (2.34, 5.17, 7.43)$ and $\vec{b} = (.07, .23, .41)$, find the norms of $\vec{a}$ and $\vec{b}$, the dot product of $\vec{a}$ and $\vec{b}$, and the angle between $\vec{a}$ and $\vec{b}$.

Solution:

$$|\vec{a}| = 9.35$$

$$|\vec{b}| = .48$$

$$\vec{a} \cdot \vec{b} = 4.40$$

$$\theta = \cos^{-1} \left(\frac{\vec{a} \cdot \vec{b}}{|\vec{a}| \cdot |\vec{b}|} \right) = 8.11^\circ = .14 \text{ radians} = 9.01 \text{ grads}$$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Initialize		g	CL·R			
3	Perform for i = 1, 2, ..., n	b_i	↑				
		a_i	Σ+				i
4	For $ \vec{a} $		BST	R/S			$ \vec{a} $
	or						
	$ \vec{b} $		GTO	0	7	R/S	$ \vec{b} $
	or						
	$\vec{a} \cdot \vec{b}$		RCL	·	5		$\vec{a} \cdot \vec{b}$
	or						
	θ		GTO	1	3	R/S	θ

SINE INTEGRAL

The sine integral is denoted by $\text{Si}(x)$ and is defined as follows:

$$\text{Si}(x) = \int_0^x \frac{\sin t}{t} dt$$

where x is a real number. Also, a Taylor's series expansion of $\text{Si}(x)$ yields

$$\text{Si}(x) = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{(2n+1)(2n+1)!}$$

This program computes successive partial sums of the series. It stops when two consecutive partial sums are equal, and displays the last partial sum as the answer.

DISPLAY		KEY ENTRY	DISPLAY		KEY ENTRY	REGISTERS
LINE	CODE		LINE	CODE		
00.			25.	02	2	R_0
01.	33	STO	26.	81	$\div$	$R_1 -x^2$
02.	03	3	27.	34	RCL	$R_2 2n+1$
03.	41	$\uparrow$	28.	03	3	R_3 Used
04.	71	$\times$	29.	71	$\times$	R_4
05.	42	CHS	30.	33	STO	R_5
06.	33	STO	31.	03	3	R_6
07.	01	1	32.	34	RCL	R_7
08.	01	1	33.	02	2	R_8
09.	33	STO	34.	81	$\div$	R_9
10.	02	2	35.	61	+	R_{10}
11.	34	RCL	36.	32	g	R_{11}
12.	03	3	37.	-00	$x=y$ 00	R_{12}
13.	34	RCL	38.	-13	GTO 13	R_{13}
14.	01	1	39.			R_{14}
15.	34	RCL	40.			R_{15}
16.	02	2	41.			R_{16}
17.	01	1	42.			R_{17}
18.	61	+	43.			R_{18}
19.	81	$\div$	44.			R_{19}
20.	31	f	45.			
21.	34	LAST X	46.			
22.	01	1	47.			
23.	61	+	48.			
24.	33	STO	49.			

Examples:

1. $Si(.69) = .67$ (15 seconds iteration)
2. $Si(9.8) = 1.67$ (50 seconds iteration)

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Calculate $Si(x)$	x	BST	R/S			$Si(x)$

COSINE INTEGRAL

The cosine integral is denoted by $\text{Ci}(x)$ and is defined as follows:

$$\text{Ci}(x) = \gamma + \ln x + \int_0^x \frac{\cos t - 1}{t} dt$$

where $x > 0$, and $\gamma = 0.5772156649$ is Euler's constant.

Also, a Taylor series expansions yields

$$\text{Ci}(x) = \gamma + \ln x + \sum_{n=1}^{\infty} \frac{(-1)^n x^{2n}}{2n (2n)!}$$

This program computes successive partial sums of the series. When two consecutive partial sums are equal, the value is used as the sum of the series.

DISPLAY		KEY ENTRY	DISPLAY		KEY ENTRY	REGISTERS
LINE	CODE		LINE	CODE		
00.			25.	81	÷	R ₀
01.	41	↑	26.	31	f	R ₁ -x ²
02.	71	x	27.	34	LAST X	R ₂ 2n
03.	42	CHS	28.	01	1	R ₃ Used
04.	33	STO	29.	61	+	R ₄ γ
05.	01	1	30.	33	STO	R ₅
06.	01	1	31.	02	2	R ₆
07.	33	STO	32.	81	÷	R ₇
08.	03	3	33.	34	RCL	R ₈
09.	00	0	34.	03	3	R ₉
10.	33	STO	35.	71	x	R ₀₀
11.	02	2	36.	33	STO	R ₀₁
12.	31	f	37.	03	3	R ₀₂
13.	34	LAST X	38.	34	RCL	R ₀₃
14.	31	f	39.	02	2	R ₀₄
15.	22	ln	40.	81	÷	R ₀₅
16.	34	RCL	41.	61	+	R ₀₆
17.	04	4	42.	32	g	R ₀₇
18.	61	+	43.	-00	x=y 00	R ₀₈
19.	34	RCL	44.	-19	GTO 19	R ₀₉
20.	01	1	45.			
21.	34	RCL	46.			
22.	02	2	47.			
23.	01	1	48.			
24.	61	+	49.			

Examples:

1. $Ci(1.38) = .46$ (20 seconds iteration)
2. $Ci(5) = -.19$ (40 seconds iteration)

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Store $\gamma = .5772156649$		.	5	7	7	
			2	1	5	6	
			6	4	9		
			STO	4			
3	Calculate $Ci(x)$	x	BST	R/S			$Ci(x)$

EXPONENTIAL INTEGRAL

The exponential integral is denoted by $Ei(x)$ and is defined as follows:

$$Ei(x) = \int_{-\infty}^x \frac{e^t}{t} dt$$

where $x > 0$.

Using a Taylor's series expansion and letting $\gamma = 0.5772156649$ be Euler's constant:

$$Ei(x) = \gamma + \ln x + \sum_{n=1}^{\infty} \frac{x^n}{n(n!)}$$

This program computes successive partial sums of the series. When two consecutive partial sums are equal, the value is used as the sum of the series.

DISPLAY			KEY ENTRY	DISPLAY			KEY ENTRY	REGISTERS
LINE	CODE			LINE	CODE			
00.				25.	34	RCL		R ₀
01.	33	STO		26.	01	1		R ₁ x
02.	01	1		27.	34	RCL		R ₂ Used
03.	01	1		28.	02	2		R ₃ Used
04.	33	STO		29.	01	1		R ₄
05.	03	3		30.	61	+		R ₅
06.	00	0		31.	33	STO		R ₆
07.	33	STO		32.	02	2		R ₇
08.	02	2		33.	81	÷		R ₈
09.	34	RCL		34.	34	RCL		R ₉
10.	01	1		35.	03	3		R ₁₀
11.	31	f		36.	71	x		R ₁₁
12.	22	ln		37.	33	STO		R ₁₂
13.	83	.		38.	03	3		R ₁₃
14.	05	5		39.	34	RCL		R ₁₄
15.	07	7		40.	02	2		R ₁₅
16.	07	7		41.	81	÷		R ₁₆
17.	02	2		42.	61	+		R ₁₇
18.	01	1		43.	32	g		R ₁₈
19.	05	5		44.	-00	x=y 00		R ₁₉
20.	06	6		45.	-25	GTO 25		
21.	06	6		46.				
22.	04	4		47.				
23.	09	9		48.				
24.	61	+		49.				

Examples:

1. $Ei(1.59) = 3.57$ (35 seconds iteration)
2. $Ei(.61) = .80$ (25 seconds iteration)

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Compute $Ei(x)$	x	BST	R/S			$Ei(x)$

NUMERICAL INTEGRATION, TRAPEZOIDAL RULE

Let $x_0, x_1, \dots, x_n$ be equally spaced points such that $x_i = x_0 + ih$ for $i = 0, 1, 2, \dots, n$, at which corresponding values $f(x_0), f(x_1), \dots, f(x_n)$ of a function $f(x)$ are known. The function need not be known explicitly but if it is, these values can be found previously by writing the function into memory and evaluating at the various points. R_2 through R_{09} could be used to store these values.

The Trapezoidal Rule is:

$$\int_{x_0}^{x_n} f(x) dx \cong \frac{h}{2} \left[f(x_0) + 2 \sum_{i=1}^{n-1} f(x_i) + f(x_n) \right]$$

Let the answer be represented by I.

DISPLAY		KEY ENTRY	DISPLAY		KEY ENTRY	REGISTERS
LINE	CODE		LINE	CODE		
00.			25.	34	RCL	R_0 $h/2$
01.	02	2	26.	00	0	R_1 Σ
02.	81	$\div$	27.	71	$\times$	R_2
03.	33	STO	28.	34	RCL	R_3
04.	00	0	29.	01	1	R_4
05.	84	R/S	30.	61	+	R_5
06.	34	RCL	31.	33	STO	R_6
07.	00	0	32.	01	1	R_7
08.	71	$\times$	33.	-24	GTO 24	R_8
09.	33	STO	34.			R_9
10.	01	1	35.			R_{00}
11.	84	R/S	36.			R_{01}
12.	34	RCL	37.			R_{02}
13.	00	0	38.			R_{03}
14.	71	$\times$	39.			R_{04}
15.	33	STO	40.			R_{05}
16.	61	+	41.			R_{06}
17.	01	1	42.			R_{07}
18.	02	2	43.			R_{08}
19.	33	STO	44.			R_{09}
20.	71	$\times$	45.			
21.	00	0	46.			
22.	34	RCL	47.			
23.	01	1	48.			
24.	84	R/S	49.			

Example:

Find the $\int_0^2 x^3 dx$ using $h = .25$.

The following data must be found first:

i	0	1	2	3	4	5	6	7	8
x_i	0	.25	.50	.75	1.00	1.25	1.50	1.75	2.00
$f(x_i)$	0	.0156	.1250	.4219	1.0000	1.9531	3.3750	5.3594	8.0000

Solution:

$$\int_0^2 x^3 dx \cong 4.0625$$

Actual solution is 4.

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Enter h	h	BST	R/S			h/2
3	Enter $f(x_0)$	$f(x_0)$	R/S				Partial Sum
4	Enter $f(x_n)$	$f(x_n)$	R/S				Partial Sum
5	Perform step 5 for $i = 1, 2, \dots, n-2$	$f(x_i)$	R/S				Partial Sum
6	Enter $f(x_{n-1})$	$f(x_{n-1})$	R/S				I

NUMERICAL INTEGRATION, SIMPSON'S RULE

Let $x_0, x_1, \dots, x_n$ be equally spaced points such that $x_i = x_0 + ih$ for $i = 0, 1, 2, \dots, n$ at which corresponding values $f(x_0), f(x_1), \dots, f(x_n)$ of a function $f(x)$ are known. This function need not be known explicitly but if it is, these values can be found previously by writing the function into memory and evaluating at the various points. R_2 through R_{99} could be used to store these values. n must be an even positive integer.

Simpson's Rule is:

$$\int_{x_0}^{x_n} f(x) dx \cong \frac{h}{3} [f(x_0) + 4f(x_1) + 2f(x_2) + \dots + 4f(x_{n-3}) + 2f(x_{n-2}) + 4f(x_{n-1}) + f(x_n)].$$

Let the solution be indicated by I.

DISPLAY		KEY ENTRY	DISPLAY		KEY ENTRY	REGISTERS
LINE	CODE		LINE	CODE		
00.			25.	71	x	R_0 $h/3$
01.	03	3	26.	34	RCL	R_1 Σ
02.	81	$\div$	27.	01	1	R_2
03.	33	STO	28.	61	+	R_3
04.	00	0	29.	33	STO	R_4
05.	84	R/S	30.	01	1	R_5
06.	34	RCL	31.	84	R/S	R_6
07.	00	0	32.	34	RCL	R_7
08.	71	x	33.	00	0	R_8
09.	33	STO	34.	71	x	R_9
10.	01	1	35.	02	2	R_{00}
11.	84	R/S	36.	71	x	R_{01}
12.	34	RCL	37.	34	RCL	R_{02}
13.	00	0	38.	01	1	R_{03}
14.	71	x	39.	61	+	R_{04}
15.	34	RCL	40.	33	STO	R_{05}
16.	01	1	41.	01	1	R_{06}
17.	61	+	42.	-20	GTO 20	R_{07}
18.	33	STO	43.			R_{08}
19.	01	1	44.			R_{09}
20.	84	R/S	45.			
21.	34	RCL	46.			
22.	00	0	47.			
23.	71	x	48.			
24.	04	4	49.			

Example:

Compute $\int_0^2 x^3 dx$ using Simpson's Rule with $h = .25$.

The following data must be found first:

i	0	1	2	3	4	5	6	7	8
x_i	0	.25	.50	.75	1.00	1.25	1.50	1.75	2.00
$f(x_i)$	0	.0156	.1250	.4219	1.0000	1.9531	3.3750	5.3594	8.0000

Solution:

$$\int_0^2 x^3 dx \cong 4.0000$$

The exact solution is 4.

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Enter h	h	BST	R/S			h/3
3	Enter $f(x_0)$	$f(x_0)$	R/S				Partial Sum
4	Enter $f(x_n)$	$f(x_n)$	R/S				Partial Sum
5	Perform for $i = 1, 2, \dots, n-1$	$f(x_i)$	R/S				Partial Sum
6	Enter $f(x_{n-1})$	$f(x_{n-1})$	R/S				I

NUMERICAL SOLUTION TO DIFFERENTIAL EQUATIONS

This program may be used to solve a wide variety of first order differential equations of the form

$$y' = f(x, y)$$

with initial values x_0, y_0 .

The solution is a numerical solution which calculates y_i for $x_i = x_0 + ih$ ($i = 1, 2, 3, \dots$). h is an increment specified by the user.

The program uses Euler's method:

$$y_{i+1} = y_i + f(x_i, y_i) h$$

$f(x, y)$ is keyed into memory starting at line 20. The user has 30 program steps and all registers except R_4, R_5 , and R_6 available to write $f(x, y)$. The user can assume x to be in the X-register and in R_6 and can assume y to be in the Y-register and in R_5 . The routine should return the value of $f(x, y)$ in the X-register and should end with GTO 08. The accuracy of this method is low unless a small step size is used.

DISPLAY		KEY ENTRY	DISPLAY		KEY ENTRY	REGISTERS
LINE	CODE		LINE	CODE		
00.			25.			R_0
01.	22	$x \leftrightarrow y$	26.			R_1
02.	33	STO	27.			R_2
03.	05	5	28.			R_3
04.	22	$x \leftrightarrow y$	29.			R_4 h
05.	33	STO	30.			R_5 y
06.	06	6	31.			R_6 x
07.	-20	GTO 20	32.			R_7
08.	34	RCL	33.			R_8
09.	04	4	34.			R_9
10.	71	x	35.			R_{10}
11.	34	RCL	36.			R_{11}
12.	05	5	37.			R_{12}
13.	61	+	38.			R_{13}
14.	34	RCL	39.			R_{14}
15.	06	6	40.			R_{15}
16.	34	RCL	41.			R_{16}
17.	04	4	42.			R_{17}
18.	61	+	43.			R_{18}
19.	-00	GTO 00	44.			R_{19}
20.			45.			
21.			46.			
22.			47.			
23.			48.			
24.			49.			

Example:

Solve numerically the differential equation

$$y' = \frac{x + 1 + 2y}{x}$$

 with the initial conditions $x_0 = 1$, $y_0 = -0.5$. Use a step size of $h = 0.1$.

Solutions:

 The keystrokes for $f(x, y)$ are: 1 + x↔y 2 × + RCL 6 ÷

x	1	1.1	1.2	1.3	1.4	1.5
y	-0.5	-0.4	-0.28	-0.15	0.01	0.18

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Switch to RUN mode		GTO	1	9		
3	Switch to PRGM						
4	Key in function						
5	Key in		GTO	0	8		
6	Switch to RUN						
7	Store step size	h	STO	4			
8	Put in initial values	y_0	↑				
		x_0	BST	R/S			x_i
9	If y value desired		$x↔y$				y_i
10	If y value has been displayed		$x↔y$				x_i
11	For next x value		R/S				x_{i+1}
12	Go to step 9						

LINEAR INTERPOLATION

If $(x_1, f(x_1))$ and $(x_2, f(x_2))$ are two points of a function $f(x)$, then the function at x_0 can be approximated by the following formula:

$$f(x_0) \cong \frac{(x_2 - x_0) f(x_1) + (x_0 - x_1) f(x_2)}{(x_2 - x_1)}$$

This is called the linear interpolation formula. Of course, x_2 cannot equal x_1 .

DISPLAY			DISPLAY			REGISTERS
LINE	CODE	KEY ENTRY	LINE	CODE	KEY ENTRY	
00.			25.	-00	GTO 00	R ₀
01.	33	STO	26.			R ₁ x ₁
02.	05	5	27.			R ₂ f(x ₁)
03.	34	RCL	28.			R ₃ x ₂
04.	03	3	29.			R ₄ f(x ₂)
05.	22	x↔y	30.			R ₅ x ₀
06.	51	—	31.			R ₆
07.	34	RCL	32.			R ₇
08.	02	2	33.			R ₈
09.	71	x	34.			R ₉
10.	34	RCL	35.			R ₀₀
11.	05	5	36.			R ₀₁
12.	34	RCL	37.			R ₀₂
13.	01	1	38.			R ₀₃
14.	51	—	39.			R ₀₄
15.	34	RCL	40.			R ₀₅
16.	04	4	41.			R ₀₆
17.	71	x	42.			R ₀₇
18.	61	+	43.			R ₀₈
19.	34	RCL	44.			R ₀₉
20.	03	3	45.			
21.	34	RCL	46.			
22.	01	1	47.			
23.	51	—	48.			
24.	81	÷	49.			

Example:

If (1.2, .30119) and (1.3, .27253) are two points of a function, find $f(1.27)$ and $f(1.29)$.

Solution:

1. $f(1.27) = .28113$ FIX 5

2. $f(1.29) = .27540$ FIX 5

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Store (x_1 , $f(x_1)$)	x_1	STO	1			
		$f(x_1)$	STO	2			
3	Store (x_2 , $f(x_2)$)	x_2	STO	3			
		$f(x_2)$	STO	4			
4	Find $f(x_0)$	x_0	BST	R/S			$f(x_0)$

QUADRATIC EQUATIONS

A general quadratic equation is of the form $ax^2 + bx + c = 0$

The equation has two roots x_1 and x_2 . Let $D = \frac{b^2 - 4ac}{4a^2}$

$$\text{If } D \geq 0 \text{ then } x_1 = \begin{cases} -\frac{b}{2a} + \sqrt{D} & \text{if } -\frac{b}{2a} \geq 0 \\ -\frac{b}{2a} - \sqrt{D} & \text{if } -\frac{b}{2a} < 0 \end{cases}$$

$$\text{and } x_2 = \frac{c}{ax_1}$$

These formulas compute the larger root (in absolute value) first. Better significance can be obtained by this method.

$$\text{If } D < 0 \text{ then } x_1, x_2 = -\frac{b}{2a} \pm \sqrt{-D} i = u \pm iv$$

The coefficient a cannot be zero.

DISPLAY			KEY ENTRY
LINE	CODE		
00.			
01.	34	RCL	
02.	02	2	
03.	34	RCL	
04.	01	1	
05.	02	2	
06.	71	x	
07.	81	÷	
08.	42	CHS	
09.	41	↑	
10.	32	g	
11.	42	x ²	
12.	34	RCL	
13.	03	3	
14.	34	RCL	
15.	01	1	
16.	81	÷	
17.	33	STO	
18.	00	0	
19.	51	—	
20.	84	R/S	
21.	00	0	
22.	31	f	
23.	-32	x ≤ y 32	
24.	23	R↓	
DISPLAY			KEY ENTRY
LINE	CODE		
25.	42	CHS	
26.	31	f	
27.	42	√x	
28.	22	x ↔ y	
29.	84	R/S	
30.	22	x ↔ y	
31.	-00	GTO 00	
32.	23	R↓	
33.	31	f	
34.	42	√x	
35.	22	x ↔ y	
36.	00	0	
37.	31	f	
38.	-43	x ≤ y 43	
39.	23	R↓	
40.	22	x ↔ y	
41.	51	—	
42.	-45	GTO 45	
43.	23	R↓	
44.	61	+	
45.	84	R/S	
46.	34	RCL	
47.	00	0	
48.	22	x ↔ y	
49.	81	÷	

REGISTERS	
R ₀	c/a
R ₁	a
R ₂	b
R ₃	c
R ₄	
R ₅	
R ₆	
R ₇	
R ₈	
R ₉	
R ₁₀	
R ₁₁	
R ₁₂	
R ₁₃	
R ₁₄	
R ₁₅	
R ₁₆	
R ₁₇	
R ₁₈	
R ₁₉	

Examples:

Find the solution to the following three equations:

1. $x^2 - 3x - 4 = 0$

2. $2x^2 + 5x + 3 = 0$

3. $2x^2 + 3x + 4 = 0$

Solutions:

1. $D = 6.25$ $x_1 = 4, x_2 = -1$

2. $D = .06$ $x_1 = -1.50, x_2 = -1.00$

3. $D = -1.44$ $x_1, x_2 = -.75 \pm 1.20i$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Store coefficients	a	STO	1			
		b	STO	2			
		c	STO	3			
3	Calculate D		BST	R/S			D*
4	If $D \geq 0$ roots are real		R/S				x_1 *
			R/S				x_2
	or						
4	If $D < 0$ roots are complex of the form $u \pm i v$		R/S				u *
			R/S				v
	* The stack must be maintained at these positions.						

SYNTHETIC DIVISION

This program divides a polynomial of the form

$$a_0 x^n + a_1 x^{n-1} + a_2 x^{n-2} + \dots + a_n$$

by a binomial of the form

$$x - x_0$$

The positive integer n must be less than or equal to 8. The answer is of the form

$$b_0 x^{n-1} + b_1 x^{n-2} + \dots + b_{n-1} + \frac{b_n}{x - x_0}$$

Be sure to input the x_0 and not the $-x_0$ of $x - x_0$.

DISPLAY			KEY ENTRY	DISPLAY			KEY ENTRY	REGISTERS
LINE	CODE			LINE	CODE			
00.				25.	61	+		R ₀ a ₀
01.	41	↑		26.	84	R/S		R ₁ a ₁
02.	41	↑		27.	71	x		R ₂ a ₂
03.	41	↑		28.	34	RCL		R ₃ a ₃
04.	34	RCL		29.	05	5		R ₄ a ₄
05.	00	0		30.	61	+		R ₅ a ₅
06.	84	R/S		31.	84	R/S		R ₆ a ₆
07.	71	x		32.	71	x		R ₇ a ₇
08.	34	RCL		33.	34	RCL		R ₈ a ₈
09.	01	1		34.	06	6		R ₉
10.	61	+		35.	61	+		R ₁₀
11.	84	R/S		36.	84	R/S		R ₁₁
12.	71	x		37.	71	x		R ₁₂
13.	34	RCL		38.	34	RCL		R ₁₃
14.	02	2		39.	07	7		R ₁₄
15.	61	+		40.	61	+		R ₁₅
16.	84	R/S		41.	84	R/S		R ₁₆
17.	71	x		42.	71	x		R ₁₇
18.	34	RCL		43.	34	RCL		R ₁₈
19.	03	3		44.	08	8		R ₁₉
20.	61	+		45.	61	+		
21.	84	R/S		46.	-00	GTO 00		
22.	71	x		47.				
23.	34	RCL		48.				
24.	04	4		49.				

Examples:

- Find the factors of 823.
- Find the factors of 221.

Solutions:

- 823 is prime. Since $\sqrt{823} = 28.69$ only 2, 3, 5, 7, 11, 13, 17, 19, 23 need be checked.
- 13 and 17

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Let $p = 2$ if x is even, $p = 3$						
	otherwise	x	STO	1			
2	Let $\max = \sqrt{x}$		f	$\sqrt{x}$			max
3			RCL	1			
4		p	$\div$				x/p
5	If x/p is an integer p is a factor	x/p	STO	1			
6	Go to step 4						
7	If x/p is prime then x/p is the only remaining factor. Stop.						
	or						
5	If x/p is not an integer let p be the next prime and go to step 3.						
	Note: Only those p 's less than or equal to $\max$ need be checked.						

POLYNOMIAL EVALUATION

A polynomial of the form

$$f(x) = a_0 x^n + a_1 x^{n-1} + \dots + a_{n-1} x + a_n$$

is evaluated by writing it in the form

$$x (... x (x (a_0 x + a_1) + a_2) + \dots + a_{n-1}) + a_n$$

n can be any positive integer.

DISPLAY			KEY ENTRY	DISPLAY			KEY ENTRY	REGISTERS
LINE	CODE			LINE	CODE			
00.				25.				R ₀
01.	33	STO		26.				R ₁ x
02.	02	2		27.				R ₂ Temporary
03.	34	RCL		28.				R ₃
04.	01	1		29.				R ₄
05.	41	↑		30.				R ₅
06.	41	↑		31.				R ₆
07.	41	↑		32.				R ₇
08.	34	RCL		33.				R ₈
09.	02	2		34.				R ₉
10.	84	R/S		35.				R ₀₀
11.	33	STO		36.				R ₀₁
12.	02	2		37.				R ₀₂
13.	44	CLX		38.				R ₀₃
14.	61	+		39.				R ₀₄
15.	71	x		40.				R ₀₅
16.	34	RCL		41.				R ₀₆
17.	02	2		42.				R ₀₇
18.	61	+		43.				R ₀₈
19.	-10	GTO 10		44.				R ₀₉
20.				45.				
21.				46.				
22.				47.				
23.				48.				
24.				49.				

Example:

Evaluate $f(x) = x^2 + 2x + 3$ for $x = 2$.

Solution:

$$f(2) = 11$$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Store x	x	STO	1			
3	Input a_0	a_0	BST	R/S			
4	Perform for $i = 1, 2, \dots, n-1$	a_i^*	R/S				Partial
5	Input a_n		R/S				$f(x)$
	* The stack must be maintained at this position.						

NUMBER IN BASE b TO A NUMBER IN BASE 10

This program consists of two subprograms. The first changes the integer part of a number in base b to a number in base 10.

$$I_{10} = i_n i_{n-1} \dots i_2 i_1 = i_n b^{n-1} + i_{n-1} b^{n-2} + \dots + i_2 b + i_1$$

This is evaluated in the form

$$b (... (b (b (i_n b + i_{n-1}) + i_{n-2}) + \dots) + i_2) + i_1$$

The second subprogram changes the fraction part of a number in base b to a number in base 10.

$$F_{10} = f_1 f_2 \dots f_m = f_1 b^{-1} + f_2 b^{-2} + \dots + f_m b^{-m}$$

The two programs together can then convert any number in base b to a number in base 10. Zeros must be entered in their proper place.

DISPLAY		KEY ENTRY	DISPLAY		KEY ENTRY	REGISTERS
LINE	CODE		LINE	CODE		
00.			25.	41	↑	R ₀
01.	33	STO	26.	41	↑	R ₁ b
02.	02	2	27.	41	↑	R ₂ Temporary
03.	34	RCL	28.	34	RCL	R ₃ Temporary
04.	01	1	29.	02	2	R ₄
05.	41	↑	30.	71	x	R ₅
06.	41	↑	31.	84	R/S	R ₆
07.	41	↑	32.	33	STO	R ₇
08.	34	RCL	33.	02	2	R ₈
09.	02	2	34.	44	CLX	R ₉
10.	84	R/S	35.	61	+	R ₀₀
11.	33	STO	36.	33	STO	R ₀₁
12.	02	2	37.	03	3	R ₀₂
13.	44	CLX	38.	44	CLX	R ₀₃
14.	61	+	39.	61	+	R ₀₄
15.	71	x	40.	71	x	R ₀₅
16.	34	RCL	41.	41	↑	R ₀₆
17.	02	2	42.	41	↑	R ₀₇
18.	61	+	43.	34	RCL	R ₀₈
19.	-10	GTO 10	44.	02	2	R ₀₉
20.	33	STO	45.	71	x	
21.	02	2	46.	34	RCL	
22.	34	RCL	47.	03	3	
23.	01	1	48.	61	+	
24.	13	1/x	49.	-31	GTO 31	

Examples:

1. Convert 101.0101_2 to a decimal number.
2. Convert $A1_{16}$ to a decimal number (in base 16, A = 10, B = 11, C = 12, D = 13, E = 14, and F = 15).

Solutions:

1. 5.31
2. 161

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Store b	b	STO	1			
3	For integer part	i_n	BST	R/S			
4	Perform for $j = n-1, n-2, \dots, 2$	i_j^*	R/S				Partial
5	Input i_1	i_1	R/S				I_{10}
	or						
3	For fractional part	f_1	GTO	2	0	R/S	
4	Perform for $j = 2, 3, \dots, m-1$	f_j^*	R/S				Partial
5	Input f_m	f_m	R/S				F_{10}
	*The stack must be maintained						
	at these positions.						

NUMBER IN BASE 10 TO NUMBER IN BASE b

This routine converts any number in base 10 to a number in base b , N_b . The user must input the greatest integer of the number displayed after the **R/S**. The greatest integer of a number is the largest integer less than or equal to the number; i.e., if

$$x = 1.5 \text{ then } GI[1.5] = 1$$

and if

$$x = -1.5 \text{ then } GI[-1.5] = -2$$

The user ends the routine when the display flashes or when the accuracy of the machine is exceeded. The variable c must be input as 1 to allow one display position per digit for $2 \leq b \leq 10$ or as 10 to allow 2 display positions for $11 \leq b \leq 100$.

DISPLAY		KEY ENTRY	DISPLAY		KEY ENTRY	REGISTERS
LINE	CODE		LINE	CODE		
00.			25.	61	+	R_0
01.	33	STO	26.	03	3	$R_1 \ b$
02.	02	2	27.	34	RCL	$R_2 \ N_{10}$
03.	00	0	28.	02	2	$R_3 \ N_b$
04.	33	STO	29.	34	RCL	$R_4 \ c$
05.	03	3	30.	01	1	$R_5 \ \text{Temporary}$
06.	23	$R \downarrow$	31.	34	RCL	R_6
07.	31	f	32.	05	5	R_7
08.	22	In	33.	12	y^x	R_8
09.	34	RCL	34.	51	-	R_9
10.	01	1	35.	33	STO	R_{10}
11.	31	f	36.	02	2	R_{11}
12.	22	In	37.	-07	GTO 07	R_{12}
13.	81	$\div$	38.			R_{13}
14.	84	R/S	39.			R_{14}
15.	33	STO	40.			R_{15}
16.	05	5	41.			R_{16}
17.	01	1	42.			R_{17}
18.	00	0	43.			R_{18}
19.	34	RCL	44.			R_{19}
20.	04	4	45.			
21.	71	x	46.			
22.	22	$x \rightarrow y$	47.			
23.	12	y^x	48.			
24.	33	STO	49.			

Examples:

- 1. Convert 18_{10} to base 17.
- 2. Convert 2.333333333 to base 2.

Solutions:

- 1. 101 ; i.e., 11_{17}
- 2. 10.01010101_2

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Store b	b	STO	1			
3	Store c	c	STO	4			
	(c = 1 if $2 \leq b \leq 10$ or c = 10 if $11 \leq b \leq 100$)						
3	Input n	n	BST	R/S			x_1
4	Repeat this step until display flashes or until x_i is more than 10 less than x_1 if c = 1 or more than 5 less than x_1 if c = 10. Input greatest integer less than or equal to x_i	$[x_i]^*$					
			R/S				x_{i+1}
5	To obtain answer		RCL	3			N_b
	* Decide on $[x_i]$ displaying x_i in Fix 9.						

NEWTON'S METHOD SOLUTION TO $f(x)=0$

Newton's method is an iterative solution technique that uses an initial guess x_0 and calculates succeeding x 's by the formula

$$x_{k+1} = x_k - \frac{f(x)}{f'(x)}$$

where $f'(x)$ is the first derivative of $f(x)$.

The user must write in $f(x)/f'(x)$ to be solved starting at line 18. He can assume x is in the X-register and R_1 . He has 31 program memory locations, the stack registers, and all storage registers except R_1 available for this evaluation. The user must end his code with GTO 04.

The code as written gives 5 place accuracy to the right of the decimal. For more or less accuracy change the constant (10^{-12}) in program memory locations 9 through 12. The constant 10^{-12} assures that the square or the change in x is less than 10^{-12} , i.e., that x is off no more than one count in the sixth decimal place, assuring that the fifth decimal place is correct.

Warning:

Newton's method may not converge and will only find one solution. A new solution may be found or convergence may be improved with a different initial guess.

DISPLAY		KEY ENTRY	DISPLAY		KEY ENTRY	REGISTERS
LINE	CODE		LINE	CODE		
00.			25.			R ₀
01.	34	RCL	26.			R ₁ x_k
02.	01	1	27.			R ₂
03.	-18	GTO 18	28.			R ₃
04.	33	STO	29.			R ₄
05.	51	—	30.			R ₅
06.	01	1	31.			R ₆
07.	41	↑	32.			R ₇
08.	71	x	33.			R ₈
09.	43	EEX	34.			R ₉
10.	01	1	35.			R ₀₀
11.	02	2	36.			R ₀₁
12.	42	CHS	37.			R ₀₂
13.	31	f	38.			R ₀₃
14.	-01	$x \leq y$ 01	39.			R ₀₄
15.	34	RCL	40.			R ₀₅
16.	01	1	41.			R ₀₆
17.	-00	GTO 00	42.			R ₀₇
18.			43.			R ₀₈
19.			44.			R ₀₉
20.			45.			
21.			46.			
22.			47.			
23.			48.			
24.			49.			

128 Newton's Method Solution to $f(x)=0$

Example:

1. Find the solution to the function

$$f(x) = x^3 - 2x - 4$$

Solution:

$$\frac{f(x)}{f'(x)} = \frac{x^3 - 2x - 4}{3x^2 - 2}$$

Keystrokes:

$\uparrow$ $\uparrow$ $\uparrow$ $\times$ $\times$ $x \div y$

2 $\times$ - 4 - $x \div y$ $\uparrow$ $\times$

3 $\times$ 2 - $\div$

$$x = 2.00$$

With an initial guess of 10, iteration time is about 30 seconds.

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program	x_0					x
2	Switch to RUN		GTO	1	7		
3	Switch to PRGM						
4	Enter $f(x)/f'(x)$						
5	Key in		GTO	0	4		
6	Switch to RUN						
7	Enter initial guess		STO	1			
8	Find solution		BST	R/S			

TRIGONOMETRIC FUNCTIONS**($\cot$, $\sec$, $\csc$, $\cot^{-1}$, $\sec^{-1}$, $\csc^{-1}$)**

The above functions can be evaluated by the following formulas:

$$1. \cot x = \frac{1}{\tan x}$$

$$2. \sec x = \frac{1}{\cos x}$$

$$3. \csc x = \frac{1}{\sin x}$$

$$4. \cot^{-1} x = \tan^{-1} \left(\frac{1}{x} \right)$$

$$5. \sec^{-1} x = \cos^{-1} \left(\frac{1}{x} \right)$$

$$6. \csc^{-1} x = \sin^{-1} \left(\frac{1}{x} \right)$$

Examples:

1. $\cot(30^\circ) = 1.73$
2. $\sec\left(\frac{\pi}{4}\right) = 1.41$
3. $\csc(100 \text{ grads}) = 1.00$
4. $\cot^{-1}(5) = 11.31^\circ$
5. $\sec^{-1}(2) = 60.00^\circ$
6. $\csc^{-1}(4) = 14.48^\circ$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	For $\cot x$	x	f	tan	1/x		$\cot x$
	or						
	$\sec x$	x	f	cos	1/x		$\sec x$
	or						
	$\csc x$	x	f	sin	1/x		$\csc x$
	or						
	$\cot^{-1} x$	x	1/x	g	$\tan^{-1}$		$\cot^{-1} x$
	or						
	$\sec^{-1} x$	x	1/x	g	$\cos^{-1}$		$\sec^{-1} x$
	or						
	$\csc^{-1} x$	x	1/x	g	$\sin^{-1}$		$\csc^{-1} x$

VERSINE, COVERSINE, HAVERSINE, EXSECANT

The above are calculated by the following formulas:

1. versine (versed sine)

$$\text{vers } \theta = 1 - \cos \theta$$

2. coversine (covered sine)

$$\text{cov } \theta = 1 - \sin \theta$$

3. haversine

$$\text{hav } \theta = \frac{1}{2} \text{vers } \theta = \sin^2 \frac{1}{2} \theta$$

4. exsecant

$$\text{exsec } \theta = \sec \theta - 1$$

The program works in any angular mode. However, if in degrees decimal degrees are assumed.

DISPLAY		KEY ENTRY	DISPLAY		KEY ENTRY	REGISTERS
LINE	CODE		LINE	CODE		
00.			25.	51	—	R ₀
01.	31	f	26.	-00	GTO 00	R ₁
02.	13	cos	27.			R ₂
03.	42	CHS	28.			R ₃
04.	01	1	29.			R ₄
05.	61	+	30.			R ₅
06.	-00	GTO 00	31.			R ₆
07.	31	f	32.			R ₇
08.	12	sin	33.			R ₈
09.	42	CHS	34.			R ₉
10.	01	1	35.			R ₀₀
11.	61	+	36.			R ₀₁
12.	-00	GTO 00	37.			R ₀₂
13.	31	f	38.			R ₀₃
14.	13	cos	39.			R ₀₄
15.	42	CHS	40.			R ₀₅
16.	01	1	41.			R ₀₆
17.	61	+	42.			R ₀₇
18.	02	2	43.			R ₀₈
19.	81	÷	44.			R ₀₉
20.	-00	GTO 00	45.			
21.	31	f	46.			
22.	13	cos	47.			
23.	13	1/x	48.			
24.	01	1	49.			

Examples:

$$\text{vers } 100^\circ = 1.1736$$

$$\text{cov } 100^\circ = .0152$$

$$\text{hav } 100^\circ = .5868$$

$$\text{exsec } 100^\circ = -6.7588$$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Versine	θ	BST	R/S			Vers θ
	or						
	Coversine	θ	GTO	0	7	R/S	Cov θ
	or						
	Haversine	θ	GTO	1	3	R/S	Hav θ
	or						
	Exsecant	θ	GTO	2	1	R/S	Exsec θ

HYPERBOLIC FUNCTIONS

This program evaluates the six hyperbolic functions by the following formulas:

$$1. \sinh x = \frac{e^x - e^{-x}}{2}$$

$$2. \cosh x = \frac{e^x + e^{-x}}{2}$$

$$3. \tanh x = \frac{e^x - e^{-x}}{e^x + e^{-x}}$$

$$4. \operatorname{csch} x = \frac{1}{\sinh x} \quad (x \neq 0)$$

$$5. \operatorname{sech} x = \frac{1}{\cosh x}$$

$$6. \operatorname{coth} x = \frac{1}{\tanh x} \quad (x \neq 0)$$

DISPLAY			KEY ENTRY	DISPLAY		KEY ENTRY	REGISTERS
LINE	CODE			LINE	CODE		
00.				25.	71	x	R ₀
01.	32	g		26.	61	+	R ₁
02.	22	e ^x		27.	81	÷	R ₂
03.	41	↑		28.	-00	GTO 00	R ₃
04.	13	1/x		29.			R ₄
05.	51	-		30.			R ₅
06.	02	2		31.			R ₆
07.	81	÷		32.			R ₇
08.	-00	GTO 00		33.			R ₈
09.	32	g		34.			R ₉
10.	22	e ^x		35.			R ₀₀
11.	41	↑		36.			R ₀₁
12.	13	1/x		37.			R ₀₂
13.	61	+		38.			R ₀₃
14.	-06	GTO 06		39.			R ₀₄
15.	32	g		40.			R ₀₅
16.	22	e ^x		41.			R ₀₆
17.	41	↑		42.			R ₀₇
18.	13	1/x		43.			R ₀₈
19.	51	-		44.			R ₀₉
20.	41	↑		45.			
21.	41	↑		46.			
22.	31	f		47.			
23.	34	LAST X		48.			
24.	02	2		49.			

Examples:

1. $\sinh 1.5 = 2.13$
2. $\cosh 5.9 = 182.52$
3. $\tanh 1.3 = .86$
4. $\operatorname{csch} 0.95 = .91$
5. $\operatorname{sech} (-3) = .10$
6. $\operatorname{coth} (-1.99) = -1.04$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
	$\sinh x$	x	BST	R/S			$\sinh x$
	or						
	$\cosh x$	x	GTO	0	9	R/S	$\cosh x$
	or						
	$\tanh x$	x	GTO	1	5	R/S	$\tanh x$
	or						
	$\operatorname{csch} x$	x	BST	R/S	1/x		$\operatorname{csch} x$
	or						
	$\operatorname{sech} x$	x	GTO	0	9	R/S	
			1/x				$\operatorname{sech} x$
	or						
	$\operatorname{coth} x$	x	GTO	1	5	R/S	$\operatorname{coth} x$
			1/x				

INVERSE HYPERBOLIC FUNCTIONS

This program evaluates the inverse hyperbolic functions by the following formulas:

$$1. \sinh^{-1} x = \ln [x + (x^2 + 1)^{1/2}]$$

$$2. \cosh^{-1} x = \ln [x + (x^2 - 1)^{1/2}] \quad x \geq 1$$

$$3. \tanh^{-1} x = \frac{1}{2} \ln \left[\frac{1+x}{1-x} \right] \quad x^2 < 1$$

$$4. \operatorname{csch}^{-1} x = \sinh^{-1} \left[\frac{1}{x} \right] \quad x \neq 0$$

$$5. \operatorname{sech}^{-1} x = \cosh^{-1} \left[\frac{1}{x} \right] \quad 0 < x \leq 1$$

$$6. \operatorname{coth}^{-1} x = \tanh^{-1} \left[\frac{1}{x} \right] \quad x^2 > 1$$

DISPLAY		KEY ENTRY	DISPLAY		KEY ENTRY	REGISTERS
LINE	CODE		LINE	CODE		
00.			25.	01	1	R ₀
01.	41	↑	26.	61	+	R ₁
02.	41	↑	27.	22	$x \leftrightarrow y$	R ₂
03.	71	x	28.	42	CHS	R ₃
04.	01	1	29.	01	1	R ₄
05.	61	+	30.	61	+	R ₅
06.	31	f	31.	81	÷	R ₆
07.	42	$\sqrt{x}$	32.	31	f	R ₇
08.	61	+	33.	22	ln	R ₈
09.	31	f	34.	02	2	R ₉
10.	22	ln	35.	81	÷	R ₀₀
11.	-00	GTO 00	36.	-00	GTO 00	R ₀₁
12.	41	↑	37.			R ₀₂
13.	41	↑	38.			R ₀₃
14.	71	x	39.			R ₀₄
15.	01	1	40.			R ₀₅
16.	51	-	41.			R ₀₆
17.	31	f	42.			R ₀₇
18.	42	$\sqrt{x}$	43.			R ₀₈
19.	61	+	44.			R ₀₉
20.	31	f	45.			
21.	22	ln	46.			
22.	-00	GTO 00	47.			
23.	41	↑	48.			
24.	41	↑	49.			

Examples:

1. $\sinh^{-1} (3.5) = 1.97$
2. $\cosh^{-1} (100) = 5.30$
3. $\tanh^{-1} (-.7) = -.87$
4. $\operatorname{csch}^{-1} (3) = .33$
5. $\operatorname{sech}^{-1} (.5) = 1.32$
6. $\operatorname{coth}^{-1} (5.4) = .19$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
	$\sinh^{-1} x$	x	BST	R/S			$\sinh^{-1} x$
	or						
	$\cosh^{-1} x$	x	GTO	1	2	R/S	$\cosh^{-1} x$
	or						
	$\tanh^{-1} x$	x	GTO	2	3	R/S	$\tanh^{-1} x$
	or						
	$\operatorname{csch}^{-1} x$	x	1/x	BST	R/S		$\operatorname{csch}^{-1} x$
	or						
	$\operatorname{sech}^{-1} x$	x	1/x	GTO	1	2	$\operatorname{sech}^{-1} x$
			R/S				
	or						
	$\operatorname{coth}^{-1} x$	x	1/x	GTO	2	3	$\operatorname{coth}^{-1} x$
			R/S				

POLYGONS INSCRIBED IN A CIRCLE

Given the radius of a circle this program calculates the length S_1 of a side and the area A_1 of a polygon of n sides inscribed in the circle. The formulas used are:

$$1. S_1 = 2r \sin \left(\frac{c}{n} \right)$$

$$2. A_1 = \frac{1}{2} n r^2 \sin \left(\frac{2c}{n} \right)$$

where

$$c = 2 \sin^{-1} 1 = \pi \text{ radians} = 180^\circ = 200 \text{ grads}$$

n = number of sides

r = radius of the circle

It must be true that n is an integer greater than 2.

DISPLAY			KEY ENTRY			REGISTERS
LINE	CODE	KEY ENTRY	LINE	CODE	KEY ENTRY	
00.			25.	34	RCL	R ₀
01.	01	1	26.	02	2	R ₁ n
02.	32	g	27.	41	↑	R ₂ r
03.	12	$\sin^{-1}$	28.	71	x	R ₃ c/n
04.	02	2	29.	71	x	R ₄
05.	71	x	30.	34	RCL	R ₅
06.	34	RCL	31.	01	1	R ₆
07.	01	1	32.	71	x	R ₇
08.	81	÷	33.	02	2	R ₈
09.	33	STO	34.	81	÷	R ₉
10.	03	3	35.	-00	GTO 00	R ₀₀
11.	31	f	36.			R ₀₁
12.	12	sin	37.			R ₀₂
13.	02	2	38.			R ₀₃
14.	71	x	39.			R ₀₄
15.	34	RCL	40.			R ₀₅
16.	02	2	41.			R ₀₆
17.	71	x	42.			R ₀₇
18.	84	R/S	43.			R ₀₈
19.	34	RCL	44.			R ₀₉
20.	03	3	45.			
21.	02	2	46.			
22.	71	x	47.			
23.	31	f	48.			
24.	12	sin	49.			

Example:

Given a circle of radius 5 find the length of a side and area of a polygon of 6 sides inscribed in the circle.

Solution:

$$S_1 = 5.00$$

$$A_1 = 64.95$$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program	n r					S ₁ A ₁
2	Store n and r		STO	1			
			STO	2			
3	Find S ₁		BST	R/S			
4	Find A ₁		R/S				

POLYGONS CIRCUMSCRIBED ABOUT A CIRCLE

Given the radius of a circle this program calculates the length S_2 of a side and the area A_2 of a polygon of n sides circumscribed about a circle. The formulas used are:

$$1. S_2 = 2r \tan (c/n)$$

$$2. A_2 = n r^2 \tan (c/n)$$

where

$$c = 2 \sin^{-1} 1 = \pi \text{ radians} = 180^\circ = 200 \text{ grads}$$

n = number of sides

r = radius of the circle

It must be true that n is an integer greater than 2.

DISPLAY		KEY ENTRY	DISPLAY		KEY ENTRY	REGISTERS
LINE	CODE		LINE	CODE		
00.			25.	01	1	R ₀
01.	01	1	26.	71	x	R ₁ n
02.	32	g	27.	-00	GTO 00	R ₂ r
03.	12	$\sin^{-1}$	28.			R ₃ r tan (c/n)
04.	02	2	29.			R ₄
05.	71	x	30.			R ₅
06.	34	RCL	31.			R ₆
07.	01	1	32.			R ₇
08.	81	$\div$	33.			R ₈
09.	31	f	34.			R ₉
10.	14	tan	35.			R ₀₀
11.	34	RCL	36.			R ₀₁
12.	02	2	37.			R ₀₂
13.	71	x	38.			R ₀₃
14.	33	STO	39.			R ₀₄
15.	03	3	40.			R ₀₅
16.	02	2	41.			R ₀₆
17.	71	x	42.			R ₀₇
18.	84	R/S	43.			R ₀₈
19.	34	RCL	44.			R ₀₉
20.	03	3	45.			
21.	34	RCL	46.			
22.	02	2	47.			
23.	71	x	48.			
24.	34	RCL	49.			

Example:

Find the length of a side and the area of a polygon of 6 sides circumscribed about a circle of radius 5.

Solution:

$$S_2 = 5.77$$

$$A_2 = 86.60$$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Store n and r	n	STO	1			
		r	STO	2			
3	Calculate S_2		BST	R/S			S_2
4	Calculate A_2		R/S				A_2

CIRCLE DETERMINED BY THREE POINTS

Let (x_1, y_1) (x_2, y_2) (x_3, y_3) be three points such that $x_1 \neq x_2$ and $x_1 \neq x_3$. If the points cannot be renumbered to satisfy this condition, the points cannot be on a circle. Let the center of the circle be (x_0, y_0) and the radius of the circle be r . Then

$$y_0 = \frac{k_2 - k_1}{n_2 - n_1}, \quad x_0 = k_2 - n_2 y_0, \quad \text{and } r = \sqrt{(x_1 - x_0)^2 + (y_1 - y_0)^2}$$

where

$$k_1 = \frac{1}{2} [(x_1 + x_2) + n_1 (y_1 + y_2)], \quad k_2 = \frac{1}{2} [(x_1 + x_3) + n_2 (y_1 + y_3)]$$

$$n_1 = \frac{y_1 - y_2}{x_1 - x_2}, \quad \text{and } n_2 = \frac{y_1 - y_3}{x_1 - x_3}$$

If $n_1 = n_2$ the points cannot form a circle.

DISPLAY		KEY ENTRY	DISPLAY		KEY ENTRY	REGISTERS
LINE	CODE		LINE	CODE		
00.			25.	02	2	R ₀
01.	34	RCL	26.	81	÷	R ₁ x ₁
02.	02	2	27.	-00	GTO 00	R ₂ y ₁
03.	34	RCL	28.	34	RCL	R ₃ x ₂ , x ₃
04.	04	4	29.	08	8	R ₄ y ₂ , y ₃
05.	51	—	30.	34	RCL	R ₅ n ₁
06.	34	RCL	31.	06	6	R ₆ k ₁
07.	01	1	32.	51	—	R ₇ n ₂
08.	34	RCL	33.	34	RCL	R ₈ k ₂
09.	03	3	34.	07	7	R ₉ y ₀
10.	51	—	35.	34	RCL	R ₀₀
11.	81	÷	36.	05	5	R ₀₁
12.	84	R/S	37.	51	—	R ₀₂
13.	34	RCL	38.	81	÷	R ₀₃
14.	02	2	39.	33	STO	R ₀₄
15.	34	RCL	40.	09	9	R ₀₅
16.	04	4	41.	84	R/S	R ₀₆
17.	61	+	42.	34	RCL	R ₀₇
18.	71	x	43.	07	7	R ₀₈
19.	34	RCL	44.	71	x	R ₀₉
20.	01	1	45.	34	RCL	
21.	61	+	46.	08	8	
22.	34	RCL	47.	22	x↔y	
23.	03	3	48.	51	—	
24.	61	+	49.	-00	GTO 00	

Examples:

- Find the equation of the circle that goes through the three points (1, 1), (3.5, -7.6), (12, 0.8).
- Find the equation of the circle that passes through the three points (0, 1), (-1, 0), (0, -1).

Solutions:

- $n_1 = -3.44$, $k_1 = 13.60$, $n_2 = -.02$, $k_2 = 6.48$
Center = (6.45, -2.08), $r = 6.26$
Equation: $(x - 6.45)^2 + (y + 2.08)^2 = (6.26)^2$
- $n_1 = 1.00$, $k_1 = 0.00$, $n_2 = -1.00$, $k_2 = 0.00$
Center = (0, 0), $r = 1$
Equation: $x^2 + y^2 = 1$
Note: (-1, 0) must be (x_1, y_1)

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Store (x_1, y_1)	x_1	STO	1			
		y_1	STO	2			
3	Store (x_2, y_2)	x_2	STO	3			
		y_2	STO	4			
4	Calculate and store n_1		BST	R/S			n_1^*
			STO	5			
5	Calculate and store k_1		R/S				k_1
			STO	6			
6	Store (x_3, y_3)	x_3	STO	3			
		y_3	STO	4			
7	Calculate and store n_2		R/S				n_2^*
			STO	7			
8	Calculate and store k_2		R/S				k_2
			STO	8			
9	Find y_0		GTO	2	8	R/S	y_0^*
10	Find x_0		R/S				x_0^*
11	Calculate r		RCL	1	-	RCL	
			9	RCL	2	-	
			g	R→P			r
	* The stack should be maintained at these points.						

EQUALLY SPACED POINTS ON A CIRCLE

Given a circle with center (x_0, y_0) and radius r , this program calculates the coordinates of equally spaced points on the circle. The user inputs the coordinates of the center, the radius, the number of points n to be spaced on the circle, and an angle θ (measured from the positive x -axis) which describes the position of the first point on the circle.

The formulas used are:

$$x_{k+1} = x_0 + r \cos(\theta + ck)$$

$$y_{k+1} = y_0 + r \sin(\theta + ck)$$

where

$$k = 0, 1, 2, \dots, n - 1$$

and

$$c = \frac{4 \sin^{-1} 1}{n} = \frac{2\pi \text{ radians}}{n} = \frac{360^\circ}{n} = \frac{400 \text{ grads}}{n}$$

The program works in any angular mode but if in degrees decimal degrees are assumed.

DISPLAY		KEY ENTRY	DISPLAY		KEY ENTRY	REGISTERS
LINE	CODE		LINE	CODE		
00.			25.	61	+	R_0
01.	13	$1/x$	26.	34	RCL	$R_1 \times_0$
02.	01	1	27.	04	4	$R_2 y_0$
03.	32	g	28.	31	f	$R_3 \theta$
04.	12	$\sin^{-1}$	29.	00	R→P	$R_4 r$
05.	04	4	30.	34	RCL	$R_5 c$
06.	71	x	31.	01	1	$R_6 k$
07.	71	x	32.	61	+	$R_7 \theta + ck$
08.	33	STO	33.	84	R/S	R_8
09.	05	5	34.	22	$x \div y$	R_9
10.	01	1	35.	34	RCL	R_{10}
11.	42	CHS	36.	02	2	R_{11}
12.	33	STO	37.	61	+	R_{12}
13.	06	6	38.	84	R/S	R_{13}
14.	01	1	39.	-14	GTO 14	R_{14}
15.	33	STO	40.			R_{15}
16.	61	+	41.			R_{16}
17.	06	6	42.			R_{17}
18.	34	RCL	43.			R_{18}
19.	03	3	44.			R_{19}
20.	34	RCL	45.			
21.	05	5	46.			
22.	34	RCL	47.			
23.	06	6	48.			
24.	71	x	49.			

146 Equally Spaced Points on a Circle


Examples:

- Find five points equally spaced around a circle with center at (4.28, 3.10) with radius 1. Start the first point at $\pi/4$ radians around the circle.
- Find three points equally spaced around a circle with center at (-3.4, 1.8) with radius 3.21. Start the first point 36° around the circle.

Solutions:

- (4.99, 3.81), (3.83, 3.99), (3.29, 2.94), (4.12, 2.11), (5.17, 2.65)
(Set calculator in radian mode.)
- (-.80, 3.69), (-6.33, 3.11), (-3.06, -1.39)

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Store (x_0, y_0)	x_0	STO	1			
		y_0	STO	2			
3	Store θ	θ	STO	3			
4	Store r	r	STO	4			
5	Enter n	n	BST	R/S			x_1^*
			R/S				y_1
6	Perform for $i = 2, \dots, n$		R/S				x_i^*
			R/S				y_i
	* Stack must be maintained at these positions.						

TRIANGLE SOLUTION B, b, c

Given two sides and a non-included angle, this program solves the triangle for the remaining parameters by the following formulas:

$$1. C = \sin^{-1} \left(\frac{c \sin B}{b} \right)$$

$$2. A = 2 \sin^{-1} 1 - (B + C) = \pi \text{ radians} - (B + C) = 180^\circ - (B + C) \\ = 200 \text{ grads} - (B + C)$$

$$3. a = \frac{b \sin A}{\sin B}$$

If B is acute ($< 90^\circ$) and $b < c$, a second set of solutions exists and is calculated by the following formulas:

$$4. C' = 2 \sin^{-1} 1 - C$$

$$5. A' = 2 \sin^{-1} 1 - (B + C')$$

148 Triangle Solution B, b, c

DISPLAY			KEY ENTRY	DISPLAY			KEY ENTRY	REGISTERS
LINE	CODE			LINE	CODE			
00.				25.	04	4		R ₀
01.	34	RCL		26.	22	$x \div y$		R ₁ B
02.	03	3		27.	51	—		R ₂ b
03.	34	RCL		28.	84	R/S		R ₃ c
04.	01	1		29.	31	f		R ₄ $2 \sin^{-1} 1$
05.	31	f		30.	12	sin		R ₅ C
06.	12	sin		31.	34	RCL		R ₆
07.	71	x		32.	02	2		R ₇
08.	34	RCL		33.	71	x		R ₈
09.	02	2		34.	34	RCL		R ₉
10.	81	$\div$		35.	01	1		R ₀₀
11.	32	g		36.	31	f		R ₀₁
12.	12	$\sin^{-1}$		37.	12	sin		R ₀₂
13.	33	STO		38.	81	$\div$		R ₀₃
14.	05	5		39.	84	R/S		R ₀₄
15.	84	R/S		40.	34	RCL		R ₀₅
16.	34	RCL		41.	04	4		R ₀₆
17.	01	1		42.	34	RCL		R ₀₇
18.	61	+		43.	05	5		R ₀₈
19.	01	1		44.	51	—		R ₀₉
20.	32	g		45.	84	R/S		
21.	12	$\sin^{-1}$		46.	-16	GTO 16		
22.	02	2		47.				
23.	71	x		48.				
24.	33	STO		49.				

Example:

Given the following two sides and non-included angle:

$$B = 33^\circ 40' \text{ (convert to decimal degrees)}$$

$$b = 31.5$$

$$c = 51.8$$

Solve the triangle.

Solution:

Since B is less than 90° and $b < c$, two sets of solutions exist.

$$C = 65.73$$

$$C' = 114.27^\circ$$


$$A = 80.60$$

$$A' = 32.06^\circ$$

$$a = 56.06$$

$$a' = 30.16$$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Store B, b, and c	B	STO	1			
		b	STO	2			
		c	STO	3			
3	Solve triangle		BST	R/S			C*
			R/S				A*
			R/S				a*
4	If $B < 90^\circ$ and $b < c$ find alternate solution						
			R/S				C'
			R/S				A'
			R/S				a'
	* The stack must be maintained at these positions.						

TRIANGLE SOLUTION a, b, c

Given three sides of a triangle this program solves the triangle for the remaining parameters by the following formulas:

$$C = \cos^{-1} \left(\frac{a^2 + b^2 - c^2}{2ab} \right)$$

$$B = \sin^{-1} \left(\frac{b \sin C}{c} \right)$$

$$A = \sin^{-1} \left(\frac{a \sin C}{c} \right)$$

Reletter if necessary to make c the largest side. The program works in any angular mode. However, if in degree mode decimal degrees are assumed.

DISPLAY		KEY ENTRY	DISPLAY		KEY ENTRY	REGISTERS
LINE	CODE		LINE	CODE		
00.			25.	84	R/S	R ₀
01.	34	RCL	26.	31	f	R ₁ a
02.	01	1	27.	12	sin	R ₂ b
03.	41	↑	28.	34	RCL	R ₃ c
04.	71	x	29.	03	3	R ₄
05.	34	RCL	30.	81	÷	R ₅
06.	02	2	31.	34	RCL	R ₆
07.	41	↑	32.	02	2	R ₇
08.	71	x	33.	22	x↗y	R ₈
09.	61	+	34.	71	x	R ₉
10.	34	RCL	35.	31	f	R ₀₀
11.	03	3	36.	34	LAST X	R ₀₁
12.	41	↑	37.	22	x↗y	R ₀₂
13.	71	x	38.	32	g	R ₀₃
14.	51	—	39.	12	sin ⁻¹	R ₀₄
15.	34	RCL	40.	84	R/S	R ₀₅
16.	01	1	41.	23	R↓	R ₀₆
17.	34	RCL	42.	34	RCL	R ₀₇
18.	02	2	43.	01	1	R ₀₈
19.	71	x	44.	71	x	R ₀₉
20.	02	2	45.	32	g	
21.	71	x	46.	12	sin ⁻¹	
22.	81	÷	47.	-00	GTO 00	
23.	32	g	48.			
24.	13	cos ⁻¹	49.			

Example:

Given the following three sides:

a = 30.3

b = 40.4

c = 62.6

Solve the triangle.


Solution:

C = 123.99°

B = 32.35°

A = 23.66°

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Store a, b, and c (c is the largest)	a	STO	1			
		b	STO	2			
		c	STO	3			
3	Find the solution		BST	R/S			C*
			R/S				B*
			R/S				A
	* The stack must be maintained at these positions.						

TRIANGLE SOLUTION a, A, C

Given two angles and an opposite side this program solves the triangle for the remaining parameters by the following formulas:

$$B = 2 \sin^{-1} 1 - (A + C) = \pi \text{ radians} - (A + C) = 180^\circ - (A + C) \\ = 200 \text{ grads} - (A + C)$$

$$b = \frac{a \sin B}{\sin A}$$

$$c = \frac{a \sin C}{\sin A}$$

The program works in any angular mode. However, if in degree mode all angles are assumed to be in decimal degrees.

DISPLAY			KEY ENTRY	DISPLAY			KEY ENTRY	REGISTERS
LINE	CODE			LINE	CODE			
00.				25.	01	1		R ₀
01.	01	1		26.	31	f		R ₁ a
02.	32	g		27.	34	LAST X		R ₂ A
03.	12	sin ⁻¹		28.	81	÷		R ₃ C
04.	02	2		29.	34	RCL		R ₄
05.	71	x		30.	03	3		R ₅
06.	34	RCL		31.	31	f		R ₆
07.	02	2		32.	12	sin		R ₇
08.	34	RCL		33.	71	x		R ₈
09.	03	3		34.	-00	GTO 00		R ₉
10.	61	+		35.				R ₁₀
11.	51	-		36.				R ₁₁
12.	84	R/S		37.				R ₁₂
13.	31	f		38.				R ₁₃
14.	12	sin		39.				R ₁₄
15.	34	RCL		40.				R ₁₅
16.	01	1		41.				R ₁₆
17.	71	x		42.				R ₁₇
18.	34	RCL		43.				R ₁₈
19.	02	2		44.				R ₁₉
20.	31	f		45.				
21.	12	sin		46.				
22.	81	÷		47.				
23.	84	R/S		48.				
24.	34	RCL		49.				

Example:

Given the following two angles and opposite side:

$$a = 17.5$$

$$A = 41.23^\circ$$

$$C = 62.20^\circ$$

Solve the triangle.

Solution:


$$B = 76.57^\circ$$

$$b = 25.83$$

$$c = 23.49$$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Store a, A, and C	a	STO	1			
		A	STO	2			
		C	STO	3			
3	Find the solution		BST	R/S			B*
			R/S				b*
			R/S				c
	* The stack must be maintained at these positions.						

TRIANGLE SOLUTION a, b, C



Given two sides and their included angle this program solves the triangle for the remaining parameters by the following formulas:

$$c = \sqrt{a^2 + b^2 - 2ab \cos C} \qquad A = \sin^{-1} \left(\frac{a \sin C}{c} \right)$$

$$B = 2 \sin^{-1} 1 - (A + C) = \pi \text{ radians} - (A + C) = 180^\circ - (A + C) \\ = 200 \text{ grads} - (A + C)$$

Reletter if necessary, to make a the smaller of a and b.

This program works in any angular mode. However, if in degrees decimal degrees are assumed.

DISPLAY			KEY ENTRY	DISPLAY			KEY ENTRY	REGISTERS
LINE	CODE			LINE	CODE			
00.				25.	34	RCL		R ₀
01.	34	RCL		26.	01	1		R ₁ a
02.	01	1		27.	34	RCL		R ₂ b
03.	34	RCL		28.	03	3		R ₃ C
04.	02	2		29.	31	f		R ₄
05.	32	g		30.	12	sin		R ₅
06.	00	R→P		31.	71	x		R ₆
07.	32	g		32.	22	x↔y		R ₇
08.	42	x ²		33.	81	÷		R ₈
09.	34	RCL		34.	32	g		R ₉
10.	01	1		35.	12	sin ⁻¹		R ₀₀
11.	34	RCL		36.	84	R/S		R ₀₁
12.	02	2		37.	01	1		R ₀₂
13.	71	x		38.	32	g		R ₀₃
14.	02	2		39.	12	sin ⁻¹		R ₀₄
15.	71	x		40.	02	2		R ₀₅
16.	34	RCL		41.	71	x		R ₀₆
17.	03	3		42.	22	x↔y		R ₀₇
18.	31	f		43.	34	RCL		R ₀₈
19.	13	cos		44.	03	3		R ₀₉
20.	71	x		45.	61	+		
21.	51	—		46.	51	—		
22.	31	f		47.	-00	GTO 00		
23.	42	√x		48.				
24.	84	R/S		49.				

Examples:

Given the following two sides and included angle:

$$a = 132$$

$$b = 224$$

$$C = 28^{\circ}40' \text{ (convert to decimal degrees first)}$$

Solve the triangle.

Solution:


$$c = 125.35$$

$$A = 30.34^{\circ}$$

$$B = 120.99^{\circ}$$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Store a, b, and C (a is smaller of a and b)						
		a	STO	1			
		b	STO	2			
		C	STO	3			
3	Solve triangle		BST	R/S			c*
			R/S				A*
			R/S				B
	* The stack must be maintained at these positions.						

TRIANGLE SOLUTION a, B, C



Given two angles and their included side this program solves the triangle for the remaining parameters by the following formulas:

$$A = 2 \sin^{-1} 1 - (B + C) = \pi \text{ radians} - (B + C) = 180^\circ - (B + C)$$

$$= 200 \text{ grads} - (B + C)$$

$$b = \frac{a \sin B}{\sin A}$$

$$c = \frac{a \sin C}{\sin A}$$

The program works in any angular mode. However, if in degrees the program assumes decimal degrees.

DISPLAY		KEY ENTRY	DISPLAY		KEY ENTRY	REGISTERS
LINE	CODE		LINE	CODE		
00.			25.	02	2	R ₀
01.	01	1	26.	31	f	R ₁ a
02.	32	g	27.	12	sin	R ₂ B
03.	12	$\sin^{-1}$	28.	71	x	R ₃ C
04.	02	2	29.	84	R/S	R ₄ A, (a/sin A)
05.	71	x	30.	34	RCL	R ₅
06.	34	RCL	31.	04	4	R ₆
07.	02	2	32.	34	RCL	R ₇
08.	34	RCL	33.	03	3	R ₈
09.	03	3	34.	31	f	R ₉
10.	61	+	35.	12	sin	R ₀₀
11.	51	—	36.	71	x	R ₀₁
12.	33	STO	37.	-00	GTO 00	R ₀₂
13.	04	4	38.			R ₀₃
14.	84	R/S	39.			R ₀₄
15.	34	RCL	40.			R ₀₅
16.	01	1	41.			R ₀₆
17.	34	RCL	42.			R ₀₇
18.	04	4	43.			R ₀₈
19.	31	f	44.			R ₀₉
20.	12	sin	45.			
21.	81	÷	46.			
22.	33	STO	47.			
23.	04	4	48.			
24.	34	RCL	49.			

Example:

Given the following two angles and their included side:

$a = 25.2$

$B = 35^{\circ}20'$ (convert B and C to decimal degrees first)

$C = 68^{\circ}30'$

Solve the triangle.

Solution:

$A = 76.17^{\circ}$

$b = 15.01$

$c = 24.15$


STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Store a, B, C	a	STO	1			
		B	STO	2			
		C	STO	3			
3	Solve triangle		BST	R/S			A*
			R/S				b*
			R/S				c
	* The stack must be maintained at these positions.						

AREA OF A TRIANGLE a, b, c

Given three sides of a triangle this program computes the area by the following formula:

$$\text{Area} = \sqrt{s(s-a)(s-b)(s-c)}$$

where $s = \frac{1}{2}(a + b + c)$



DISPLAY			KEY ENTRY	DISPLAY			KEY ENTRY	REGISTERS
LINE	CODE			LINE	CODE			
00.				25.	03	3		R ₀
01.	34	RCL		26.	51	—		R ₁ a
02.	01	1		27.	71	x		R ₂ b
03.	34	RCL		28.	31	f		R ₃ c
04.	02	2		29.	42	$\sqrt{x}$		R ₄
05.	61	+		30.	-00	GTO 00		R ₅
06.	34	RCL		31.				R ₆
07.	03	3		32.				R ₇
08.	61	+		33.				R ₈
09.	02	2		34.				R ₉
10.	81	÷		35.				R ₀₀
11.	41	↑		36.				R ₀₁
12.	41	↑		37.				R ₀₂
13.	41	↑		38.				R ₀₃
14.	34	RCL		39.				R ₀₄
15.	01	1		40.				R ₀₅
16.	51	—		41.				R ₀₆
17.	71	x		42.				R ₀₇
18.	22	$x \rightarrow y$		43.				R ₀₈
19.	34	RCL		44.				R ₀₉
20.	02	2		45.				
21.	51	—		46.				
22.	71	x		47.				
23.	22	$x \rightarrow y$		48.				
24.	34	RCL		49.				

Example:

Find the area of a triangle with the following three sides:

$$a = 5.31$$

$$b = 7.09$$


$$c = 8.86$$

Solution:

$$\text{Area} = 18.82$$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Store a, b, c	a	STO	1			
		b	STO	2			
		c	STO	3			
3	Find area		BST	R/S			Area

AREA OF A TRIANGLE a, b, C



Given two sides and an included angle of a triangle this program computes the area by the following formula:

Area = 1/2 ab sin C

The angle C can be in any angular mode but if in degrees it is assumed to be in decimal degrees.

DISPLAY		KEY ENTRY	DISPLAY		KEY ENTRY	REGISTERS
LINE	CODE		LINE	CODE		
00.			25.			R ₀
01.	34	RCL	26.			R ₁ a
02.	03	3	27.			R ₂ b
03.	31	f	28.			R ₃ C
04.	12	sin	29.			R ₄
05.	34	RCL	30.			R ₅
06.	01	1	31.			R ₆
07.	71	x	32.			R ₇
08.	34	RCL	33.			R ₈
09.	02	2	34.			R ₉
10.	71	x	35.			R ₀₀
11.	02	2	36.			R ₀₁
12.	81	÷	37.			R ₀₂
13.	-00	GTO 00	38.			R ₀₃
14.			39.			R ₀₄
15.			40.			R ₀₅
16.			41.			R ₀₆
17.			42.			R ₀₇
18.			43.			R ₀₈
19.			44.			R ₀₉
20.			45.			
21.			46.			
22.			47.			
23.			48.			
24.			49.			

Example:

Find the area of the triangle with the following two sides and included angle.

$$a = 5.3174$$

$$b = 7.0898$$


$$C = 45^\circ$$

Solution:

$$\text{Area} = 13.33$$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Store data	a	STO	1			
		b	STO	2			
		C	STO	3			
3	Find area		BST	R/S			Area

AREA OF A TRIANGLE a, B, C



Given two angles and an included side of a triangle this program computes the area by the following formula:

$$\text{Area} = \frac{a^2 \sin B \sin C}{2 \sin (B + C)}$$

Angles B and C can be in any angular mode. If in degrees all angles are assumed to be decimal degrees.

DISPLAY			KEY ENTRY	DISPLAY			KEY ENTRY	REGISTERS
LINE	CODE			LINE	CODE			
00.				25.	-00	GTO 00		R ₀
01.	34	RCL		26.				R ₁ a
02.	01	1		27.				R ₂ B
03.	32	g		28.				R ₃ C
04.	42	x ²		29.				R ₄
05.	02	2		30.				R ₅
06.	81	÷		31.				R ₆
07.	34	RCL		32.				R ₇
08.	02	2		33.				R ₈
09.	31	f		34.				R ₉
10.	12	sin		35.				R ₀₀
11.	71	x		36.				R ₀₁
12.	34	RCL		37.				R ₀₂
13.	03	3		38.				R ₀₃
14.	31	f		39.				R ₀₄
15.	12	sin		40.				R ₀₅
16.	71	x		41.				R ₀₆
17.	34	RCL		42.				R ₀₇
18.	02	2		43.				R ₀₈
19.	34	RCL		44.				R ₀₉
20.	03	3		45.				
21.	61	+		46.				
22.	31	f		47.				
23.	12	sin		48.				
24.	81	÷		49.				

Example:

Given the following two angles and included side find the area of the triangle.

$$a = 14.625$$

$$B = 70.54^\circ$$

$$C = 62.96^\circ$$

Solution:

$$\text{Area} = 123.82$$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Input data	a	STO	1			
		B	STO	2			
		C	STO	3			
3	Find area		BST	R/S			Area

164 Area of a Triangle $[(x_1, y_1), (x_2, y_2), (x_3, y_3)]$

AREA OF A TRIANGLE $[(x_1, y_1), (x_2, y_2), (x_3, y_3)]$

Given the coordinates of the vertices of a triangle, the area is found by the following formulas:

Area = $\frac{1}{2}$ Determinant of D where

$$D = \begin{bmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{bmatrix}$$

Therefore,

$$\text{Area} = \frac{1}{2} [x_1 (y_2 - y_3) + x_2 (y_3 - y_1) + x_3 (y_1 - y_2)]$$

DISPLAY			KEY ENTRY	DISPLAY			KEY ENTRY	REGISTERS
LINE	CODE			LINE	CODE			
00.				25.	71	x		R ₀
01.	34	RCL		26.	61	+		R ₁ x ₁
02.	01	1		27.	02	2		R ₂ y ₁
03.	34	RCL		28.	81	÷		R ₃ x ₂
04.	04	4		29.	-00	GTO 00		R ₄ y ₂
05.	34	RCL		30.				R ₅ x ₃
06.	06	6		31.				R ₆ y ₃
07.	51	—		32.				R ₇
08.	71	x		33.				R ₈
09.	34	RCL		34.				R ₉
10.	03	3		35.				R ₀₀
11.	34	RCL		36.				R ₀₁
12.	06	6		37.				R ₀₂
13.	34	RCL		38.				R ₀₃
14.	02	2		39.				R ₀₄
15.	51	—		40.				R ₀₅
16.	71	x		41.				R ₀₆
17.	61	+		42.				R ₀₇
18.	34	RCL		43.				R ₀₈
19.	05	5		44.				R ₀₉
20.	34	RCL		45.				
21.	02	2		46.				
22.	34	RCL		47.				
23.	04	4		48.				
24.	51	—		49.				

Example:

Find the area of the triangle with the following x-y coordinate vertices.

(0, 0)

(4, 0)

(4, 3)

Solution:

Area = 6

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Enter coordinates of vertices	x_1	STO	1			
		y_1	STO	2			
		x_2	STO	3			
		y_2	STO	4			
		x_3	STO	5			
		y_3	STO	6			
3	Find area		BST	R/S			Area

AREA OF A POLYGON

If the x-y coordinates of the vertices of a polygon are known, the area can be found by the following formula:

$$\text{Area} = \frac{1}{2} [(x_1 + x_2)(y_1 - y_2) + (x_2 + x_3)(y_2 - y_3) + \dots + (x_n + x_1)(y_n - y_1)]$$

Traverse the coordinates clockwise for a positive area.

DISPLAY		KEY ENTRY	DISPLAY		KEY ENTRY	REGISTERS
LINE	CODE		LINE	CODE		
00.			25.	81	÷	R ₀
01.	33	STO	26.	33	STO	R ₁ x _{i-1}
02.	02	2	27.	61	+	R ₂ y _{i-1}
03.	23	R↓	28.	05	5	R ₃ x _i
04.	33	STO	29.	34	RCL	R ₄ y _i
05.	01	1	30.	04	4	R ₅ ΣAREA
06.	00	0	31.	33	STO	R ₆
07.	33	STO	32.	02	2	R ₇
08.	05	5	33.	34	RCL	R ₈
09.	84	R/S	34.	03	3	R ₉
10.	33	STO	35.	33	STO	R ₀₀
11.	04	4	36.	01	1	R ₀₁
12.	23	R↓	37.	34	RCL	R ₀₂
13.	33	STO	38.	05	5	R ₀₃
14.	03	3	39.	-09	GTO 09	R ₀₄
15.	34	RCL	40.			R ₀₅
16.	01	1	41.			R ₀₆
17.	61	+	42.			R ₀₇
18.	34	RCL	43.			R ₀₈
19.	02	2	44.			R ₀₉
20.	34	RCL	45.			
21.	04	4	46.			
22.	51	-	47.			
23.	71	x	48.			
24.	02	2	49.			


Example:

Find the area of the polygon with the following x-y coordinate vertices.

Point	Coordinates (x, y)
1	(1, 3)
2	(2, 4)
3	(5, 4)
4	(5, 7)
5	(7, 5)
6	(7, 1)
7	(3, 1)

Solution:

Area = 19.50



STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						0.00
2	Input (x ₁ , y ₁)	x ₁ *	↑				
		y ₁ *	BST	R/S			
3	Input (x _i , y _i) for i = 2, ..., n	x _i	↑				Intermediate
		y _i	R/S				
4	Input (x ₁ , y ₁) again	x ₁	↑				Area
		y ₁	R/S				
	*If x ₁ and y ₁ are quite complicated it may be convenient to store them in R ₆ and R ₇ and recall them when needed.						

SPHERICAL TRIANGLE SOLUTION A, b, c

Given two sides and an included angle of a spherical triangle this program finds the other side by the following formula:

$$a = \cos^{-1} (\cos b \cos c + \sin b \sin c \cos A)$$

The program "Spherical Triangle Solution a, b, c" can then be used to find the other angles.

The program works in any angular mode. However, if in degree mode all angles are assumed to be decimal degrees.

DISPLAY		KEY ENTRY	DISPLAY		KEY ENTRY	REGISTERS
LINE	CODE		LINE	CODE		
00.			25.	13	$\cos^{-1}$	R ₀
01.	34	RCL	26.	33	STO	R ₁ a
02.	04	4	27.	01	1	R ₂ b
03.	34	RCL	28.	-00	GTO 00	R ₃ c
04.	03	3	29.			R ₄ A
05.	31	f	30.			R ₅
06.	12	sin	31.			R ₆
07.	31	f	32.			R ₇
08.	00	R←P	33.			R ₈
09.	34	RCL	34.			R ₉
10.	02	2	35.			R ₁₀
11.	31	f	36.			R ₁₁
12.	12	sin	37.			R ₁₂
13.	71	x	38.			R ₁₃
14.	34	RCL	39.			R ₁₄
15.	02	2	40.			R ₁₅
16.	31	f	41.			R ₁₆
17.	13	cos	42.			R ₁₇
18.	34	RCL	43.			R ₁₈
19.	03	3	44.			R ₁₉
20.	31	f	45.			
21.	13	cos	46.			
22.	71	x	47.			
23.	61	+	48.			
24.	32	g	49.			

Example:

Given the following two sides and included angle, find the remaining parameters of a spherical triangle.

$$A = 30^\circ, \quad b = 50.5^\circ, \quad c = 47.3^\circ$$

Solution:

$$a = 22.71^\circ$$

After running the program Spherical Triangle Solution a, b, and c

$$B = 87.88^\circ$$

$$C = 72.13^\circ$$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Store A, b, c	b	STO	2			
		c	STO	3			
		A	STO	4			
3	Find solution		BST	R/S			a
4	To find B and C the program now						
	has a, b, and c in the correct						
	registers to run Spherical						
	Triangle Solution a, b, and c						

SPHERICAL TRIANGLE SOLUTION a, b, c

Given the three sides of a spherical triangle this program calculates the angles by the following formula:

$$A = \cos^{-1} \left(\frac{\cos a - \cos b \cos c}{\sin b \sin c} \right)$$

$$B = \cos^{-1} \left(\frac{\cos b - \cos a \cos c}{\sin a \sin c} \right)$$

$$C = \cos^{-1} \left(\frac{\cos c - \cos a \cos b}{\sin a \sin b} \right)$$

The program works in any angular mode. However, if in degree mode all angles are assumed to be in decimal degrees.

DISPLAY		KEY ENTRY	DISPLAY		KEY ENTRY	REGISTERS
LINE	CODE		LINE	CODE		
00.			25.	32	g	R ₀
01.	34	RCL	26.	13	cos ⁻¹	R ₁ a
02.	01	1	27.	84	R/S	R ₂ b
03.	31	f	28.	34	RCL	R ₃ c
04.	13	cos	29.	01	1	R ₄
05.	34	RCL	30.	34	RCL	R ₅
06.	02	2	31.	02	2	R ₆
07.	31	f	32.	34	RCL	R ₇
08.	13	cos	33.	03	3	R ₈
09.	34	RCL	34.	33	STO	R ₉
10.	03	3	35.	02	2	R ₀₀
11.	31	f	36.	23	R↓	R ₀₁
12.	13	cos	37.	33	STO	R ₀₂
13.	71	x	38.	01	1	R ₀₃
14.	51	—	39.	23	R↓	R ₀₄
15.	34	RCL	40.	33	STO	R ₀₅
16.	02	2	41.	03	3	R ₀₆
17.	31	f	42.	-01	GTO 01	R ₀₇
18.	12	sin	43.			R ₀₈
19.	34	RCL	44.			R ₀₉
20.	03	3	45.			
21.	31	f	46.			
22.	12	sin	47.			
23.	71	x	48.			
24.	81	÷	49.			

Examples:

Given the following three sides of a spherical triangle calculate the three angles.

$$a = 1.12^\circ, \quad b = 52.38^\circ, \quad \text{and } c = 53.42^\circ$$

Solution:

$$A = .52^\circ, \quad B = 21.63^\circ, \quad C = 158.05^\circ$$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Store a, b, and c	a	STO	1			
		b	STO	2			
		c	STO	3			
3	Solve for A, B, and C		BST	R/S			A
			R/S				B
			R/S				C

SPHERICAL TRIANGLE SOLUTION A, B, C

Given the three angles of a spherical triangle this program calculates the sides by the following formulas:

$$a = \cos^{-1} \left(\frac{\cos A + \cos B \cos C}{\sin B \sin C} \right)$$

$$b = \cos^{-1} \left(\frac{\cos B + \cos A \cos C}{\sin A \sin C} \right)$$

$$c = \cos^{-1} \left(\frac{\cos C + \cos A \cos B}{\sin A \sin B} \right)$$

The program works in any angular mode. However, if in degree mode all angles are assumed to be in decimal degrees.

DISPLAY			KEY ENTRY	DISPLAY			KEY ENTRY	REGISTERS
LINE	CODE			LINE	CODE			
00.				25.	32	g		R ₀
01.	34	RCL		26.	13	cos ⁻¹		R ₁ A
02.	01	1		27.	84	R/S		R ₂ B
03.	31	f		28.	34	RCL		R ₃ C
04.	13	cos		29.	03	3		R ₄
05.	34	RCL		30.	34	RCL		R ₅
06.	02	2		31.	02	2		R ₆
07.	31	f		32.	34	RCL		R ₇
08.	13	cos		33.	01	1		R ₈
09.	34	RCL		34.	33	STO		R ₉
10.	03	3		35.	03	3		R ₀₀
11.	31	f		36.	23	R↓		R ₀₁
12.	13	cos		37.	33	STO		R ₀₂
13.	71	x		38.	01	1		R ₀₃
14.	61	+		39.	23	R↓		R ₀₄
15.	34	RCL		40.	33	STO		R ₀₅
16.	02	2		41.	02	2		R ₀₆
17.	31	f		42.	-01	GTO 01		R ₀₇
18.	12	sin		43.				R ₀₈
19.	34	RCL		44.				R ₀₉
20.	03	3		45.				
21.	31	f		46.				
22.	12	sin		47.				
23.	71	x		48.				
24.	81	÷		49.				

Example:

Given the following three angles of a spherical triangle find the three sides.

$A = .52^\circ, \quad B = 21.63^\circ, \quad C = 158.05^\circ$

Solution:

$a = 1.10^\circ, \quad b = 51.51^\circ, \quad c = 52.53^\circ$

STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Store A, B, and C	A	STO	1			
		B	STO	2			
		C	STO	3			
3	Calculate a, b, and c		BST	R/S			a
			R/S				b
			R/S				c

TRANSLATION AND/OR ROTATION OF COORDINATE AXIS

Let (x, y) be coordinates in the old system and let (x_0, y_0) be the center of a new coordinate system rotated through an angle of θ . The new coordinates are (x', y') and are calculated by the following formulas:

$$1. \quad x' = (x - x_0) \cos \theta + (y - y_0) \sin \theta$$

$$2. \quad y' = -(x - x_0) \sin \theta + (y - y_0) \cos \theta$$

For no rotation put in $\theta = 0$.

For no translation put in $(x_0, y_0) = (0, 0)$

DISPLAY		KEY ENTRY	DISPLAY		KEY ENTRY	REGISTERS
LINE	CODE		LINE	CODE		
00.			25.	51	—	R ₀
01.	34	RCL	26.	22	$x \leftrightarrow y$	R ₁ θ
02.	02	2	27.	34	RCL	R ₂ x_0
03.	51	—	28.	04	4	R ₃ y_0
04.	34	RCL	29.	61	+	R ₄ $(x - x_0) \cos \theta$
05.	01	1	30.	-00	GTO 00	R ₅ $(x - x_0) \sin \theta$
06.	22	$x \leftrightarrow y$	31.			R ₆
07.	31	f	32.			R ₇
08.	00	R←P	33.			R ₈
09.	33	STO	34.			R ₉
10.	04	4	35.			R ₁₀
11.	23	R↓	36.			R ₁₁
12.	33	STO	37.			R ₁₂
13.	05	5	38.			R ₁₃
14.	23	R↓	39.			R ₁₄
15.	34	RCL	40.			R ₁₅
16.	03	3	41.			R ₁₆
17.	51	—	42.			R ₁₇
18.	34	RCL	43.			R ₁₈
19.	01	1	44.			R ₁₉
20.	22	$x \leftrightarrow y$	45.			
21.	31	f	46.			
22.	00	R←P	47.			
23.	34	RCL	48.			
24.	05	5	49.			

Examples:

1. Translate the point $(1, 1)$ to a new coordinate system with center at $(1, 2)$.
($\theta = 0$)
2. Translate the point $(1, 3)$ to a new coordinate system with center at $(-1, 4)$ at 30° to the old system.

Solutions:

1. $(0, -1)$
2. $(1.23, -1.87)$


STEP	INSTRUCTIONS	INPUT DATA/UNITS	KEYS				OUTPUT DATA/UNITS
1	Enter program						
2	Store constants	θ	STO	1			
		x_0	STO	2			
		y_0	STO	3			
3	Enter coordinates	y	$\uparrow$				
		x	BST	R/S			x'
			$x \leftrightarrow y$				y'
4	For a new point go to step 3.						

INDEX

- Accumulated Interest 54
- Angles 147, 150, 152, 154, 156
- Angles Between Vectors 96
- Arc Cosecant 132
- Arc Cotangent 132
- Arc Secant 132
- Area 158, 160, 162, 164, 166
- Axis 174
- Base Conversion 122, 124
- Circle 138, 140, 142, 144
- Circle Determined by Three Points 142
- Complex Arithmetic 6
- Complex Functions 9, 12, 14, 16
- Complex Hyperbolic 24, 26, 28
- Complex Inverse Hyperbolic 36, 38, 40
- Complex Inverse Trigonometric 30, 32, 34
- Complex Polynomial Evaluation 42
- Complex Trigonometric 18, 20, 22
- Compounded Amount 44
- Coordinate Axis 174
- Cosecant 130
- Cosh 134
- Cosine Integral 102
- Cotangent 130
- Coversine 132
- Cross Product 86
- Day of the Week 75
- Days Between Dates 75
- Depreciation 68, 70, 72
- Determinant 78, 80
- Differential Equations 110
- Discounted Cash Flow Analysis 66
- Division, Synthetic 116
- Dot Product of Vectors 96
- Equally Spaced Points on a Circle 144
- Equations 88, 91, 114, 126
- Exponential Integral 104
- Exsecant 132
- Factoring Integers 118
- Function Solution 126
- Haversine 132
- Hyperbolic Functions 134
- Integers 118, 122, 124
- Integration 108, 110
- Interest 44, 48, 54, 57, 60
- Interpolation 112
- Inverse of a Matrix 78, 82
- Iterative Solutions to Equations 126
- Linear Interpolation 112
- Matrix 78, 80, 82, 94
- Newton's Method 126
- Norm of Vectors 96
- Numerical Integration 106, 108
- Numerical Solution to Differential Equations 110
- Payment 44, 51, 54, 63
- Polygons 138, 140
- Polynomial Evaluation 120
- Present Value 51
- Primes 118
- Quadratic Equations 114
- Rotation 174
- Secant 130
- Simpson's Rule 108
- Simultaneous Equations 88, 91
- Sine Integral 100
- Sinh 134
- Spherical Triangles 168, 170, 172
- Synthetic Division 116
- Tanh 134
- Time Periods 44, 48, 51, 54, 63
- Translation 174
- Trapezoidal Rule 106
- Triangle Area 158, 160, 162, 164
- Triangle Solution 147, 150, 152, 154, 156
- Trigonometric Functions 130, 132
- Vector Cross Product 96
- Vectors 96, 86
- Versine 132
- Weekday 75



Sales and service from 172 offices in 65 countries.
19310 Pruneridge Avenue, Cupertino, California 95014



Scan Copyright ©
The Museum of HP Calculators
www.hpmuseum.org

Original content used with permission.

Thank you for supporting the Museum of HP
Calculators by purchasing this Scan!

Please to not make copies of this scan or
make it available on file sharing services.